



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

### Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

### About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>





2044 106 408 578

P232i2

c.1

v.1

W. G. FARLOW

P232i2 c.1 v.1

Harvard University



FARLOW  
REFERENCE LIBRARY  
OF  
CRYPTOGAMIC BOTANY















# INDEX BRYOLOGICUS

SIVE

## ENUMERATIO MUSCORUM

AD DIEM ULTIMAM ANNI 1900 COGNITORUM

ADJUNCTIS

*Synonymia distributioneque geographica*  
LOCUPLETISSIMIS

QUEM CONSCRIPSIT

**E. G. PARIS, O. O.**

---

**Editio secunda .**

**Vol. I.**

---

PARIS V<sup>me</sup>

LIBRAIRIE SCIENTIFIQUE A. HERMANN

6 et 12, Rue de la Sorbonne.

---

1904.

Handwritten text, possibly a signature or initials, located in the top left corner.

**DIES**  
**PROMULGATIONIS FASCICULORUM**  
**I-VI**

<b>Fasciculus</b>	<b>I (p. 1-64).....</b>	<b>1<sup>a</sup> Décembri 1903.</b>
—	<b>II (p. 65-128).....</b>	<b>2<sup>a</sup> Januar. 1904.</b>
—	<b>III (p. 129-192).....</b>	<b>5<sup>a</sup> Februar. 1904.</b>
—	<b>IV (p. 193-256).....</b>	<b>2<sup>a</sup> Mart. 1904.</b>
—	<b>V (p. 257-320).....</b>	<b>1<sup>a</sup> April. 1904.</b>
—	<b>VI (p. 321-3 ).....</b>	<b>1<sup>a</sup> Magi. 1904.</b>





# INDEX BRYOLOGICUS.

---

**ABIETINELLA** C. M. in *Nuov. Giorn. bot. ital.* 1896 (nom.)  
Spec. omn. sunt *THUIDIA* ejusd. nom.

**Acamptodous** Duby *Choix de Crypt. ex.* 1880, p. 9 (sect. *vel* subgen. *LEPIDOPILI*).

**ACANTHOCLADIUM** Mitt. *Austral. M.* p. 37 (1878). — 18 spec.

*Armittii* Broth. et Geh. — *Mastopoma*.

*Bessoni* (*Ren. et Card.*) Broth. in litt.

*Microthamnium Bessoni* R. C. in *Bull. Soc. roy. bot. Belg.* 1893, I, p. 123;  
*Ren. M. Masc. Mad.* n. 23; *Par. Ind.* p. 807 (1896).

*Sematophyllum Bessoni* eor. in *Ren. Prodr. fl. bryol. Mad. etc.* p. 234  
(1897); *Par. Suppl. Ind.* p. 306 (1900).

Ster. — *Afr.* : Madagascar.

*Crossii* Broth. et Geh. in *Oesv af Finska Vet.-Akad. Foerh.*  
1893, p. 55.

2. — *Pac.* : Austral. (Manning River).

*extenuatum* (*Brid.*) Mitt. in *Trans. and Proceed. of the roy. Soc. of Victoria*, 1883, p. 85.

*Hypnum crinitum* H. f. et W. *Fl. of the N. Zeal.* II, p. 114, t. 91, f. 1 (1855).  
*H. extenuatum* *Brid. Bryol. univ.* II, p. 484 (1827); *C. M. Syn.* II,

p. 391 p.p. (1851); *H. f. et W. Fl. Tasman.* II, p. 314 (1860) et *Handb. of the N. Zeal. Fl.* p. 481 (1867); *Par. Ind.* p. 635 (1895).

*H. glaucescens* Hsch. in *Sieber M. Nov. Holland.*

*Rhynchostegium glaucescens* Mitt. in op. cit. p. 80.

*Sematophyllum extenuatum* *Par. Suppl. Ind.* p. 306 (1900).

2. Cortic. — *Pac.* : Austral. or, Nov. Seeland., Tasman., ins. Auckland.

*flexile* (*Ren. et Card.*) Broth. in litt.

*Hypnum alamarzartrense* Kiaër?

*Microthamnium flexile* R. C. in *Bull. Soc. roy. bot. Belg.* 1890, I, p. 185;  
*Ren. M. Masc. Mad.* n. 18; *Par. Ind.* p. 808 (1896).

*Sematophyllum flexile* eor. in *Ren. Prodr. fl. bryol. Mad. etc.* p. 325 (1897);  
*Par. Suppl. Ind.* p. 307.

Ster. — *Afr.* : Madagascar.

*hamatum* C. M. — *Mastopoma*.

*Jungneri* Broth. in *Engler's Bot. Jahrb.* 1897, p. 278.

*Sematophyllum Jungneri* *Par. Ind.* p. 1166 (1897).

1. Cortic. — *Afr.* : Cameroon.

*lanceifolium* (*Harv.*) Broth. in litt.

*Hypnum Korthalsii* Dz. *Mk. in Ann. sc. nat.* 1844, p. 307 et *M. fr. Arch.*  
*ind.* p. 11 (1844); *C. M. Syn.* II, p. 497 (1851).

- H. lancifolium* C. M. op. et l. cc.; *Bryol. jav. II*, p. 141, t. 239 (1861-70).  
*H. trimegustum* Mont. in *Lond. Journ. of bot.* 1844, p. 633 et in *Ann. sc. nat.* 1845, p. 87; *Syll.* p. 10 (1856).  
*Neckera lancifolia* Harv. in *Lond. Journ. of bot.* 1840, p. 14 et in *Hook. Ic. pl. rar.* t. 21, f. 5 (1841).  
*Sematophyllum lancifolium* Jaeg. *Ad. II*, p. 445 (1875-76); *Par. Ind.* p. 1166 (1897).  
*Stereodon lancifolius* Mitt. *M. Ind. or.* p. 110 (1859).
2. — As. : Nepal., Sumatra, Java, Borneo.
- pedunculatum* Mitt. in *Proced. of the Linn. Soc. of N. S. Wales*, 1882, I, p. 102.  
*Hypnum pedunculatum* Par. *Ind.* p. 666 (1895).  
*Sematophyllum pedunculatum* ej. *Suppl. Ind.* p. 307 (1900).
2. fr. ign. — Pac. : Polynes., Nov. Hebrid. (Aneitum).
- Prionodontella* Broth. in *Warburg, Monsonia*, I, p. 177 (1899).  
*Sematophyllum Prionodontella* Par. *Suppl. Ind.* p. 307 (1900).
2. fr. ign. — As. : Mindanao.
- pseudo-tanytrichum* (*Bryol. jav.*) Broth. in litt.  
*Hypnum pseudo-tanytrichum* *Bryol. jav. II*, p. 150, t. 248 (1861-70); *Par. Ind.* p. 672 (1896).  
*H. Hornschuchii* Dz. et Mh. in *Ann. sc. nat.* 1844, p. 307.
2. Silv. — As. : M. Java.
- rigidicaule* (C. M.) Broth. in *Engler's Bot. Jahrb.* 1897, p. 278.  
*Heterophyllum rigidicaule* C. M. in *Dusen M. Camer.* n. 794.  
*Rhaphidostegium rigidicaule* Par. *Ind.* p. 1103 (1897).
2. Cortic. — Afr. : Cameroon.
- rigidum* (Hsch. et Reinw.) Broth. in *Warburg, Monsonia*, I, p. 51 (1899).  
*Hypnum rigidum* H. et R. in *Nov. Act. Acad. Caes. Leop. XIV, II, Suppl.* p. 731 (1829); *Bryol. jav. II*, p. 140, t. 237 et 238 (1861-70); C. M. *Syn. II*, p. 498 (1851).  
*Sematophyllum rigidum* Mitt. *Fl. Vit.* p. 398 (1871); *Par. Ind.* p. 1169 (1897).
2. Cortic. — As. : Sumatra, M. Java, Borneo. Pac. : Nov. Caledon., Nov. Guin.
- » *var. Braunianum* (*Bryol. jav.*) Par.  
*Hypnum rigidum var. Braunianum* *Bryol. jav. l. c.*  
*Sematophyllum rigidum var. Braunianum* Jaeg. *Ad. II*, p. 445 (1875-76); *Par. op. et l. cc.*
- As. : Java.
- » *var. convolutum* (*Bryol. jav.*) Par.  
*Hypnum rigidum var. convolutum* *Bryol. jav. l. c.*  
*Sematophyllum rigidum var. convolutum* Jaeg. et Par. *op. et l. cc.*
- As. : — Java.
- Rutenbergii* (C. M.) Broth. in litt.  
 ? *Hylocomium Rutenbergii* Par. *Ind.* p. 591 (1895).  
*Hypnum Rutenbergii* C. M. in *Abhandl. ... Brem. VII*, p. 213 (1881).  
*Sematophyllum Rutenbergii* Ren. Pr. *fl. bryol. Madag. etc.*, p. 234 (1897).
- Afr. : Madagascar.



*Strangei* Mitt. et F. v. Müll. in *Proced. of the Linn. Soc. of N. S. Wales*, 1882, I, p. 102.

Hypnum *Strangei* Par. Ind. p. 687 (1896).

Sematophyllum *Strangei* ej. *Suppl. Ind.* p. 308 (1900).

— Pac. : Nov. Caled. (ins. Pinorum).

subtrichocolea (C. M.) Broth. in litt.

Sematophyllum subtrichocolea Par. Ind. p. 1171 (1897).

Trismegistia subtrichocolea C. M. in *Dusen M. Camer.* n. 30.

— Afr. : Cameroon.

surculare (Mitt.) Broth. in litt.

Hypnum surculare Jaeg. Ad. II, p. 608 (1877-78); Par. Ind. p. 690 (1896).

Stereodon surcularis Mitt. M. Ind. or. p. 112 (1859).

2 ?

— As. : Khasia.

tanytrichum (Mont) Broth. in litt.

Hypnum tanytrichum Mont. in *Ann. sc. nat.* 1845, IV, p. 88 et *Syll.* p. 9 (1856); C. M. *Syn.* II, p. 391 (1851); *Bryol. jav.* II, p. 151, t. 249 (1861-70); Par. Ind. p. 690 (1896).

Sematophyllum tanytrichum Par. *Suppl. Ind.* p. 308 (1900).

Stereodon tanytrichus Mitt. op. et l. cc.

2.

— As. : Bhotan, Sikkim, Java.

Trichocolea (C. M.) Broth. in litt.

Hypnum Trichocolea C. M. in *Fl.* 1888, p. 417.

Sematophyllum Trichocolea Par. Ind. p. 1171 (1897).

Lignic. — Afr. : Kilima N'Djaro, Usambara.

trichocoleoides (C. M.) Broth. in litt.

Hypnum trichocoleoides C. M. in *Fl.* 1886, p. 283.

Sematophyllum trichocoleoides Par. Ind. p. 1171 (1897).

Ster.

— Afr. : Cameroon, ins. Sao Thoma.

uncinifolium Broth. — Mastopoma.

ACANTHODIUM Mitt. in *Journ. of the Linn. Soc.* 1868, p. 182.

*deflexifolium* Mitt. — Sematophyllum.

*papillatum* Mitt. — Trichosteleum.

*rigidum* Mitt. — Sematophyllum complanatum ?

ACAULON C. M. in *Bot. Zeit.* 1847, p. 99, *Syn. I*, p. 21 (1849) et *II*, p. 519 (1851).

Spec. omn. non indicatae sunt SPHÆRANGIA ejusd. nom.

*carniolicum* C. M. — Phascum.

*Floërkeanum* C. M. — Microbryum.

*Floërkeanum* var. β C. M. — Microbryum *Floërkeanum* var. *badium*.

*minus* Jaeg. — Sphærangium muticum var. *minus*.

*muticum* C. M. } — Sphærangium muticum p.p.

                          } — Sphærangium capense p.p.

*pellucidum* Fleisch. — Sphærangium mediterraneum.

*piligerum* Limpr. — Sphærangium muticum var. *cuspidatum*.

*triquetrum* var. Sull. et Lesq. — Sphærangium rufescens.

*turgidum* Mitt. — Sphærangium apiculatum var. *turgidum*.

Achyrophyllum Mitt. in *Journ. of the Linn. Soc.* 1859, p. 88 (sect. STERIODONTIS = PTYCHOMNION).

**Achrolepis** Lindb. in *Act. Soc. sc. Fenn.*, X, 1872, p. 276 (sect. MYURELLÆ, fors. gen. propr.)

ACIDODONTIUM *Schwaegr. Suppl. II, P. II, p. 152* (1827). — 14 spec.

brachypodium (C. M.) *Broth. Natur. Pflanzenfam. p. 561.*

Brachymenium brachypodium *Par. Suppl. Ind. p. 38* (1900).

Bryum brachypodium C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 22.

2. — *Am. merid.* : And. Boliv.

exaltatum (*Spruce*) *Jaeg. Ad. I, p. 572* (1873-74).

Bryum exaltatum *Spr. in Mitt. M. austr. am. p. 285* (1869); *Par. Ind. p. 181* (1894).

2. Cortic. — *Am. merid.* : And. Quitens.

floresianum C. M. in *Bull. Soc. roy. bot. Belg.* 1892, I, p. 162.

Bryum floresianum *Par. op. cit. p. 183.*

2. — *Am. sept.* : And. Costa Ricens.

heteroneuron (*Spruce*) *Broth. op. et l. cc.*

Brachymenium heteroneuron *Jaeg. op. cit. p. 581*; *Par. op. cit. p. 122.*

Bryum heteroneuron *Spr. in Mitt. op. cit. p. 281.*

2. Cortic. — *Am. merid.* : And. Quitens.

*Kunthii* } Hsch. — Brachymenium systylium.  
              } *Schwaegr.* — A. megalocarpum.

lonchotrachelon (C. M.) *Broth. op. et l. cc.*

Brachymenium lonchotrachelon *Par. Suppl. Ind. l. cc.*

Bryum lonchotrachelon C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 23.

2. — *Am. merid.* : And. Boliv.

longifolium (W. P. Sch.) *Broth. op. et l. cc.*

Brachymenium longifolium W. P. Sch. in *Mandon Pl. Boliv. n. 1662*;  
*Par. Ind. p. 123.*

Bryum longifolium C. M. in *op. et l. cc.*

2. — *Am. merid.* : And. Boliv.

macropoma (C. M.) *Broth. op. et l. cc.*

Brachymenium macropoma *Par. Ind. p. 124* (1894).

Bryum macropoma C. M. in *Linn.* 1878-79, p. 290.

2. Cortic. — *Am. merid.* : Argentin, subtropic.

megalocarpum (*Hook.*) *Ren. et Card.*

A. *Kunthii* *Schwaegr. Suppl. II, P. II, p. 152, t. 196* (1827).

Bryum megalocarpum *Hook. in Kunth. Syn. pl. Equinoct. p. 59* (1822-28);

C. M. *Syn. I, p. 264* (1849); *Mitt. op. cit. p. 286*; *Par. Ind. p. 194.*

Megalangium *Kunthii* *Brid. Bryol. univ. II, p. 29, t. Suppl. 3* (in ic. sub *Macrothecio*) (1827).

2. Cortic. — *Am. merid.* : And. Nov. Granat. et Quitens.

ramicola (*Spruce*) *Jaeg. op. et l. cc.*

Bryum ramicola *Spr. in Mitt. op. cit. p. 286*; *Par. Ind. p. 209.*

2. Cortic. — *Am. merid.* : And. Quitens.

rhamphostegium (*Hpe.*) *Jaeg. op. cit. p. 571.*

Bryum rhamphostegium *Par. Ind. p. 210.*

Cladodium rhamphostegium *Hpe. in Ann. sc. nat. V ser., IV, p. 339* (1866-67).

2. Cortic ? — *Am. merid.* : And. Nov. Granat.

seminerve *Hook. f. et Wils. in Lond. Journ. of bot.* 1846, p. 451, t. 164.

*Bryum seminerve* C. M. *Syn.* I, p. 264 (1849); *Mitt. op. cit.* p. 285; *Mitt. op. cit.* p. 285; *Par. Ind.* p. 214.

2. Cortic. — *Am. sept.* : *Jamaic. Am. merid.* : And. Quitens.

*Sprucei* (*Mitt.*) *Jaeg. op. cit.* p. 572.

*Bryum Sprucei* *Mitt. op. cit.* p. 285; *Par. Ind.* p. 215.

2. Cortic ? — *Am. merid.* : And. Quitens.

subrotundum *Hook. f. et Wils. op. et l. cc.*

*Brachymerium subrotundum* *Tayl. in eod. op.* p. 56.

*Bryum subrotundum* *Mitt. op. cit.* p. 287.

*Bryum Taylori* C. M. *Syn.* I, p. 264 (1849); *Par. Ind.* p. 218.

2. Cortic. — *Am. merid.* : And. Quitens.

trachyticola (C. M.) *Broth. op. et l. cc.*

*Brachymerium trachyticola* *Jaeg. op. cit.* p. 582; *Par. Ind.* p. 127.

*Bryum trachyticola* C. M. in *Linn.* 1874, p. 583.

2. Rupic. — *Am. merid.* : And. Quitens.

ACIRHYNCHUM *Hpe. in sched.*

*cupressinum* *Hpe.* — *Rhaphidostegium sphærotheca.*

**Acisphagnum** C. M. in *Linn.* 1874, p. 548 (nom.) et in *Fl.* 1887, p. 404  
(*Sphagna cuspidata*).

**Accosphagnum** C. M. in *Fl.* 1887, p. 405 (*Sphagna sericea*).

**Acoleos** C. M. in *Linn.* 1874, p. 509 (sect. *BARTRAMIÆ*).

ACOSTA C. M. in *Linn.* 1848, p. 91.

*cuspidata* C. M. — *Leucomium cuspidatifolium.*

*Meyeniana* C. M. — *Ectropothecium.*

*Suartzii* C. M. — *Ectropothecium subdenticulatum.*

ACROCERATIUM *Mitt. in Trans. and Proced. of the roy. Soc. of Victoria,*  
1883, p. 36 (erratum ! = *ACROCLADIUM*).

**ACROCLADIUM** *Mitt. M. Austr. Am.* pp. 22 et 531 (1869). — 4 spec.

*Auberti* (*Schwaegr.*) *Jaeg. Ad. II,* p. 509 (1876-77).

*Hypnum Auberti* *Schwaegr. Suppl. I, P. II,* p. 200 (1816); C. M. *Syn. II,*  
p. 262 (1851).

*H. Stereodon Auberti* *Brid. Bryol. univ. II,* p. 50 (1827).

*Leskea nitens* *Brid. Sp. M. II,* p. 556 (1812).

2. — *Afr.* : *Ins. Borbon., Franciaë, Comor. magn.,*  
*Madagascar.*

*auriculatum* *Mitt. p.p.* \ — *Lembophyllum.*

*cuspidatum* *Lindb.* — *Hypnum.*

*cuspidatum* *Lindb.* — *Hypnum.*

*mucronatum* (C. M.) *Jaeg. op. et l. cc.*

*Hypnum mucronatum* C. M. *op. et l. cc.*

2. — *Afr.* : *Pr. B. Sp.*



*Phyllogonium* Jaeg. — *A. politum*.

*politum* (Hook. f. et Wils.) Mitt. *M. austr. am.* p. 531 (1869).

*A. Phyllogonium* Jaeg. op. et l. cc.

*Hypnum Phyllogonium* C. M. Syn. II, p. 262 (fid. Mitt.).

*H. politum* H. f. et W. in *Lond. Journ. of bot.* 1844, p. 353 et *Fl. Antarct. II*, p. 416, t. 154, f. 2 (1847); *C. M. op. cit.* p. 263.

*Phyllogonium callichroum* Mont. in *Ann. sc. nat.* 1845, IV, p. 87.

2. *Lignic.*, *rupic.* et ad terr. humos. — *Afr.* : Pr. B. Sp., ins. Kerguelen, Marion. *Am. merid.* : And. Nov. Granat. et Quitens., Chile, Fuegia, ins. Eremitae. *Pac.* : Austral., Nov. Seeland., Tasman.

*trichocladium* Bosw. in *Journ. of bot.* 1892, p. 99.

*Ster.* — *Pac.* : Austral.

**ACROCRYPHÆA**<sup>(1)</sup> *Bryol. eur. V. Mont. Cryphaeae.* ( ) 10 spec.

*brachyodus* Broth. in litt.

*A. brevidens* Broth. in *Mosen M. Brasil* n. 80.

— *Am. merid.* : Brasil or. (Minas Geraes).

*caripensis* Hpe. — *A. squarrosula*.

*Coffeae* (C. M.) Par. *Suppl. Ind.* p. I (1900).

*Cryphaea Coffeae* C. M. in *Hedwig.* 1888, p. 240.

*Cr. filiformis* Sull. in *Wright M. cubens.* n. 67.

*Cr. Gardneri* Mitt. *M. austr. am.* p. 416 p.p. (fid. C. M.).

Ad *Coffeam arabicam*. — *Am. sept.* : Cuba.

*concavifolia* (Griff.) *Bryol. jav.* II, p. 196 (1861-70).

*A. javanica* *Bryol. eur.* l. c.

*Cryphaea concavifolia* Mitt. *M. Ind. or.* p. 125 (1859).

*Orthotrichum concavifolium* Griff. *Not.* p. 400 et *l.c. pl. asiat. II*, t. 76, fig. 3 (1841).

*Schœnobryum julaceum* Dz. et Mk. *M. frond. ined. Archip. ind.* p. 184 t. 60 (1854).

— *As.* : Nepal., Ceylan, Java, Celebes.

*corymbosa* β Hpe. — *A. squarrosula*.

*corymbosula* Par. et W. P. Sch. *Mon. Cryph. ined. et in hb. W. P. Sch. (nunc in hb. Kew.)*.

— *Am. sept.* : Mexic.

*ferruginea* (Lindb.) Par. et W. P. Sch. *ll. cc.*

*A. Gardneri* Jaeg. *Ad. II*, p. 94 (1874-75).

*Cryphaea ferruginea* Lindb. *mem. ab Aongstr. in Oefv. af K. Vet.-Akad. Foerh.* 1876, p. 81.

*Cr. Gardneri* Mitt. in *Journ. of the Linn. Soc.* 1859 et *M. austr. am.* p. 415 (1869) p. p.

*Grimmia julacea* Hsch. *Fl. bras.* p. 7, t. I, f. I, p. p. (1840).

— *Am. sept.* : Cuba. *Am. merid.* : And. Nov.

Granat, Brasil. austr. or.

*Gardneri* Jaeg. — *A. ferruginea*.

*javanica* *Bryol. eur.* — *A. concavifolia*.

(1) Spec. omn. monoicae et, ut videtur, corticolae.

julacea (*Hsch.*) *W. P. Sch. ll. cc.*

*Cryphaea julacea* *Mitt. M. austr. am. p. 415 (1869).*

*Cr. maynensis* *Tayl. in Spr. M. Amaz. n. 1279,*

*Grimmia julacea* *Hsch. op. et l. cc. p.p.*

*Pilotrichum julaceum* *C. M. Syn. II, p. 173 p.p. (1851).*

— *Am. merid.* : Columb., And. Quitens. et  
Peruv., Brasil. austr. or., Paraguay.

« *var. costa-ricensis* *Ren. et Card. in Bull. Soc. roy. bot. Belg, 1894,*  
*I, p. 175.*

— *Am. sept.* : Costa-Rica.

mexicana *W. P. Sch. ll. cc.*

*Cryphaea Leiboldii* *C. M. in Linn. 1844, p. 680.*

*Pilotrichum julaceum* *C. M. op. et l. cc. p.p.*

— *Am. sept.* : Mexic.

paraguensis *Besch. in Rev. bryol. 1885, p. 18 (nom.) et in Journ. de*  
*bot. 1891, p.*

— *Am. merid.* : Paraguay.

rubricaulis (*Mitt.*) *Jaeg. Ad. II, p. 95 (1874-75).*

*Cryphaea rubricaulis* *Mitt. M. austr. am. p. 415 (1869).*

— *Am. merid.* : And. Quitens., Brasil centr.

squarrosula (*C. M.*) *W. P. Sch. ll. cc.*

*A. caripensis* *Hpe. Enum. M. Brasil. p. 38 (1879).*

*A. corymbosa*  $\beta$  *Hpe. in Krause M. Ecuador.*

*Cryphaea caripensis* *C. M. in Linn. 1847, p. 210.*

*Cr. filiformis* *C. M. in op. et l. cc.*

*Moritzia caripensis* *Hpe. Enum. M. Brasil. p. 82.*

*Pilotrichum filiforme*  $\beta$  *squarrosulum* *C. M. Syn. II, p. 169 (1851).*

— *Am. sept.* : Mexic. *Am. merid.* : Venezuela.

ACROPORIUM *Mitt. in Journ. of the Linn. Soc. 1868, p. 182 (gen. propr.)*  
*et M. austr. am. p. 477 (1869) (sect. SEMATOPHYLLI).*

*Spec. omn. non indicatae sunt SEMATOPHYLLA ejusd. nom.*

*lumprophyllum* *Mitt. — S. Mittenii.*

**ACROSCHISMA** *Hook. f. et Wils. in Lond. Journ. of bot. 1844, pp. 536.*  
*et 538. — 2 spec.*

*andensis* *Spr. — A. Wilsoni.*

*densifolia* *Wils. in Kew Journ. bot. IX, p. 289 (1857).*

*Andreaea densifolia* *Mitt. M. Ind. or. p. 7 (1859).*

— *As.* : Sikkim.

*Wilsoni* *Hook f. et Wils. in op. et ll. cc., et Fl. antarct II, p. 379,*  
*t. 151, f. 3 (1844).*

*A. andensis* *Spr. in Proceed. of the Linn. Soc.*

*Andreaea Wilsoni* *C. M. Syn. I, p. 11 (1849) et II, p. 517 (1851).*

1. *Rupic?* — *Am. merid.* : And. Peruv., Patag., Fuegia, ins. Eremitae.  
*Pac.* : Ins. Auckland, Campbell.

**Acrosphagnum** *C. M. in Fl. 1887, p. 405 (Sphagna mucronata).*

**ACTINODONTIUM** Schwaegr. Suppl. II, P. II, I, p. 175, t. 174 (1826).  
6 spec.

*ascendens* Schwaegr. *op. et l. cc.* : *Brid. Bryol. univ. II*, p. 736 (1827) ;  
*C. M. Syn. II*, p. 74 (1851) ; *Bryol. jav. II*, p. 36, t. 159 (1861-70).

1 ou Syn.

— *As.* : M. Java.

*Duseni* (C. M.) *Broth. in Engler's Bot. Jahrb.* 1897, p. 260.

*Lepidopilum Duseni* C. M. in *Dusen M. Camer. n.* 40 ; *Broth. in op. cit.*  
1894, p. 201 (descr.).

*Lepidopilum integrum* C. M. in *Coll. cit.*

1. Cortic. — *Afr.* : Cameroon.

*hirsutum* Besch. *Fl. bryol. Réunion, etc.*, p. 137 (1880-81).

1. Ad arb. — *Afr.* : Ins. Bourbon.

« *var. ramosum* Besch. *op. et l. cc.*

Ubi typ.

*rhapidostegum* (C. M.) *Bryol. jav. II*, p. 37, t. 160 (1861-70).

*Hookeria rhapidostega* C. M. *Syn. II*, p. 677 (1851).

1 (C. M.), 2 (*Bryol. jav.*). — *As.* : Java.

*Sprucei* (Mitt.) *Jaeg. Ad. II*, p. 229 (1874-75).

*Lepidopilum Sprucei* Mitt. *M. austr. am. p.* 370 (1869).

2. Silv. — *Am. merid.* : And. Quitens.

*streptopogonaceum* (C. M.) *Broth. in op. et l. cc.*

*Lepidopilum streptopogonaceum* C. M. in *Dusen M. Camer. n.* 107.

1 et Syn. Cortic. — *Afr.* : Cameroon.

**Acuminella** C. M. *Syn I*, p. 752 (sect. SCHLOTHEMIÆ) (1849).

**ADELOTHECIUM** Mitt. *M. austr. am. p.p.* 18 et 301 (1869).

*bogotense* } Hpe. — *Distichophyllum Geheebii*.  
                  } Mitt. — *Hookeria*.

**Adelphodon** Lindb. in *Act. Soc. sc. Fenn.* 1872, p. 245 (sect. LESQUEREUXIÆ).

**AEROBRYUM** Dz. et Mk. in *Nederl. Kruidk. Arch.* II, p. 279 ( ) .  
24 spec.

*Bauerae* C. M. *ms.*

— *Pac.* : Austral.

« *var. gracilis* C. M. in *Geheeb Neue Beitr. zur Moosfl. v. Neu-Guin.* p. 11 (1889).

— *Pac.* : Nov. Guin. sept.

*capense* (W. P. Sch.) C. M. in *Linn.* 1876, p. 262.

*Meteorium capense* W. P. Sch. in *Breut. M. capens.*

*Neckera capensis* C. M. in *Bot. Zeit.* 1858, p. 165.

2 ?

— *Afr.* : Pr. B. Sp.

« *var. rupestris* C. M. in *Rehm. M. Afr. austr. n.* 322 b.

Ut et ubi typ.

- capillicaule* Ren. et Card. in Bull. Soc. roy. bot. Belg. 1898, I, p. 120.  
Ster. — Afr. : Madagascar.
- Conferva* C. M. in Linn. 1878-79, p. 492.  
— Am. merid. : Venezuela.
- crispicuspe* Besch. in Journ. de bot. 1891, p.  
— Afr. : Ins. Seychell.
- cymatocheilos* (C. M.) Par. Ind. p. 9 (1894).  
Eriocladium cymatocheilos C. M. in Fl. 1890, p. 493.  
Ster. Silv. — Afr. : Kilima N'Djaro.
- hokinense* Besch. in Ann. sc. nat. 1892, I, p. 74.  
2. fr. ign. Silv. — As. : Yunnan.
- integrifolium* Besch. in op. et l. cc.  
2. fr. ign. Silv. — As. : Yunnan.
- javanicum* (Bryol. jav.) C. M. in Linn. 1876, p. 262.  
Meteorium javanicum Bryol. jav. II, p. 87, t. 203 (1861-70).  
2. Ad ramul. arb. — As. : Java.
- lanosulum* C. M. in Linn. 1876, p. 260.  
Eriocladium comorense Hpe in sched.  
2. Filic. — Afr. : Ins. Comor., Madagascar.
- lanosum* Mitt. in Journ. of the Linn. Soc. 1868, p. 170.  
Meteorium lanosum Mitt. M. Ind. or. p. 90 (1859).  
M. longissimum Dz. et Mk. M. frond. ined. Archip. ind. p. 159, t. 48 ?  
fid. Mitt (1854).  
M. longissimum  $\beta$  tenue Bryol. jav. II, p. 87, t. 202, fid. Mitt. (1861-70).  
M. vitianum Sull. ? fid. Mitt.  
2. — As. : Tonkin., Hong-Kong, Sikkim, Ceylan, Sumatra.
- longipendulum* (C. M.) Par. Ind. p. 9 (1894).  
Eriocladium longipendulum C. M. in Dusen M. Camer. n. 35 ; Dusen in K. Sv. Vet.-Akad. Foehr. 1895, n. 2.  
2. Ster. — Afr. : Cameroon.
- longissimum* (Dz. et Mk.) C. M. in Linn. 1879, p. 260.  
Meteorium longissimum Dz. Mk. M. frond. ined. Archip. ind. p. 159, t. 48 (1854), fid. Mitt. ; Bryol. jav. l.c.  
Neckera Dozyana C. M. Syn. II, pp. 141 et 672 (1851).  
N. longissima Dz. et Mk. in Ann. sc. nat. 1844, p. 313  
2. Ad arbor. — As. : Ins. Molucc., Batjan, Celebes, Borneo, Sumbawa, Java, Banca, Sumatra.
- nanopteris* (C. M.) Par. Ind. p. 9 (1894).  
Eriocladium nanopteris C. M. in Dusen M. Camer. n. 268.  
— Afr. : Cameroon.
- plumarium* (Hpe) Par. op. cit. p. 10.  
Eriocladium plumarium Hpe Enum. M. Brasil. p. 41 (1879).  
Ster. — Am. merid. : Brasil. austr. br.
- prostratulum* C. M. in Linn. 1873, p. 260.  
Neckera prostratula C. M. in op. cit. 1874, p. 560.  
Ster. — As. : Ins. Philippin.

- pseudo-capense C. M. in *Besch. Fl. bryol. Réunion, etc.*, p. 127 (1880-81).  
 Meteorium capense Aongstr. in *Oefv. of K. Vet.-Akad. Foerh.* 1875, n. 3, p. 143.  
 Papillaria pseudo-capensis C. M. *ms. ap. Aongstr. l. c.*  
 2. fr. ign. — Afr. : Choa, Madagascar, ins. Sancta Maria, Mayotte, Bourbon., Franciae.
- pseudo-lanosum Broth. et Geh. in *Bibl. bot.* 1898, p. 17.  
 Ster. — Pac. : Nov. Guinea batav. (M. Arfak).
- speciosum Dz. et Mk. in *Nederl. Kruidk. Arch.* 11, p. 79 ( ) et *Bryol. jav.* 11, p. 83, t. 198 (1861-70).  
 Meteorium speciosum Mitt. *M. Ind. or.* p. 87 (1859).  
 2. Ad *Filices* arbor. — As. : Celebes, Amboine, Khasia, Ceylan.
- striatulum (Mitt.) Par. *op. et l. cc.*  
 Meteorium striatulum Mitt. *Fl. Vit.* p. 395 (1871).  
 Ster. Terr. — Pac. : Nov. Hebrid., ins. Austral.
- subpiligerum (Hpe) C. M. in *Linn.* 1876, p. 261 (in *adn. ad A. lanosulum*).  
 Neckera subpiligera Hpe in *Linn.* 1871, p. 217.  
 Papillaria subpiligera *Besch. Fl. bryol. Réunion, etc.*, p. 124.  
 Ster. — Afr. : Madagascar.
- trachypterum (C. M.) Par. *op. et l. cc.*  
 Eriocladium trachypterum C. M. in *Dusen M. Camer. n.* 450 ; *Dusen in K. Sv. Vet.-Akad. Foerh.* 1895, n. 2.  
 2. Ster. Secus torr. — Afr. : Cameroon.
- vitianum (Sull.) C. M. in *Linn.* 1876, p. 262.  
 Meteorium vitianum Sull. in *Am. Expl. Exped.* 1859, p. 22, t. 21.  
 — Pac. : Ins. Viti, Taïti.
- Wallichii C. M. — Papillaria.
- Warburgii Broth. in *Warburg, Monsonia, I*, p. 50 (1899).  
 Ster. — As. : Ins. Batjan.
- AFZELIA** Ehrh. *Pl. crypt. exs.* (1790).  
*cirrata* Ehrh. — Dicranoweisia.  
*crispa* Ehrh. — Weisia viridula.  
*heteromalla* Ehrh. — Ditrichum homomallum.  
*lanceolata* Ehrh. — Pottia.  
*inclinata* Ehrh. — Distichium.  
*pulvinata* Ehrh. — Grimmia.  
*pusilla* Ehrh. — Seligeria.
- Alaria** Kindb. *Check List of Eur. and N. Amer. M.* p. 74 (nom., sect. HYPNI) (1894).
- Aloidella** C. M. *Syn. II*, p. 202 (1851, sect. POLYTRICHI); W. P. Sch. *Syn. II* ed p. 188 (1876, sect. BARBULÆ = ALOINA).
- ALOINA** C. M. *Syn. II*, p. 596 (sect. BARBULÆ) (1851) : Kindb. *Laubm. Schwed. u. Norw.* p. 136 excl. n. 5 (1883). — 16 spec.

## aloides (Koch) Kindb. op. et l. cc.

- Barbula aloides Bruch et Fűrnr. in Fl. 1829, P. II, p. 598; Bryol. eur. fasc. 13-15 (vol. II), Mon. p. 15, t. 139 (1842); C. M. Syn. I, p. 506 (1849), W. P. Sch. Syn. II ed. p. 191 (1876); Par. Ind. p. 60 (1894).  
 Bryum rigidum Dicks. Dried. pl. n. 47 (1790).  
 Tortula aloides Aongstr. in Nov. Act. Soc. Upsal. XII, p. 377 (1844).  
 T. rigida Sm. Fl. brit. III, p. 1249 (1805).  
 Trichostomum aloides Koch mss.; Schultz in Nov. Act. Acad. Caes. Leop. XI, p. 197 (1823).

2. Muric. et ad terr. argill. — Eur. : Med. et merid.; Britann., ins. Melit. As. : Minor. Afr. : M. Sinai, Alger., Canar. Am. sept. : Terra nov.

## ambigua (Bryol. eur.) Limpr. Laubm. Deutschl.... I, p. 638 (1888).

- A. ericaefolia Kindb. op. cit. p. 137.  
 Barbula ambigua Bryol. eur. fasc. 13-15 (vol. II), Mon. p. 14, t. 139 (1842); C. M. op. et l. cc.; W. P. Sch. op. cit. p. 190; Par. op. et l. cc.  
 B. rigida Hedw. Deser. I, p. 65, t. 25, f. 3 et 5 (p.p. quoad ic. l. fid. W. P. Sch. l. c.) (1785); Brid. Bryol. univ. I, pp. 528 et 824 excl. syn. Schultz (1826).  
 Bryum acaulon, Erica tenuifoliae Ger. folio Dill. Hist. M. p. 388, t. 49, f. 55 (1741) et Hb.  
 Br. ericaefolium Neck. in Act. Acad. Theod. palat. II, p. 451 (1770).  
 Tortula ambigua Aongstr. in Nov. Act. Soc. Upsal. XII, p. 376 (1844).  
 T. ericaefolia Lindb. M. scand. p. 20 (1879).  
 T. rigida Brid. Sp. M. I, p. 16, t. 2 (1806).

2. Locis denud. argill. — Eur. : Praecip. med. et merid. : Britann., Norveg., ins. Melit. As. : Minor, Persia, Jenissei super. Afr. : Alger., Tunet. Am. sept. : M. Rupestr., Columb. brit., Illinois, Calif.

## anthropophila (C. M.) Broth. Natur. Pflanzenfam. p. 428.

- Barbula anthropophila C. M. in Nuov. Giorn. bot. ital. 1896, p. 99; Par. Suppl. Ind. p. 20 (1900).

2. Ad tecta. — As. : China centr. (prov. Schen-Si).

## brevirostris (Hook. et Grev.) Kindb. op. cit. p. 137.

- Barbula brevirostris Bruch, Müll. in Fl. 1829, p. 406 et Fűrnr. l. c. p. 599; Bryol. eur. l. c. Mon. p. 16, t. 138; C. M. op. cit. p. 597; W. P. Sch. op. cit. p. 189; Par. op. cit. p. 61.  
 B. rigida γ brevirostris Brid. Bryol. univ. I, p. 824 (1826).  
 Tortula brevirostris H. et Gr. in Brewster Edinb. Journ. I, p. 289 (1824).  
 T. enervis Hartm. Fl. Skand. II ed. (1832).

2. Ad terr. argillac. — Eur. : Spitzberg., Scet., Scand., Dania, Borussia., Saxon., Austr. inf., Salisb., Tirol., Helvet., m. Cenis. As. : Jenissei. Am. sept. : Groenland., sin. Hudson, M. Rupestr., Columb. brit.

## « var. Broidleri (Limpr.) Limpr. Laubm. Deutschl.... I, p. 636 (1888).

- Barbula Broidleri Limpr. in LVII, Jahresber. d. Schles. Gesells., p. 310 (1879).

— Eur. : Lungau.

## « var. elongata Aongstr. in Act. Soc. sc. Upsal., 1844, p. 375.

-- Eur. : Scand.

## calceolifolia (Spr.) Broth. op. et l. cc.

- Barbula calceolifolia Spr. mss.; Par. Ind. p. 65 (1894).  
 Tortula calceolifolia Mitt. M. austr. am. p. 157 (1869).

Terr. — Am. merid. : And. Quitens.

catillum (C. M.) Broth. op. et l. cc.

Barbula catillum C. M. in Linn. 1878-79, p. 329; Par. op. cit. p. 69.

2. — Am. merid. : Alp. Argentin. subtropic.

cucullatifolia (C. M.) Broth. op. et l. cc.

Barbula cucullatifolia C. M. in op. cit. p. 331; Par. op. cit. p. 69.

2. — Am. merid. Alp. Argentin. Tucuman.

ericaefolia Lindb. — A. ambigua.

galeata (C. M.) Broth. op. et l. cc.

Barbula galeata C. M. in op. cit. p. 330; Par. op. cit. p. 75.

Fl. et fr. ign. — Am. merid. : Alp. Argentin, Saltens.

hamulus (C. M.) Broth. op. et l. cc.

Barbula hamulus C. M. in Bull. hb. Boissier, 1897, p. 192; Par. Suppl. Ind. p. 25 (1900).

2. — Am. sept. : Guatemala.

leptotheca (W. P. Sch.) Broth. op. et l. cc.

Barbula leptotheca W. P. Sch. in Savatier M. Jap. n. 230; Besch. in Ann. sc. nat. 1893, p. 336; Par. Ind. p. 81 (1894).

2. — As. : Japon. (Nippon centr.)

macrorhyncha (Kindb.) Broth. op. et l. cc.

Barbula macrorhyncha Kindb. in Mac. Cat. p. 50 (1892); Par. op. cit. p. 82.

2. Terr. — Am. sept. : Ontario.

obliquifolia (C. M.) Broth. op. et l. cc.

Barbula obliquifolia C. M. in Nuov. Giorn. bot. ital. 1898, p. 178; Par. Suppl. Ind. p. 28 (1900).

2. — As. : China centr. (prov. Schen Si).

pusilla Kindb. — Pterygoneuron cavifolium.

recurvipatula (C. M.) Broth. op. et l. cc.

Barbula recurvipatula C. M. in Linn. 1880-82, p. 424; Par. Ind. p. 91 (1894).

Fl. et fr. ign. — Am. merid. : Argentin. Cordob.

rigida (Hedw. p.p., Schultz) Kindb. op. et l. cc.

A. stellata Kindb. Eur. a. N. Amer. Bryin. p. 270 (1897).

Barbula rigida Hedw. Deser. I, p. 65 p.p.; t. 25, f. 16 (1787); Schultz Rec. de Barbula p. 196, t. 32, f. 1 (1823); Bryol. eur. fasc. 13-15 (vol. II), Mon. p. 13, t. 131 (1842); C. M. Syn. I, p. 596 (1849); W. P. Sch. Syn. II, p. 189 (1876); Par. op. cit. p. 91.

Bryum stellatum Schreb. Spic. fl. lips. p. 80 (1771).

Desmatodon rigidus Mitt. M. Ind. or. p. 38 (1859).

Tortula enervis Hook. et Grev. in Brewster Edinb. Journ. I, p. 288 (1824).

T. rigida Schrad. Spic. fl. germ. p. 66 (1794).

T. stellata Lindb. de Tortul. p. 233 (1864).

2. Ad mur. et rup. schist. calc. et arenac. semidecomp. — Eur. : Temper. et calidior. As. : Alatau, Caucas., Tibet occid. Afr. : Alger. Am. sept. : Niagara, Illinois, M. Rupestr., Colorado.

« var. mucronulata (Bryol. eur.) Par.

Barbula rigida var. mucronulata Bryol. eur., W. P. Sch. et Par. op. et l. cc.

Ut typ. — Eur. : liisd. loc. — Am. sept. : Calif. merid.

« *var. obtusa* (*Jur.*) *Par.*

*Barbula rigida var. obtusa Par. op. et l. cc.*

*Tortula rigida var. obtusa Jur. Laubm. Oesterr. Ung. p. 126 (1882).*

*Linic. — Eur. : Cum typ.*

« *var. pilifera* (*Bryol. eur.*) *Par.*

*Barbula rigida var. pilifera Bryol. eur., W. P. Sch. et Par. II. cc.*

*Tortula bifrons de Not. Sp. de Tortul. ital. n. 17 (1831) et Syll. n. 176 (1838).*

*T. rigida β piligera de Not. Musc. ital. p. 19, t. 4 (1862).*

*Ut typ. — Eur. : liisd. loc. Am. sept. : M. Rupestr.*

*sedifolia* (*C. M.*) *Broth. op. et l. cc.*

*Barbula sedifolia C. M. in Linn. 1878-79, p. 327; Par. op. cit. p. 91.*

*Fl. et fr. ign. — Am. merid. : Argentin. Cordob.*

*stellata* *Kindb. — A. rigida.*

*Sullivantiana* (*C. M.*) *Broth. op. et l. cc.*

*Barbula Sullivantiana C. M. in Hedwig, 1898, p. 130; Par. Suppl. Ind. p. 32 (1900).*

2. — *Pac. : Austral. or. (Victoria).*

***Aloma* C. M. in Dusen M. Camer. (sect. FISSIDENTIS).**

***Alomidium* C. M. in Fl. 1897, p. 328 (nom., sect. CONOMITRII sensu Müller.)**

***Alophos* Ren. et Card. in Bull. Soc. roy. bot. Belg., 1899, II, p. 225 sect. LYELLIE).**

**ALSIA Sull. in Proceed. of the Amer. Acad..... III, p. 185 (1854). — 3 spec.**

*abietina* (*Hook.*) *Sull. Ic. M. p. 115, t. 72 b (1864); Lesq. et Jam. Man. p. 279 (1884).*

*A. circinata Sull. Ic. M. t. 72; Lesq. in Trans of the Amer. phil. Soc. XIII, nom. ( ); Par. Ind. p. 11 (1894).*

*A. Macounii Kindb. in Bull. Torr. bot. Club, XVII, p. 275 (1890).*

*Leptodon abietinus Kindb. Eur. a. N. Amer. Bryin. p. 20 (1892).*

*L. circinatus Sull. in Pac. R. Rep. IV, p. 189, t. I (mala).*

*Neckera abietina Hook. M. ex. t. 27 (1818-20); C. M. Syn. II, p. 142 (1851).*

*Pilotrichum abietinum Brid. Bryol. univ. II, p. 258 (1827).*

2. *Ad arb. — Am. sept. : Ins. Vancouver, Columb. brit., Washington, Oregon, Idaho, Calif.*

*californica* (*Hook. et Arn.*) *Sull. in Proceed. of the Amer. Acad..... I. c. et M. of U. St. (1856); Lesq. et Jam. Man. p. 280.*

*Antitrichia pseudo-californica Kindb. Check List of Eur. and N. Am. M. (1894).*

*Neckera californica Hook. et Arn. in Beechey's Voy. 1841, p. 162; C. M. Syn. II, p. 117 (1851).*

2. *Ad arb. — Am. sept. : Ins. Vancouver, Washington, Calif.*

« *var. flagellifera* *Ren. et Card. in Bot. Gaz. 1889, p. 97.*

*Ut typ. — Am. sept. : Calif.*

*circinata* *Sull. — A. abietina.*

*longipes Sull. Ic. M. Suppl. p. 85, t. 53 (1874); Lesq. et Jam. op. et l. cc.*

2. *Rupic. — Am. sept. : Calif.*



**AMBLYODON** P. B. Prodr. p. 41 p.p. (1805). — 1 spec.

*dealbatus* (Dicks.) P. B. op. et l. cc.; *Bryol. eur. fasc.* 10 (vol. IV), t. 307 (1841); C. M. Syn. I ed. p. 127 (1849); W. P. Sch. Syn. II ed. p. 496.

*Bryum Celsii* L. hb. p.p.

*Br. dealbatum* Dicks. *Pl. crypt. fasc.* II, p. 8, t. 5, f. 3 (1790).

*Meesea dealbata* Sw. *M. Suec. pp.* 44 et 91, t. 5, f. 10 (1799).

*M. Macounii* Aust. in *Bull. Torr. bot. Club*, V, p. 22 (1878).

l et Syn. Loc. uligin., torfac., ad terr. et rup. humid. — *Eur.* : Reg. alpina per. zon. intermed. et sept., rar. merid. *As.* : Caucas., Alatau. *Am. sept.* : Sin. Hudson, Anticosti, Nov. Scot., New Brunswick, Gaspes., Wisconsin, M. Rupestr., Colorado.

“ *var. americanus* Ren. et Card. in *Bull. hb. Boissier*, 1896, p. 13 et *Obs et rectif.* p. 15.

— *Am. sept.* : Minnesota.

*longisetus* P. B. — *Meesea*.

*minor* P. B. — *Meesea uliginosa var. minor*.

*stygius* P. B. — *Cinclidium*.

*uliginosus* P. B. — *Meesea*.

**Amblyophyllum** C. M. Syn. I, p. 286 (sect. *BRV*).

**Amblyothallia** C. M. in *Bull. hb. Boissier*, 1897, p. 172 (nom., sect. *FISSIDENTIS*).

**AMBLYSTEGIUM** Bryol. eur. fasc. 55-56 (Vol. VI) (1853). — 68 spec.

Spec. omn. non indicatae sunt *HYPNA* ejusd. nom.

*adnatum* (Hedw.) Jaeg. Ad. II, p. 547 (1877-78).

*Hypnum adnatum* Hedw. *Sp. M.* p. 248, t. 64 (1801); C. M. Syn. II, p. 339 (1851); Sull. *Io. M.* p. 197, t. 121 (1864); Lesq. et Jam. *Man.* p. 375 (1884).

*H. Stereodon adnatus* Brid. *Bryol. univ.* II, p. 191 (1827).

*Stereodon adnatus* Mitt. in *Trans. of the Linn. Soc.* 1891, p. 180.

l. Cortic. et rupic. — *As.* : Japon. *Am. sept.* : New Brunswick, Ontario, Montana, Dakota, M. Rupestr., Texas; reg. sept., centr. et or.

“ *var. gracile* (Sull.) Par. Ind. p. 12 (1894).

*Hypnum adnatum var. gracile* Sull. *M. bor. am.* II ed. p. 86 (1865).

Ad rup. arenac. — *Am. sept.* : Ohio.

*aduncum* Lindb. — *Hypnum uncinatum*.

*var. legitimum* Sanio. — *Hypnum Sendtneri*.

*var. orthothecioides* Lindb. — *Hypnum orthothecioides*.

*afro-serpens* C. M. in *Dusen M. Camer. n.* 867.

— *Afr.* : Cameroon.

*alaskanum* Lesq. et Jam. — *Hylocomium*.

*ambiguum* de Not. — *A. Kochii*.

*Anzianum* de Not. — *A. Sprucei var. Anzianum*.

*argillicola* Lindb. in *Broth. Enum. M. Caucas. p.* 102 (1892).

l. In argill. humid. — *As.* : Caucas. centr.

- austro-fluviatile (*C. M.*) *Par. Ind.* p. 12 (1894).  
*Hypnum austro-fluviatile C. M. Bryol. austr. Georg.* p. 45 (1890).  
 Ster. Aquat. — *Am. merid.* : Austro-Georg.
- badium*  $\times$  *latifolium* Arn. — *H. badium*  $\times$  *Wilseni*.  
*brachypelmatum C. M. in Hedwig.* 1897, p. 131.  
 l. Paludic. — *Am. merid.* : Argentin. temper.  
 Ster. Aquat. — *Am. merid.* : Austr. Georg.
- Burnati Aman in Bull. hb. Boissier*, 1895, pp. 442-444 c. ic.  
 — *Afr.* : Ægypt.
- campicolum (C. M.) Pur. Ind.* p. 12 (1893).  
*Hypnum campicolum C. M. in Linn.* 1878-79, p. 444.  
 l. Muric. et terr. — *Am. merid.* : Argentin. Cordob.
- capillatum* Mitt. — *Thuidium*.  
*Cashii du Buys.* — *A. serpens* var. *depauperatum*.  
*chilense Ltz. in Bot. Zeit.* 1866, p. 188.  
 Ster. — *Am. merid.* : Chile austr.
- chloropelma C. M. in Hildebrandt M. Mad. n.* 2076.  
 — *Afr.* : Madagasc. or.
- chrysophyllum* var. *Zemliae* Jens. — *Hypnum Zemliae*.
- Closteri (Aust.) Curd. in litt.*  
*Amblystegium Holzingeri R. C. in Bot. Gaz.* 1894, p. 240, t. XXI, f. 6.  
*Hypnum Klosteri Aust. M. Appal. n.* 439.  
 — *Am. sept.* : M. Appalach., Virgin., distr. Columb.
- cochlearifolium* Lindb. — *Hypnum Goulardi*.
- compactum (C. M.) Jaeg. Ad. II*, p. 554 (1877-78).  
*A. serratum Bryol. eur. fasc.* 55-56 (*Vol. VI*), *Mon. p. II in adn.* (1853) ?  
*Sull. M. of U. St. p.* 782 ? (1856).  
*Hypnum compactum C. M. Syn. II*, p. 408 (1851) ; *Sull. Ic. M. p.* 201, t. 123 (1864) ; *Lesq. et Jam. Man. p.* 375 (1884).  
*H. serpens* var. *compactum Hook. in Drumm. M. bor. am. I ed. n.* 188.  
*Stereodon compactus Mitt. in Proceed. of the Linn. Soc.* 1864, p. 43.  
 l. Lignic., rupic. et terr. — *Eur.* : Anglia, Gall. centr. (Arvern.)  
*Am. sept.* : Terra nov., S. W. Canada, Wisconsin, Minnesota, Montana, Nevada, Utah, M. Rupestr., Washington, Calif.
- complanatulum* Lindb. — *Hypnum aduncum* var. *complanatulum*.  
*compressum* Mitt. — *Pseudo-Leskea*.
- confervoides (Brid.) Bryol. eur. fasc.* 55-56 (*Vol. VI*) *Mon. p. 6, t.* 562 (1853) ; *W. P. Sch. Syn. II ed. p.* 707 (1876) ; *R. du Buysson Essai sur le g. Amblystegium, II ed., p.* 9 (1889).  
*Hypnum Conierva Schwaegr. Suppl. II, P. I, fasc. 2, p.* 158, t. 142 (1823).  
*H. confervoides Brid. Sp. M. II, p.* 153 (1812) ; *C. M. Syn. II, p.* 142 (1851).  
*H. jungermannioides Brid. Bryol. univ. II, p.* 549, *fid. C. M.* (1827).  
*H. Stereodon confervoides ej. op. cit. p.* 563.  
*Leskea confervoides Spr. in Lond. Journ. of bot.* 1845, p. 182.  
*Stereodon confervoides Lindb. M. scand. p.* 38 (1879).  
 l. Ad saxa præpr. calc. — *Eur.* : E reg. campestr. usq. in subalp. passim.  
*As.* : Caucas. *Am. sept.* : New Brunswick, Nov. Anglia, Ohio, Ontario, M. Rupestr.

*convolutifolium* Jaeg. — *Rhynchostegium*.

*cordifolium* var. *coloratum* Arn. — *H. cordifolium* var. *coloratum*.

*curvicaule* (Jur.) Dix. et Jam. *Stud. Handb.* p. 447 (1896).

*A. flicinum* *curvicaule* Lindb. *M. scand.* p. 32 (1879).

*A. flicinum* var. *curvicaule* Mol. *Bay. Laubm.* p. 238 (1875).

*Hypnum* *curvicaule* Jur. in *Verh. d. K.K. zool. bot. Ges. in Wien* 1864, p. 103; *Par. Ind.* p. 628 (1895).

*H. flicinum* var. *curvicaule* Berggr. in *K. Sv. Vet.-Akad. Handl.* 1875, n. 7, p. 86.

? *Limnobium* *curvicaule* de Not. *Epil.* p. 164 (1869).

2. ♂ et fr. ign. Ad. rup. humid. — *Eur.*: Spitzberg, ins. Ursor., Alp. Julian., Rhaet., Bavar., Stir., Carinth., Salisb., Tirol, Tatra, etc.

\* var. *strictum* Dix. et Jam. *op. cit.* p. 448.

Ut typ. — *Eur.*: Scotia (Ben Lavers).

*curvipes* Gumb. in *Bryol. eur. l. c.* p. 14, t. 569; *W. P. Sch. op. cit.* p. 717.

*A. Kochii* var. *curvipes* Husn. *Muscol. gall.* p. 363 (1894).

*A. riparium* var. *curvipes* R. du Buysson, *op. cit.* p. 20.

1. In stagn. exs. — *Eur.*: Fennia? Siles., Austr. infer., Palatin. As.: Jenissei, Caucas. (Tiflis). *Am. sept.*: Ottawa, Ontario.

*densissimum* Card. in *Rev. bryol.* 1900, p. 106.

Ster. — *Am. merid.*: Fret. Gerlache.

*densum* Milde in *Bot. Zeit.* 1864, *Beit.* p. 21.

*A. compactum* (fid. Dixon).

*A. serpens* var. *cryptarum* Arnold. (Sauter) in *Rab. Bryoth. eur.* n. 841.

*Brachythecium* *densum* Jur. in *Coll. cit.* n. 995.

*Hypnum* *densum* Milde *Bryol. siles.* p. 360 (1869).

*Rhynchostegium* *tenellum* var. *brevifolium* Lindb. in *Coll. cit.* n. 750.

2. Ad. rup. calcar. — *Eur.*: Suecia, Palatin. infer., Austr. inf., Francon., Stir., Carinth., Tirol., Catalaun.

*dissitifolium* Kindb. in *Mac. Cat.* p. 220 (1892).

2? ♀ ign. Ad. rup. madid. — *Am. sept.*: Niagara, Owen Sound.

*distantifolium* Kindb. in *op. cit.* p. 222.

Ster. Rupic. — *Am. sept.*: Terra nov., Arkansas.

*enerve* *Bryol. eur. fasc.* 55-56 (Vol. VI), *Mon. p.* 8, t. 563 (1853); *W. P. Sch. op. cit.* p. 709.

*Hypnum* *incurvatum* forma (fid. Limpr. *Laubm. Deutschl.... III*, p. 294) (1897).

*Stereodon* *enervis* Lindb. *M. scand.* p. 38 (1879).

1. Rupic. — *Eur.* — Norvegia (Dovre).

*exannulatum* de Not. var. *alpicola* ej. — *Hypnum* *exannulatum* var. *purpurascens*.

*Falconeri* Mitt. in *Hb. of the Late East Ind. Comp.* n. 656.

— As.: N. W. Himalaya.

*fallaciosum* Lindb. — *Hypnum* *polygamum* var. *fallaciosum*.

*fallax* (Brid.) Milde *Bryol. siles.* p. 325 (1869); Lindb. *M. scand.* p. 32 (1879).

*A. flicinum* var. *crassinervium* Ren. in *Husn. Musc. gall.* p. 362 (1884-90) (1).

---

(1) Cf. Limpr. *Laubm. Deutschl.... III*, p. 309 (1897) et Card. in *Bull. hb Boissier*, 1899, p. 324.

- A. filicinum var. fallax *Lindb. M. scand.* p. 35.  
 A. filicinum var. Vallis Clausae *Dixon et Jam. Stud. Handb.* p. 446 (1896).  
 A. fluviatile var. fallax *Bryol. eur. fasc. 55-56 (Vol. VI), Mon.* p. 12.  
 A. formianum *Dna Fior. Mazz. in Atti dell' Acad. pontif. d. nuovi Lincei*, 1874, c. 10. (1).  
 A. irriguum var. fallax *W. P. Sch. Syn. I ed.* p. 501 (1860); *Rab. Bryoth. eur.* n. 342.  
 A. Vallis Clausae *Husn. Muscol. gall.* p. 361, t. 103 (1) (1893).  
 Hypnum fallax *Brid. Musc. rec. II*, p. 66, t. 2, f. I (1801).  
 H. filicinum var. fallax *H. et T. Muscol. brit.* p. 109 (1818).  
 H. filicinum var. Vallis Clausae *Wils. Bryol. brit.* p. 393 (1855).  
 H. formianum *W. P. Sch. Syn. II ed.* p. 741 (1876).  
 H. Vallis Clausae var. fallax *Houl. M. Fr. I*, p. 51 (1834).  
 Thuidium fallax *Kindb. Laubm. Schwed. u. Norw.* p. 10 (1883).

2. Ad saxa praecip. calc. saepius inund. — *Eur.* : Fere tota. *As.* : Jenissei, Caucas. *Afr.* : Alger. *Am. sept.* : Lac. Super., Pensilv.

« var. nigricans *Sauter Fl. Herzogth. Salz. III*, p. 67 (1870).

A. Vallis Clausae var. atro-virens *Husn. op. et l. cc.*

Ad scaturig. frigid. — *Eur.* : Salisb., Gall. occid.

« var. spinifolium (*W. P. Sch.*) *Limpr. Laubm. Deutschl. .... III*, p. 308 (1897).

A. irriguum var. spinifolium *W. P. Sch. Syn. II ed.* p. 713 (1876); *R. du Buysson, op. cit.* p. 18.

A. noterophilum (*Sull.*) *Holzinger in Bull. geol. et nat. Hist. Survey Minnesota*, IX, p. 293.

Hypnum noterophyllum *Sull. M. of U. St.* p. 78 (1856); *Sull. et Lesq. M. bor. am. I ed.* n. 348 (spec. propria, fid. *Limpr. Nachtraege*, p. 830).

H. Vallis Clausae var. spinifolium *Husn. op. et l. cc.*

Ut typ. — *Eur.* : Westphal., Helvet., Gall., Graecia. *Am. sept.* : Ubi typ.

fenestratum *Kindb. in Mac. Cat.* p. 217 (1892).

A. tenuifolium *Kindb. in Out. Nat. IV*, p. 65 (1889).

Ster. Ad. terr. humid. — *Am. sept.* : Ontario.

• filicinum (*L.*) *de Not. Cronac. II*, p. 25 (1867); *Milde Bryol. siles.* p. 325 (1869).

Hypnum affine *Hoffm. Deutsch. Fl. II*, p. 61 (1796).

H. compressum *Brid. Musc. rec. II, P. II*, p. 58 (1801).

H. dubium *Neck. Meth. M.* p. 161 (1771).

H. extricatum *Hoffm. op. et l. cc.*

H. filicinum *L. Sp. pl.* p. 1125 (1753); *Brid. Bryol. univ. II*, p. 527 (1827); *C. M. Syn. II*, p. 419 (1851); *Bryol. eur. fasc. 55-56 (Vol. VI), Mon.* p. 40, t. 609 (1853); *W. P. Sch. Syn. II ed.* p. 740 (1876); *Par. Ind.* p. 637 (1895).

H. filicinum  $\alpha$  spicatum *Weis Fl. goetting.* p. 229 (1770).

H. filicinum  $\beta$  pendulum *Brid. op. cit.* p. 582.

H. filicinum  $\gamma$  pinnatum *ej. op. et l. cc.*

H. flaccidum ( *Schleich. in sched.*

H. glareosum (

*Stereodon filicinum Mitt. in Proceed. of the Linn. Soc.* 1861, p. 43.

*Thuidium filicinum Kindb. Laubm. Schwed. u. Norw.* p. 10 (1883).

(1) Cf. *Limpr. Laubm. Deutschl. .... III*, p. 309 (1897) et *Card. in Bull. hb. Boissier*, 1899, p. 324.

2. Ad saxa, mur., terr. et lign. madid. in aquos. calcar. — *Eur.* : Spitzberg, reg. campestr. et montan. *As.* : Japon., Saghalien, Ochotsk, Jenissei, Yunnan, Himalaya et Tibet. occ., Pandjab, Kashmir, Tchitral, Afghan., Alatau, Turkestan., Luristan., Kurdistan., *Syr. Afr.* : Alger. *Am. sept.* : Fret. Hudson, Anticosti, Terra nov., Canada merid.; de Nova Scot. ad Saskatchevan, Alaska, ins. Vancouver, Columb. brit., Washington, Calif., M. Rupestr., reg. sept., centr. et or. *Pac.* : Nov. Seelandia.
- « *var. aciculinum* (*C. M. et Kindb.*) *Par.*  
*Hypnum filicinum var. aciculinum C. M. et Kindb. in Mac. Cat. p. 231 (1892).*  
 Ad rup. madid. — *Am. sept.* : Columb. brit.
- « *var. elatum* (*W. P. Sch.*) *Limpr. Laubm. Deutschl. .... III, p. 304 (1897).*  
*Hypnum filicinum var. elatum W. P. Sch. op. cit. p. 740.*  
 In spongios. — *Eur.* : Alp. Salisb. *Am. sept.* : Hic illic.
- « *var. filiforme* (*Berggr.*) *Par.*  
*Hypnum filicinum var. filiforme Berggr. M. Spetsb. n. 136 b. et in K. Sv. Vet. Akad. Föerh. XIII, N. 7, p. 87 (1875).*  
 Aquat. — *Eur.* : Ins. Ursor.
- « *var. floridanum* (*Ren.*) *Par.*  
*Hypnum filicinum var. floridanum Ren. in Lesq. et Jam. Man. p. 387 (1884).*  
 — *Am. sept.* : Florida.
- « *var. gracilescens* (*Bland.*) *Limpr. op. et l. cc.*  
*Hypnum dubium Dicks. Pl. crypt. fasc. III, p. 10 (1793).*  
*H. filicinum var. gracilescens W. P. Sch. Syn. I ed. p. 614 (1860).*  
*H. gracilescens Bland. in Brid. Sp. M. II, p. 211 (1812).*  
 Ad rup. minus humid. — *Eur. et Am. sept.* : Cum typo.
- « *var. gracillimum* *Sael. in Act. Soc. p. Faun. et Fl. fenn. VI, n. 4 (M. Lapon. Kolaënsis).*  
 — *Eur.* : Karelia.
- « *var. lacustre* (*Leiner*) *Par.*  
*Hypnum filicinum var. lacustre Lein. in Rab. Bryoth. eur. n. 399.*  
 — *Eur.* : Boden See.
- « *var. locarnense* *de Not. Epil. p. 115 (1869).*  
*Hypnum filicinum f. falcata Boul. M. Fr. I, p. 49 (1884).*  
 — *Eur.* : Gall. merid., Ital. super.
- « *var. patulum* (*Roell*) *Par.*  
*Hypnum filicinum var. patulum Roell in Hedwig. 1897, Beibl. II, p. 42.*  
 In rivul. — *Eur.* : Hispan. merid.
- « *var. pendulum* (*Brockm.*) *Par.*  
*Hypnum filicinum var. pendulum Brockm. Laubm. Mecklemb. p. 142 (1870).*  
 — *Eur.* : Mecklenbourg.
- « *var. prolixum* (*Vill.*) *de Not. Epil. p. 151 (1869).*  
*Hypnum filicinum var. elongatum von Klinggr. Leb. u. Laubm. W. u. Ostpreuss. p. 282 (1893).*  
*H. filicinum var. submersa Pfeff. Bryogeog. Stud. p. 89 ?*  
*H. prolixum Vill. Pl. Dauph. III, p. 914 (1786).*  
 Ad cataract. — *Eur.* : Passim.

« *var. saxicola* (M. T. Lge.) Par.

*Hypnum filicinum var. saxicola* M. T. Lge. in *Lot. Tidsskr.* 1869, p. 36.

— *Eur.* : Dania.

« *var. supra-alpinum* (Mol.) Par.

*Hypnum filicinum var. supra-alpinum* Mol. *Alg. Moosst.* p. 102 (1865)

— *Eur.* : Alp. Algov.

« *var. tenue* (Berggr.) Par.

*Hypnum filicinum var. tenue* Berggr. *ll. cc. n.* 136 c.

Rupic. — *Eur.* : Spitzberg, ins. Ursor.

« *var. trichodes* (Brid.) Par.

*Hypnum filicinum f. tenuis* Boul. *M. Fr. I*, p. 50 (1884).

*H. filicinum var. trichodes* Stend. *Nom. crypt.* p. 211 (1821); W. P. Sch. *op. et l. c.*

*H. repens filicinum, trichodes palustre* Dill. *Cat. Giss.* p. 218 (1718), *Hist. M. p.* 286, t. 36, f. 21 (1741) et *Hb.*

*H. trichodes* Brid. *Sp. M. II*, p. 235 (1812).

Ad scatur. herb., terr. et saxa. — *Eur. et Am. sept.* : Ubi typ.

<i>filicinum</i> varr.	{	<i>crassinervium</i> Ren.	{	<i>A. fallax.</i>
		<i>fallax</i> Lindb.		
		<i>Vallis Clausae</i> Dix. et Jam.		

*finmarchicum* (Ltz.) R. du Buyss. *op. cit.* p. 18.

*A. serpens* ? (Limpr. Laubm. *Deutschl...* III, p. 294).

*Hypnum finmarchicum* Ltz. in *Hartm. Skand. Fl. X ed.* p. 20 (1871).

l. — *Eur.* : Finmarck.

*floridanum* R. C. — *A. riparium var. floridanum.*

*fluitans* Mitt. (1876). — *Hypnum austro-fluitans.*

var. *exannulatum* Lindb. — *Hypnum exannulatum.*

Rotae Lindb. et Arn. — *Hypnum exannulatum var. Rotae.*

*fluitans* × *pseudo-stramineum* Lindb. — *Hypnum pseudo-stramineum.*

*fluviatile* (Sic.) *Bryol. eur. fasc.* 62-64, *Suppl. t. I* (1855), *sed non Bryol. eur. fasc.* 55-56 (*Vol. VI*), t. 567 (1853); W. P. Sch. *op. cit.* p. 713.

*Hypnum arhrizon* Brid. *Bryol. univ. II*, p. 535 (1827).

*H. caucasicum* ej. l. c.

*H. diffusum* Bland. in *Brid. Sp. M. II*, p. 243 (1812); *Brid. op. cit.* p. 538 (1827).

*H. filicinum* ± *lanatum* *Brid. op. cit.* p. 531.

*H. flagelliforme* ej. *Sp. M. II*, p. 240.

*H. fluviatile* Sw. *Disp. M. Succ.* p. 63 (1799); *Brid. op. cit.* p. 532; C. M. *Syn. II*, p. 420 (1851).

*H. fluviatile* ♂ *flagelliforme* *Brid. op. cit.* p. 533.

*H. irriguum* \* *fluviatile* Boul. *M. Fr. I*, p. 73 (1884).

*H. luridum* Sw. *op. cit.* p. 68.

*H. palustre var. fluviatile* Wahlenb. *Fl. succ. II ed.* p. 732 (1833).

*H. Seligeri* Brid. *Musc. rec. II, P. II*, p. 67 (1801).

*H. Vallis Clausae* Brid. *Sp. M. II*, p. 238 (1812) et *Bryol. univ. II*, p. 531 (1827) (1.)

l. Ad saxa rivul. — *Eur.* : Reg. campestr. et mont. Scand., German., Britann., Gall. sept.; Liguria. *Am. sept.* : Terr. nov., S. W. Canada, New. Jersey, Wisconsin, Alabama.

(1) Fid. C. M. l. c.; cf. Renauld. in *Rev. bryol.* 1885, p. 55.

« *var. brevifolium* Ren. et Card. in Bot. Gaz. 1900, p. 22.  
— *Am. sept.* : Minnesota.

*var. fallax* Bryol. eur. — *A. fallax*.

« *var. julaceum* R. du Buysson op. et l. cc.  
Ut et ubi typ. in Eur.

*fluviatile* Bryol. eur. — *A. irriguum*.

*var. irriguum* R. du Buyss. — *A. irriguum*.

*formianum* D<sup>na</sup> Fior. Mazz. — *A. fallax*.

*glaucum* Lindb. — *Hypnum commutatum*.

*var. sulcatum* Lindb. — *Hypnum sulcatum*.

*glaucum* × *falcatum* Lindb. — *Hypnum falcatum*.  
*gracile* Jur. — *Pseudo-Leskea*.

*Haplocladium* C. M. in Hedvig. 1897, p. 130.

*Tamariscella muda* C. M. in sched.

1. Cortic. et terr. — *Am. merid.* : Paraguay.

*Hausmanni* de Not. in sched. ; W. P. Sch. Syn. II ed. p. 720 (1876)

*A. riparium* var. abbreviatum (non Bryol. eur.) de Not. Epil. p. 147 (1869) ;  
*Husn. Musc. gall.* p. 363 (1894) ; *Braithw. Brit. Mossfl. III*, p. 30 (1896).

*A. riparium* var. *Hausmanni* de Not. in sched. ; Vent. et Bott. Enum.  
crit. p. 14 (1884).

1. Ad arb. rad. et puteor. ductus. — *Eur.* : Austria infer., Tirol.

*Holzingeri* R. C. — *A. closteri*.

*homalostegium* (C. M.) Jaeg. Ad. II, p. 547 (1877-78) (1).

*Hypnum homalostegium* C. M. in Fl. 1873.

*Pseudo-Leskeella homalostegia* Kindb. Eur. a. N. Am. Bryin. p. 48 (1897).

1. Ad arb. — *Am. sept.* : Alabama.

*hygrophiloides* W. P. Sch. in Savatier M. Japon. n. 715.

1 ? Terr. — *As.* : Japon.

*hygrophilum* (Jur.) W. P. Sch. Syn. II ed. p. 715 (1876).

*A. hygrophilum* var. *limosum* W. P. Sch. et R. du Buyss. op. cc.  
pp. 716 et 21.

*A. riparium* var. *hygrophilum* Bryhn in Norsk. Mag. for Naturv. 1891.

*A. saxatile* W. P. Sch. Syn. I ed. p. 595 (1860).

*Campylium hygrophilum* Kindb. Eur. a. N. Am. Bryin. p. 121 (1897).

*Hypnum fragile* Brid. Sp. M. II, p. 198 (1812) fide Hpe. Fl. hercyn.  
p. 361 (1873).

II. *hygrophilum* Jur. in. Rab. Bryoth. eur. N. 649 (1863) ; Milde, Bryol.  
siles, p. 340 (1869).

II. *serpens* L. herb. teste W. P. Sch.

H. Sommerfeldtii var. *Mol. Bayer. Laubm.* p. 249 (1875).

1. Paludic. et ad terr. aren. humid. — *Eur.* : Boruss., Siles., Harz,  
Thuring., Hassia, Stir., vall. Rhen. med. *Am. sept.* : Idaho ; Illinois ?  
distr. of Columb ?

*hygrophilum* H. M. — *Hypnum polygamum* var. minor.

var. *limosum* W. P. Sch. — *A. hygrophilum*.

*incurvatum* β *Blyttii* Kindb. — *Hypnum Blyttii*.

*intermedium* Lindb. — *Hypnum*.

var. *revolvens* Sanio. — *Hypnum revolvens*.

var. *verum* Sanio. — *Hypnum intermedium*.

(1) In Lesq. et Jam. Man. deest.

irriguum (Wils.) *Bryol. eur. fasc. 62-64, Suppl. ad gen. Amblystegium* (1855); *W. P. Sch. Syn. II ed. p. 712* (1876).

*A. fluviatile* *Bryol. eur. fasc. 55-56 (Vol. VI), Mon. p. 11, t. 566 excl. var. γ* (1853).

*A. fluviatile* var. *irriguum* *R. du Buyss. op. cit. p. 18* (1889).

*Hypnum fluviatile* plur. auct. (non Sic.)

*H. irriguum* *Wils. Bryol. brit. p. 361, t. 2* (1865).

*H. tenax* *Hedw. Sp. M. p. 271, t. 72* (1801); *Brid. Bryol. univ. II, p. 496* (1827) *fid. Cardot.*

1. *Ad saxa praecip. silic., mur. molend. irrig.* — *Eur.* : Fere tota, rarius in sept. : *As.* : *M. Altaï, Caucas. occ. Afr.* : *Alger. Am. sept.* : *New Jersey, Virgin. occ., Ontario, Minnesota, Montana, Calif.*

var. *fallax* *W. P. Sch.* — *A. fallax.*

« var. *flaccidum* *de Not. Epil. p. 158* (1869).

*Ad aquaeduct.* — *Eur.* : *Longobard. (Mediolan.)*

var. *spinifolium* *W. P. Sch.* — *A. fallax* var. *spinifolium.*

« var. *tenellum* *W. P. Sch. Syn. II, ed. p. 713.*

— *Eur.* : *Passim.*

*Juratzkanum* *W. P. Sch. Syn. I ed. p. 693* (1860) *et II ed. p. 710.*

*A. serpens* var. *Juratzkanum* *R. du Buyss. op. cit. p. 18* (1889).

*A. Juratskæ* *Kindb. in Mac. Cat. n. 218* (1892).

*Eurhynchium Juratzkanum* *R. Farnetti in Att. d. Istit. bot. d. real. Univ. d. Pavia* (1893).

*Hypnum Juratzkanum* *Lindb. in C. Hartm. Skand. Fl. IX ed. (1864); Boul. M. Est Fr. p. 294* (1892) *p.p.*

*H. Juratzkæ* *Mac. Canad. M. n. 466.*

1. *Ad saxa humidiusc.* — *Eur.* : *German. sept., Siles., Hungar., Bohem., Salisb., Stir., Austria infer., Tirol., Ital. super., vall. Rhen. med., Gall. austr. occ., Pyren., Astur. As.* : *Caucas. Am. sept.* : *Ottawa, Ontario, Idaho, Montana, Dakota, Wisconsin, Ohio, distr. of Columb., Columb. brit., Maryland, Virgin.*

« var. *arenaceum* *Warnst. in Verh. d. K. K. zool. bot. Gesells. in Wien. p.*

*Ad rup. arenac. humid.* — *Eur.* : *Tirol. merid.*

*Kneiffii* *Bryol. eur.* — *Hypnum aduncum* var. *Kneiffii.*

*Kochii* *Bryol. eur. fasc. 55-56 (Vol. VI), Mon. p. 13, t. 568* (1853); *W. P. Sch. op. cit. p. 716.*

*A. ambiguum* *de Not. Epil. p. 144* (1869).

*A. riparium* var. *Kochii* *R. du Buyss. op. cit. p. 20* (1889).

*A. trichopodium* var. *Kochii* *Lindb. in Act. Soc. sc. Fenn. 1872, p. 275.*

*Hypnum ambiguum* *Jaeg. Ad. II, p. 569* (1877-78).

*H. riparium* \* *Kochii* *Boul. M. Fr. I, p. 71* (1884).

*Limnobia ambiguum* *de Not. Epil. p. 163.*

*L. palustre* *ej. in Erb. critt. ital. n. 906.*

1. *Ad rad. Salic. et terr. humid. vel. limos.* — *Eur.* : *German. sept., Siles., Hungar., Bohem., Morav., Harz, Thuring., Austr. infer., Stir., Salisb., Tirol., Helvet., Voges., Wetsphal. As.* : *Amour. Am. sept.* : *Minnesota ?, distr. of Columb., Indiana, Illinois, Kansas, Ludovic.*

« var. *arvense* *Warnst. in Verh. d. K. K. zool. bot. Gesells. in Wien, 1900, p.*

*Ad terr. humid.* — *Eur.* : *Tirol. merid.*

var. *curvipes* *Husn.* — *A. curvipes.*



« *var. Loeskeanum* Warnst. in *Verh. d. bot. Ver. d. Prov. Brandenb.* 1899, p. 76.

— *Eur.* : Lusatia infer.

*latifolium* Lindb. et Arn. — *Hypnum Wilsoni* ?

*var. jenissense* Sanio. — *Hypnum Wilsoni var. jenissense*.

*leptophyllum* W. P. Sch. *Syn. II ed. p.* 719 (1876); *R. du Buyss. op. cit. p.* 15.

*Hypnum leptophyllum* W. P. Sch. in litt.; Ravaut in *Bull. Soc. bot. Fr.* 1867, p. 200 (?)

1. Lignic. et ad rup. humid. — *Eur.* : Austria infer.

« *var. cylindricum* R. du Buyss. *Essai, I ed. p.* 18 et *op. cit. p.* 16.

*Hypnum leptophyllum* Husn. *Muscol. gall. p.* 359, t. 102 (?) (1894).

Ubi typ. — *Eur.* : Gall. (Arvern., Delphinat.).

*Lescurii* (Sull.) *Jaeg. Ad. II p.* 554 (1877-78).

*Hypnum Lescurii* Sull. *M. of U. St. p.* 79 (1856) et *Io. M. p.* 203, t. 24 (1864); *Lesq. et Jam. Man. p.* 376 (1884).

*Platyloma Lescurii* Kindb. *Eur. and N. Am. Bryin. p.* 70 (1897).

1. Ad rup. madid. — *Am. sept.* : Reg. sept. et or.

*longisetum* W. P. Sch. — *A. madeirense*.

*lucidum* Jaeg. — *Plagiothecium*.

*lycopodioides*  $\beta$  *vernicosum* Lindb. et Arn. — *Hypnum vernicosum*.

*madeirense* (Mitt.) *Jaeg. op. cit. p.* 549.

*A. longisetum* W. P. Sch. in *Mandon Pl. Mad. n.* 43 (*ex ips. ?*).

*Hypnum madeirense* Mitt. in *Journ. of the Linn. Soc.* 1863, p. 6.

1. Rupic. et terr. — *Afr.* : Ins. Canar., Madeir.

*minutissimum* (Sull. et Lesq.) *Jaeg. op. cit. p.* 547.

*Hypnum minutissimum* Sull. et Lesq. in *Sull. M. of U. St. p.* 78 (1856), *Sull. Io. M. p.* 195, t. 120 (1864); *Lesq. et Jam. Man. p.* 371 (1884).

2. Ad rup. et terr. humid. — *Am. sept.* : Ontario, M. Rupestr., Columb. brit., Illinois, Ohio, New Jersey, Pensilv., etc.

*molle var. alpinum* Lindb. — *Hypnum alpinum*.

*Mülleri* (Hpe. et C. M.) *Jaeg. op. cit. p.* 555.

*A. riparium* (fid. Mitt.).

*Hypnum Mülleri* Hpe. et C. M. in *Linn.* 1853, p. 504.

— *Pac.* : Austral. or.

*noterophilum* Holz. — *A. fallax var. spinifolium*.

*obtusulum* (Mitt.) *Jaeg. op. cit. p.* 551.

*Hypnum obtusulum* Mitt. *M. Ind. or. p.* 82 (1859).

2 ? — *As.* : W. Himalaya.

*octodiceroides* C. M. in *Hedwig.* 1898, p. 265.

*Hypnum riparium* Sull. in *Wright M. cubens. n.* 129.

Ster. Loc. humid. — *Am. sept.* : Cuba.

*oligorhizon* Gümb. in *Bryol. eur. fasc.* 55-56 (*Vol. VI*), *Mon. p.* 13, t. 568 (1853); W. P. Sch. *op. cit. p.* 712.

*A. radicale* Milde *Bryol. silcs. p.* 324 p.p. (1869).

*A. varium* R. du Buyss. *Essai, II ed. p.* 14.

*A. varium*  $\beta$  *oligorhizon* Lindb. *M. Scand. p.* 32 (1879).

1. Ad rup. molass. umbr. — *Eur.* : Fennia, Bavar.

orthocladon (*P. B.*) *Jaeg. Ad. II* p. 555 (1877-78).

*A. varium* var. orthocladon *Husn. Musc. gall.* p. 359 (1894).

*Hypnum* orthocladon *P. B. Prodr.* p. 72 (1805); *Brit. Bryol. univ. II*, p. 537 (1827); *Sull. Ic. M.* p. 199, t. 122 (1864); *Lesq. et Jam. Man.* p. 374 (1884).

*H. serpens* var. orthocladon *Aust. M. Appal.*

*H. serpens* var. *varium* *C. M. Syn. II*, p. 412 p.p. (1851).

*H. varium* *H. f. et W. in Drumm. M. bor. am. II* ed., n. 142.

1. Lignic., rupic. et terr. — *Am. sept.* : New Brunswick, Ontario, Idaho, Oregon, reg. sept., centr. et or.

*orthothecioides* Lindb. — *Hypnum uncinatum* var. *orthothecioides*.

*pachyrhizon* Lindb. — *A. porphyrhizon*.

*polycarpum* Vent. et Bott. — *Hypnum aduncum* var. *polycarpum*.

*polygamum* var. *stagnatum* Braithw. — *H. polygamum* var. *fallaciosum*

*porphyrhizon* (Lindb.) *W. P. Sch. Syn. II* ed. p. 715 (1876).

*A. pachyrhizon* Lindb. *M. Scand.* p. 32 (1879).

*A. radicale* ♂ serotinum (Lindb.) Braithw. *Brit. Mossft. III*, p. 26 (1896).

*A. serpens* ♂ serotinum Lindb. in *Hartm. Skand. fl. X* ed. II, p. 20 (1871).

*Hypnum* porphyrhizon *ej. in Hartm. Skand. fl. IX* ed. II, p. 12 (1864).

*H. serpens* ♂ serotinum *ej. in op. et l. cc.*

*H. radicale* *Lesq. et Jam. Man.* p. 373 p.p. (*fid. Mac.*); *Drumm. M. bor. am. I* ed. n. 189.

1. Lignic. — *Eur.* : Suec., Dania, Fennia. *Am. sept.* : Labrador, Anticosti, ins. Miquelon, Canada merid., reg. sept. occ., m. Rupes-tr., Columb. brit., Kansas.

*protensum* Lindb. — *Hypnum stellatum* var. *protensum*.

pseudo-confervoides *Kindb. in Rev. bryol.* 1895, p. 84.

1. Ad rup. calcar. — *Am. sept.* : Canada.

pulchellum *C. M. in Ule Bryoth. brasil. n.* 73.

— *Am. merid.* : Brasil. subtropic. or.

radicale (*P. B.*) *Mitt. (non Bryol. eur.) in Journ. of the Linn. Soc.* 1869, p. 569.

*A. serpens* var. radicale *Aust. M. Appal.*

*Hypnum bergenense* *Aust. M. Appal. n.* 391.

*H. humile* *P. B. Prodr.* p. 65 (1805).

*H. radicale* *ej. op. cit.* p. 68; *Schwaegr. Suppl. I, P. II*, p. 255, t. 90 (1816).

*H. Stereodon radicalis* *Brid. Bryol. univ. II*, p. 651 (1827).

*Stereodon radicalis* *Mitt. in Proceed. of the Linn. Soc.* 1864, p. 43.

1. Ad lign. putr., terr. et rad. arb. in humid. — *Eur.* : Angl., Suec., Fennia, Brandeb., Siles., Bade, Tirol. *As.* : Caucas. occ. et centr. *Am. sept.* : Anticosti, ins. Miquelon, Canada merid., Columb. brit., reg. sept., centr., or. et merid.

« var. *Lesquereuxii* *Par. Ind.* p. 18 (1894).

*A. radicale* *Sull. et Lesq. M. bor. am. I* ed. n. 224?

*A. varium* var. radicale *R. C. Fl. ins. Miquelon*, p. 53 1888.

Ut typ. — *Am. sept.* : Ins. Miquelon.

radicale { *Bryol. eur.* — *A. varium*.  
Milde p.p. — *A. oligorhizon*.  
Sull. et Lesq. — *A. radicale* var. *Lesquereuxii* ?

*rigescens* *Limpr. Laubm. Deutschl.... III*, p. 317 (1897).

*A. serpens* *var. rigidiusculum* *Arn. M. As. bor. II*, p. 113 (1890).

1. *Ad mur. et rup. arenac.* — *Eur.* : Norveg. bor., Lusat. *As.* : Jenuissei med., Obi.

*riparioides* (*Hpe.*) *Jaeg. Ad. II*, p. 553 (1877-78).

*Serpo-Hypnum riparioides* *Hpe Symb.* 1870, p. 291.

1. *Lignic.* — *Am. merid.* : Brasil. austr. or.

*riparium* (*L.*) *Bryol. eur. fasc.* 55-56 (*Vol. VI*), *Mon.* p. 14, t. 570 (1853); *W. P. Sch. op. cit.* p. 717; *R. du Buyss. op. cit.* p. 18.

*Fissidens Weberi* *Brid. Musc. rec. II, P. I*, p. 146 (1798).

*Hypnum aquaticum* *flagellis et teretibus et pinnatis* *Dill. Hist. M.* p. 308, t. 40, fig. 44 (1741).

*H. flexipes* *Brid. Bryol. univ. II*, p. 419 (1827).

*H. hornum* *Web. Spic. fl. goett.* p. 44 ? (1778).

*H. laxifolium* *Brid. Sp. M. II*, p. 91 (1812).

*H. oligophyllum* *ej. Musc. rec. II*, p. 181 (1801).

*H. palustre lutescens*, etc. *Vill. Pl. Dauph. III*, p. 918 (1786).

*H. palustre nigrescens* *ej. op. et l. c.*

*H. riparium* *L. Sp. pl.* p. 1129 (1753); *Hedw. Descr. IV*, p. 7, t. 3 (1793); *Brid. Bryol. univ. II*, p. 412; *C. M. Syn. II*, p. 321 (1851).

*H. riparium* { *var. homomallum* }  
                  { *var. oligophyllum* } *Brid. op. cit.* pp. 413-415.  
                  { *var. pallidum* }

*H. uncinatum* *var. plumosum* *Mac. Canad. M. n.* 371.

*Skitophyllum Weberi* *la Pyl. in Journ. bot. Desr.* 1813, V, p. 62.

*Stereodon riparius* *Mitt. in Journ. of the Linn. Soc.* 1869, p. 569.

1. *Ad saxa, ligna fabref. et putresc.*, terr. humid. — *Eur.* : Tota usq. in arctic. *As.* : Japon., Tonkin., Tibet. occ. *Afr.* : Alger., Madeir., ins. Canar., Kerguelen. *Am. sept.* : Groenland., Terr. nov., Canada merid., reg. sept. occid., Alberta, Wisconsin, M. Rupestr., Columb. brit., ins. Vancouver, tota temper., Ludovic., Cuba. *Pac.* : Austral. or. (Victoria).

« *var. abbreviatum* *Bryol. eur. l. c. t.* 571; *W. P. Sch. op. cit.* p. 718.

*Hypnum riparium* *var. abbreviatum* *Lesq. et Jam. Man.* p. 377.

*In siccior.* — *Eur. et Am. sept.* : Hic illic c. typ.

*var. abbreviatum* *de Not.* — *A. Haussmanni.*

« *var. angustifolium* *Warnst. in Verh. d. bot. Ver. d. Prov. Brandenb.* 1899, p. 78.

— *Eur.* : Lusatia, inf.

« *var. brachythecioides* *Schiffn. in Oesterr. bot. Zeitschr. n. II*, p. 24 *Sep. Abdr.* (1896).

*In herbos.* — *Eur.* : Bohem.

*var. curvipes* *R. du B.* — *A. curvipes.*

« *var. elongatum* *Bryol. eur. et W. P. Sch. ll. cc.*; *R. du Buyss. op. cit.* p. 19.

*A. riparium* *var. splendens* *de Not. in Erb. critt. ital. n.* 161 *et Epil.* p. 116 (1869).

*Hypnum riparium* *var. distichum* *Boul. M. Fr.* p. 76 p.p. (1884).

*In udis.* — *Eur.* : Hic illic c. typ. *As.* : Japon.

« *var. Felisii* *de Not. Cron. II*, p. 25 (1867); *W. P. Sch. et R. du Buyss. ll. cc.*

*Ad arb. rad. in humid.* — *Eur.* : Ut praeced.

- *var. flaccidum* (Lesq. et Jam.) Ren. et Card. *M. Am. sept.* p. 58.  
Hypnum riparium *var. flaccidum* L. et J. Man. p. 377.  
Paludic. et in stagn. — *Am. sept.* : Reg. sept.
- *var. floridanum* Ren. et Card. in *Bot. Gaz.* 1889, p. 98.  
A. floridanum *eor. in Rev. bryol.* 1893, p. 24.  
Hypnum Siphon *P. B. Prodr.* p. 70 (1805); *Brid. Bryol. univ. II*, p. 391 (1897) *fid. Cardot.*  
Lignic. et cortic. — *Am. sept.* : Florid., Ludovic.
- *var. fluitans* (Lesq. et Jam.) Ren. et Card. *M. Am. sept.* p. 58.  
Hypnum riparium *var. fluitans* L. et J. op. et l. cc.  
Aquat. — *Am. sept.* : Ontario, lac. Super., Assiniboine, Columb. brit., reg. sept.
- *var. genuense* de Not. *Cron.* p. 25 (1867).  
Ad aquaed. — *Eur.* : Ital. sept. (Genua).  
*var. Haussmanni* de Not. — A. Haussmanni.  
*var. hygrophilum* Bryhn. — A. hygrophilum.
- *var. inundatum* W. P. Sch. et R. du Buyss. *II. cc.*  
In stagn. — *Eur.* : Hic illic c. typ.  
*var. Kochii* R. du B. — A. Kochii.
- *var. longifolium* (Schultz) W. P. Sch. et R. du Buyss. *II. cc.*  
Hypnum longifolium *Schultz Fl. Starg.* p. 335 (1806).  
H. riparium ♀ petrophilum *Brid. Sp. M. II*, p. 115 (1812).  
H. riparium ♂ longifolium *ej. Bryol. univ. II*, p. 414 (1827).  
In humid. praecip. ad ligna robusta. — *Eur.* : Hic illic c. typ. As. :  
Jenissei sup. et med. *Am. sept.* : Washington, ins. Vancouver.
- *var. longinerve* Card. et Ther. in *Bot. Gaz.* 1900, p. 22.  
Aquat. — *Am. sept.* : Arkansas.
- *var. pratense* (Brid.) Par. *Ind.* p. 20 (1893).  
Hypnum riparium *var. pratense* *Brid. Bryol. univ. II*, p. 415 (*forsan spec. propr.*).  
— *Am. sept.* : Terra nov.
- *var. serratum* Ren. et Card. in *Bot. Gaz.* 1889, p. 98.  
Ad arb. — *Am. sept.* : Kansas.
- *var. serrulatum* Warnst. in *Verh. d. K. K. zool. bot. Gesells. in Wien*, 1900, p.  
Muric. — *Eur.* : Tirol. merid.  
*var. splendens* de Not. — A. riparium *var. elongatum*.
- *var. stagnorum* M. T. Lge. in *Bot. Tidsskr.* 1869, p. 36.  
— *Am. sept.* : Dania.
- *var. subsecundum* *Bryol. eur.*, W. P. Sch. et R. du Buyss. *II. cc.*  
In siccor. — *Eur.* : Hic illic c. typ.
- *var. tenue* Jur. in *Rab. Bryoth. eur. n.* 1245.  
Ad. rad. *Betul.* in turfo. — *Eur.* : Salisb.

« *var. trichopodium* (Schultz) Bryol. eur., W. P. Sch. et R. du Buyss. op. et ll. cc.

A. tenuisetum Lindb. M. Scand. p. 32 (1879).

A. trichopodium C. Hartm. Skand. Fl. X ed., II, p. 19 (1871); Braithw. Brit. Mossfl. III, p. 30 excl. var. ♂ (1896).

Hypnum riparium γ Schultzii Fiedl. Syn. Laubm. Mecklenb. p. 111 (1844).

Hypnum riparium ε trichopodium Brid. Bryol. univ. II, p. 415 (1827).

H. Schultzii Stand. M. exs. fasc. III, n. 150 (1804).

H. tenuisetum Lindb. in Hartm. Skand. Fl. IX ed. (1864).

H. trichopodium Schultz Fl. Starg. p. 324 (1806).

In prat. uligin. — Eur. : Rariss. : Mecklenb., Brandeb. Am. sept. : Minnesota.

*rivicola* (Mitt.) Jaeg. Ad. II, p. 549 (1877-78).

Hypnum rivicola Mitt. M. Ind. or. p. 83 (1859).

l. — As. : Tibet occ.

*rotulare* Lindb. — Hypnum alpestre.

*Rotae* de Not. — Hypnum exannulatum var. *Rotae*.

*saxatile* W. P. Sch. — A. hygrophilum.

*Schlotthaueri* Ren. et Card. in Bot. Centralbl. 1890, n. 51 (ut subsp.

A. serpentis).

l. — Am. sept. : Yellowstone Park.

*Sendtneri* } var. *giganteum* de Not. — Hypnum Wilsoni.

} var. *hamifolium* Bomans. et Broth. — Hypnum hamifolium.

*schenisianum* C. M. in Nuov. Giorn. bot. ital. 1898, p. 205.

l ? — As. : China centr. (prov. Schen-Si).

*serpens* (L.) Bryol. eur. fasc. 55-56 (Vol. VI), Mon. p. 9, t. 564 (1853); W. P. Sch. et R. du Buyss. op. cit. pp. 709 et 17.

Bryum repens Gmel. Syst. nat. II, p. 1333 (1768).

Gymnostomum serpens Schrank Baier. Fl. II. p. 439 (1789).

Hypnum capillare Froehl. in hb. Bridel.

H. clavellatum Schrank op. cit. p. 431.

H. contextum Hedw. Sp. M. p. 273, t. 72, f. 5-12 (1801).

H. intextum Voit M. Herbisp. p. 106 (1812) et in Sturm Deutsch. Fl. II, p. 11 p.p. (1892-20).

H. repens Poll. Pl. Palat. III, n. 1051, γ. II (1777).

H. repens trichodes terrestre minimum, capitulis majusculis oblongis erectis Dill. Cat. Giss. p. 216 (1718).

H. serpens L. Sp. pl. p. 1130 (1753); C. M. Syn. II, p. 411 (1851).

H. spinulosum Hedw. op. cit. p. 269, t. 69, f. 5-10.

H. subtile (non Hedw.) Dicks. Pl. crypt. fasc. IV, p. 17 (1801).

H. trichodes serpens, setis et capsulis longis erectis Dill. Hist. M. p. 329, t. 42, f. 61 (1741).

H. viride Lam. et DC. Fl. fr. p. 536 (1805).

H. Stereodon serpens (incl. ? var. aurantiacus, contextus, repens, spinulosus, viridis (1), brachyclados, byssoides, clavatus, crassinervius, falciformis, patentissimus, pensilvanicus, subtilis (2), crispabilis, cylindricus, excentricus, macrocarpus (3) Brid. Bryol. univ. II, pp. 642-649.

Neckera serpens Willd. Prodr. fl. Berol. n. 935 (1787).

Trichostomum repens Schrank op. cit. p. 450.

(1) Ad synonyma supradicta referenda.

(2) « Quorum plures hujus speciei non sunt ». W. P. Sch. Syn. I ed. p. 592.

(3) Fide Schumann Fl. Sael. II, p. 83.

1. Ad saxa, lign. putresc. arb. rad., rarius ad terr. in toto fere terr. orbe. — *Eur.* : Tota, rarius in calidior. et in arctic. *As.* : Japon., Baltistan, Kashmir, Tehitral, Tibet occ. *Afr.* : Alger., etc. *Am. sept.* : Groenland, Labrador, Canada tot., Columb. brit., reg. sept., centr., occ. et or. *Am. merid.* : And. Quitens. et Peruv. *Pac.* : Nov. Seeland. sept.
- « *var. angustifolium* Har. Lindb. in *Limpr. Laubm. Deutschl.... III*, p. 321 (1897).  
Lignic. — *Eur.* : Lappon., Suec.
- « *var. Columbiae* Kindb. in *Bull. Torr. bot. Club, XVII*, p. 279 (ut subspec.) (1890).  
— *Am. sept.* : Columb. brit.  
*var. cryptartum* Arn. — *A. densum*.
- « *var. depauperatum* (Boul.) Braithw. *Brit. Mossfl. III*, p. 25 (1896).  
*A. Cashii* R. du Buyss. *Essai*, p. 8 (1889); *Par. Ind.* p. 12 (1894).  
*Hypnum serpens var. depauperatum* Boul. *M. Fr. I*, p. 19 (1884).  
Ad arb. rad. et terr. arenos. — *Eur.* : Anglia; hic illic.  
*var. Juratzkanum* R. du B. — *A. Juratzaknum*.
- « *var. major* R. du Buyss. *op. et l. cc.*  
Ad rad. arb. — *Eur.* : Hic illic.  
*var. orthocladon* Aust. — *A. orthocladon*.
- « *var. pinnatum* W. P. Sch. *op. et l. cc.*  
Ad rup. arenac. — *Eur.* : Sudet.  
*var. radicale* Aust. — *A. radicale*.  
*var. rigidiusculum* Lindb. et Arn. — *A. rigescens*.
- « *var. rigidum* T. Jens. *Bryol. dan. p.* 192 (1856).  
— *Eur.* : Dania.
- « *var. serrulatum* Breidl. in *Oesterr. bot. Zeit.* 1889; *Limpr. op. cit.* p. 321.  
— *Eur.* : Stiria. *As.* : Caucas.
- « *var. subnervae* Ren. et Card. in *Bot. Gaz.* 1900, p. 22.  
— *Am. sept.* : Terra nov.
- « *var. tenue* (Schräd.) *Bryol. eur. et W. P. Sch. II. cc.*  
*Hypnum serpens var. tenue* Brid. *Musc. rec. II, P. II*, p. 114 (1801).  
*H. tenue* Schräd. *Krypt. Gew. n.* 80 (1796).  
*H. Stereodon serpens var. tenue* Brid. *Bryol. univ. II*, p. 616 (1827).  
Ad arb. trunc. — *Eur.* : Reg. montana.
- « *var. xanthodictyon* Kindb. in *Mac. Cat. p.* 218 (1892).  
*A. serpens* \* *lutescens* ej. mss.  
Ad saxa madid. — *Am. sept.* : Anticosti.
- « *var. β* (Hook. f. et Wils.) *Par. Ind. p.* 21 (1893).  
*Hypnum serpens var. β* H. f. et W. *Fl. antarct. p.* 138 (1844) et *Handb. of the N. Zeal. Fl. p.* 479 (1867).  
— *Pac.* : Ins. Auckland.  
*serpens* \* *lutescens* Kindb. — *A. serpens var. xanthodictyon*.  
*serratum* *Bryol. eur.* — *A. compactum* ?  
*sinensi-subtile* C. M. in *Nuov. Giorn. bot. ital.* 1896, p. 123.  
1. — *As.* : China centr. (prov. Schen-Si.)

*Smithii* Lindb. — *Hypnum arcticum*.

*sparsifolium* (Hpe) *Jaeg. Ad. II*, p. 552 (1877-78).

*Serpo-Hypnum sparsifolium* Hpe *Symb.* 1870, p. 290.

l. Lignic. — *Am. merid.* : Brasil. austr. or.

*sparsile* (Mitt.) *Jaeg. op. cit.* p. 354.

*Hypnum sparsile* Mitt. *M. Ind. or.* p. 82 (1839).

l. — *As.* : Assam super.

*speirophyllum* Kindb. in *Olt. Nat. IV*, p. 64 (1889) et in *Mac. Cat.* p. 217 (1892).

2 ? *Ster. Rupic.* — *Am. sept.* : New Brunswick.

*Sprucei* (Bruch) *Bryol. eur. fasc.* 55-56 (Vol. VI), *Mon.* p. 5, t. 56i (1853); *W. P. Sch. et R. du Buyss. op. cit.* pp. 705 et 5.

*Hypnum confervoides* H. f. et W. in *Drumm. M. bor. am.* 1 ed. n. 190.

*H. Jungermannia* Hpe in *Sched.*

*H. jungermannoides* Brid. *Sp. M. II*, p. 225 (1812) *fid. Limpr.*

*H. Sprucei* (Br.) *Spruce in Lond. Journ. bot. II*, p. 180 (1845) et *M. Pyr. exs.* n. 62 (1847); *C. M. Syn. II*, p. 415 (1851).

*Leskea Sprucei* Bruch. in *litt.*

*L. tenella* W. P. Sch. in *Drumm. M. bor. am.*

*Platydicta Sprucei* Berk. *Handb. brit. M.* p. 145 (1863).

*Stereodon Sprucei* Lindb. in *Broth. Enum. M. Cauc.* p. 129 (1892).

2. Ad rup. umbr. humid. — *Eur.* : Spitzberg, ins. Ursor., reg. mont., subalp. et alp. Scand., Britann., Westphal., Tatra, Salisb., Stir., Carinth., Bavar., Tirol., Helvet., Gall. centr. (Arvern.), l'yren. *As.* : Penins. Tschutschica, Jenissei, Obi, Caucas. centr. *Am. sept.* : Groenland, Canada mer. occ., Massachussets, Wisconsin, Montana, M. Rupestr., Columb. brit., ins. Vancouver, Washington, Idaho, Colorado, Nov. Mexic.

α *var.* *Anzianum* (de Not.) *R. du Buyss. op. cit.* p. 7.

*A. Anzianum* de Not. *Epil.* p. 156 (1869).

— *Eur.* : Longobard.

*subcompactum* C. M. et Kindb. in *Mac. Cat.* p. 221 (1892).

*Hypnum compactum* Mac. *Canad. M.* p. 324.

l. Ad rup. et arb. rad. — *Am. sept.* : Niagara, Montana, M. Rupestr., Columb. brit.

*subserpens* W. P. Sch. in *Savat. M. Jap.* n. 709.

*Thuidium capillatum* (fid. E. S. Salmon).

l. Ad terr. calcar. — *As.* : Japon.

*subsulcatum* de Not. — *Hypnum sulcatum var. subsulcatum*.

*subtile* (Hedw.) *Bryol. eur. fasc.* 55-56 (Vol. VI), *Mon.* p. 4, t. 561 (1853); *W. P. Sch. et R. du Buyss. op. cit.* pp. 706 et 7.

*Hypnum*: *minimum*, foliis ovato-lanceolatis, capitulis erectis, operculis conicis Hall. *Hist. Stirp. Helvet.* n. 1757 (1761).

*H. subtile* Hoffm. *Deutsch. Fl. II*, p. 70 (1796); *C. M. Syn. II*, p. 415 (1851).

*H. Stereodon serpens var. subtilis* Brid. *Bryol. univ. II*, p. 649 (1827).

*Leskea subtilis* Hedw. *Descript. IV*, p. 23, t. 9 (1793).

*Neckera subtilis* Brid. *Musc. rec. II, P. II*, p. 45 (1801).

1. Ad arb. silvat. praecip. *Fagos*, rar. ad. saxa. — *Eur.* : Reg. mont. et subalp., rar. campestr.; deest in Britann. *As.* : Caucas. occ. et centr., Kashmir. *Am. sept.* : Ottawa, Ontario, reg. sept. or., Wisconsin, Minnesota.

var. *tenuissimum* Limpr. — *A. tenuissimum*.

*tenuifolium* Kindb. — *A. fenestratum*.

*tenuisetum* Lindb. — *A. riparium* var. *trichopodium*.

*tenuissimum* (Gümb.) *Bryol. eur. fasc.* 55-56 (*Vol. VI*), *Mon. p.* 7, t. 562 (1853); *W. P. Sch. et R. du Buyss. op. cc. pp.* 707 et 7.

*A. subtile* var. *tenuissimum* Limpr. *Laubm. Deutschl...* III, p. 301 (1897).

1. Cortic. — *Eur.* : Fennia, Bavar. (Monach.) *As.* : Tibet occ.

thuidioides *W. P. Sch. in Savatier M. Japon. n.* 720.

1. Terr. — *As.* : Japon.

tibetanum (*Mitt.*) *Par. Ind. p.* 23 (1893).

*Hypnum tibetanum* Mitt. *M. Ind. or. p.* 83 (1859).

Ster. — *As.* : Tibet. occ.

? torrentium *Besch. in Bull. Soc. bot. Fr.* 1898, p. 125.

Ster. Ad rup. irror. — *Pac.* : Taïti.

*trichopodium* C. Hartm. — *A. riparium* var. *trichopodium*.

var. *Kochii* Lindb. — *A. Kochii*.

*uncinatum* Mitt. (1896). — *Hypnum austro-uncinatum*.

*vacillans* Sull.  *Ic. M. Suppl. p.* 96, t. 72 (1874).

*Hypnum vacillans* Lesq. et Jam. *Man. p.* 377 (1881).

1. Aquat. — *Am. sept.* : New Jersey, M. Alb., Ontario.

*Vallis Clausae* Husn. — *A. fallax*.

var. *atro-virens* Husn. — *A. fallax* var. *nigricans*.

var. *spinifolium* Husn. — *A. fallax* var. *spinifolium*.

varium (*Hedw.*) Lindb. *M. scand. p.* 32 (1879).

*A. radicale* *Bryol. eur. fasc.* 55-56 (*Vol. VI*), *Mon. p.* 10, t. 565 (1853).

*Hypnum debile* Brid. *Sp. M. II, p.* 250 (1812).

*H. humile* *P. B. Prodr. p.* 68 (1805) (1).

*H. inordinatum* Brid. *Sp. M. II, p.* 182 et *Bryol. univ. II, p.* 483 (1827) (1).

*H. pachypoma* *Schwaegr. Suppl. I, P. II, p.* 259 (1816) et III, *P. II, II, t.* 287 (1829-30); *Brid. Bryol. univ. II, p.* 538 1827 (1).

*H. radicale* (non *P. B.*) Wils. *Bryol. brit. p.* 365 t. 25 (1855).

*H. serpens* β varium *C. M. Syn. II, p.* 412 p.p (1851).

*H. varium* *P. B. Prodr. p.* 72 (1805).

*H. Stereodon* varium *Brid. Bryol. univ. II, p.* 652.

*Leskea varia* *Hedw. Sp. M. p.* 216, t. 53, f. 15-20 (1801).

*Rigodium varium* Lindb. *Laubm. Schwed. u. Norw. p.* 14 (1883).

*Stereodon* varium Mitt. in *Proced. of the Linn. Soc.* 1864, p. 43.

1. Ad terr., rup. et rad. arb. in humid. — *Eur.* : German. sept., Siles., Bohem., Austr. infer., Hungar., Stir., Carinth., Tirol., Helvet., vall. Rhen. med. *As.* : Jenissei med., Pers., Caucas. occ. et centr. *Afr.* : Madeir. *Am. sept.* : Anticosti, ins. Miquelon, Canada merid., Columb. brit, Washington, Idaho, Montana, reg. sept., centr. or. et merid. *Am. merid.* : And. Quitens ?

(1) Forsan ad *A. orthocladon* referenda. — De syn. *A. radicalis* cf. Lindb. *M. scand. p.* 32, *R. du Buyss. in Rev. de bot.* IV, n. 42-43 et Limpr. *Laubm. Deutschl...* III, pp. 316 et 323.



- varium* R. du Buyss. — *A. oligorhizon*.  
 var. *radicale* R. C. — *A. radicale* var. *Lesquereuxii*.  
 var. *oligorhizon* Lindb. — *A. oligorhizon*.  
*vernicosum* var. *majus* Lindb. — *Hypnum vernicosum* var. *turgidum*.  
*viridulum* Lindb. — *Hypnum norvegicum*.  
*Wilsoni* Lindb. — *Hypnum Wilsoni*.  
 var. *hamatum* Lindb. — *Hypnum hamifolium*.  
*Wollei* Kindb. — *Leskea*.

*yesoanum* Ren. et Card. in *Faurie Pl. Jap. n.* 8622.  
 Ster. Aquat. ? — As. : Japon. (Iac Kushiro).

**Amblytropis** Mitt. *M. austr. am.* p. 339 (sect. *HOOKERIE*) (1869).

**Amphidiopsis** C. M. in Linn. 1878-79, p. 332 (sect. *BARBULÆ*).

**AMPHIDIUM** Nees in Sturm. *Deutsch. Fl. II*, p. 17 (1819); *C. M. Syn. I*, p. 667 (1849) et *II*, p. 636 (1851) (sect. *ZYGODONTIS*); *W. P. Sch. Coroll.* p. 39 (1856) (1)

- lapponicum* W. P. Sch. — *Amphoridium*.  
*Mougeotii* W. P. Sch. — *Amphoridium*.  
*pulvinatum* Nees. — *Zygodon Forsteri*.

**AMPHORIDIUM** W. P. Sch. *Syn. I* ed. p. 247 (1860). — 16 spec.

- africanum* C. M. in *Rehm. M. Afr. austr. n.* 143.  
 Ster. — Afr. : Orange.

*anoëctangioides* (C. M.) Broth. *Natür. Pflanzenfam.* p. 460.

- Zygodon anoëctangioides* C. M. in *Fl.* 1896, p. 449; *Par. Ind.* p. 1371 (1898).  
 Ster. — Pac. : Ins. Hawai.

*caespitosum* (Mitt.) Jaeg. *Ad. I*, p. 385 (1872-73); *Lesq. et Jam. Man.* p. 160 (1864).

- Didymodon caespitosus* Mitt. in *Journ. of the Linn. Soc.* 1864, p. 18.  
 Ster. ? Rupic. — Am. sept. : Columb. brit., ins. Vancouver.

*californicum* (Hpe) Jaeg. *op. et l. cc.*; *Lesq. et Jam. op. cit.* p. 159.

- A. Sullivantii* Mac. *Canad. M. n.* 111.  
*Zygodon californicus* Hpe in *Bot. Zcit.* 1862, p. 361; *Sull. Ic. M. Suppl.* p. 47, t. 82 (1874).

2. Rupic. — Am. sept. : Columb. brit., ins. Vancouver, Washington, Calif.

*canariense* W. P. Sch. — *A. curvipes*.

*compactum* (C. M.) *Par. Suppl. Ind.* p. 7 (1900).

- Zygodon compactus* C. M. in *Hedw.* 1898, p. 134.  
 Ster. — Pac. : Nov. Seeland. austr.

*crispatum* (Kindb.) *Par. op. et l. cc.*

- Zygodon crispatus* Kindb. in *Rev. bryol.* 1896, p. 21.  
 Rupic. — Am. sept. : Columb. brit.

*curvipes* (C. M.) Jaeg. *Ad. I*, p. 386 (1872-73).

- A. canariense* W. P. Sch. in *Mandon Pl. Madeir. n.* 13.  
*Anoëctangium lapponicum* Kze. in *Holl. Pl. Madeir. in Fl.* 1830.  
*Rhabdoweisia curvipes* Mitt. in *Godman's Nat. hist. of Azor.* p. 291 (1870).  
 2. Rupic. — Afr. : Madeir.

---

(1) Cf. Limpr. *Laubm. Deutschl. .... II*, p. 5.

cyathicarpum (Mont.) Jaeg. op. et l. cc.

Didymodon cyathicarpus Mitt. in Journ. of the Linn. Soc. 1859, p. 70.  
Gymnostomum linearifolium Tayl. in Lond. Journ. of bot. 1846, p. 42.  
Oncophorus cyathicarpus Mitt. in Voy H. S. M. Challenger, Bot. III, p. 78 ( ).

Rhabdowisia cyathicarpa ej. in Journ. of the Linn. Soc. XII, p. 54.  
Syrrophodon tortuosus Hsch. in Linn. 1811, p. 117.  
Zygodon cyathicarpus Mont. in Ann. sc. nat. 1845, p. 106 et Syll. p. 37 (1856); C. M. Syn. I, p. 682 (1849).

1. Terr. — Afr. : Abyss., Kihina N'jDaro, Cameroon, Pr. B. Sp. Am. sept. : (fid. Handb. of the N. Zeal. Fl., absque loco). Am. merid. : M. Pichincha, Chile, ins. Juan Fernandez. Pac. : Austral. or. (Victoria), Nov. Seeland., Tasmaan.

gracile de Not. — Zygodon.

integrifolium (C. M.) Par. Suppl. Ind. p. 8 (1900).

Zygodon integrifolius C. M. in Trans. and Proceed. of the N. Zeal. Inst. 1892, p. 192, t. XXXVII et in Hedwig. 1898, p. 133.

2. Rupic. — Pac. : Nov. Seeland. austr.

kilima n'djaricum (C. M.) Par. Ind. p. 24 (1893).

Zygodon kilima n'djaricus C. M. in Fl. 1890, p. 482.

Ster. — Afr. : Kilima N'Djaro.

lapponicum (Hedw.) W. P. Sch. op. et l. cc.

Amphidium lapponicum W. P. Sch. in Bryol. eur. Consp. III, p. 3 (1855) et Coroll. p. 39 (1856).

Anictangium lapponicum Hedw. Sp. M. p. 40 (1801).

A. striatum Brid. Sp. M. I, p. 25 (1806).

Bryum lapponicum Dicks. Pl. crypt. fasc. IV, p. 10 (1801).

Br. surculo filiformi ramoso, foliis subulatis, capsulis piriformibus sessilibus, operculis arcuatis conicis Ström in Skr. af Nat.-hist. Selsk. 1, 2, p. 33 et t. XI, n. 3 (1791) fid. Sw.

Cylicocarpus lapponicus Lindb. in Oefv. af Vet.-Akad. Foerh. 1862, n. 10.

Didymodon lapponicus Mitt. in Journ. of the Linn. Soc. 1864, p. 18.

Gymnostomum lapponicum Hedw. M. frond. III, p. 10, t. 5\* (1792).

G. striatum Roehl. Deutsch. Fl. III, p. 37 (1813).

Hedwigia lapponica Hedw. op. cit. p. 31 animadv.; Brid. Musc. rec. II, p. 33 (1788).

Hymenostylium lapponicum Funck in Bot. Zeit. I, 1837.

Schistidium striatum Brid. Mant. M. p. 22 (1819).

Zygodon lapponicus Bryol. eur. fasc. 4, (Vol. III), Mon. p. 6, t. 206 (1839).

1. In fss. rup.; calcar. fugit. — Eur. : Reg. subalp. usq. in supraalp. : Spitzberg. As. : Penins. Tschutschica, Kashmir, Tchitral, Gilgit, Caucas. centr. Am. sept. : Groenland, Anticosti, S. W. Canada, reg. sept., Wyoming, Idaho, M. Rupestr., Columb. brit., ins. Vancouver, Washington, Calif.

Mougeotii (Bryol. eur.) W. P. Sch. op. cit. p. 248.

Amphidium Mougeotii W. P. Sch. op. cit. p. 40.

Anoetangium Mougeotii Lindb. M. scand. p. 29 (1879).

Barbula Woodii W. P. Sch. Syn. II ed. p. 207 (1876).

Cylicocarpus Mougeotii Lindb. in Oefv. af Vet.-Akad. Foerh. 1862, n. 10.

Gymnostomum Mougeotii Bruch mss.; Hüb. Muscol. germ. p. 59 (1333).

Hymenostylium Mougeotii Garov. Bryol. austr. exc. p. 18 (1840).

Zygodon Mougeotii Bryol. eur. II, cc. p. 7 t. 206.

2. In fss. rup.; calcar. fugit. — Eur. : Vulgatiss., rariss. fructif., copiose, in Arvernica; Scot., Salisb., etc. As. : Caucas. Am. sept. : Fret. Scoresby, prom. Breton, m. Alb., Pensilv., Idaho, Alaska, m. Rupestr., Columb. brit., ins. Vancouver, Washington, Oregon.

*Peckii* Sull. — *Anoëctangium*.

*remotidens* (C. M.) Par. *Suppl. Ind.* p. 8 (1900).

*Zygodon remotidens* C. M. in *Hedwig*. 1898, p. 133.

1. — *Pac.* : Austral. or. (Victoria).

*rupestre* de Not. — *Zygodon viridissimus* var. *rupestris*.

*subcyathicarpum* (C. M.) Par.

*Zygodon subcyathicarpus* C. M. in *Rehm. M. Afr. austr.* n. 146; *Par. Ind.* p. 1378 (1898).

— *Afr.* : Prom. B. Sp.

*sublapponicum* (C. M.) Par. *op. et l. cc.*

*Zygodon sublapponicus* C. M. in *Nuov. Giorn. bot. ital.* 1898, p. 86.

1. — *As.* : China centr. (prov. Schen-Si).

*Sullivantii* (C. M.) *Lesq. et Jam. Man.* p. 159.

*Syrrophodon* ? *excelsus* Sull. *M. Ailegh. n.* 170 (1816).

*Zygodon Sullivantii* C. M. *Syn. I.* p. 679 (1849); *Sull. M. of U. St.* p. 32 (1856) et *l. c.* *M. I.* p. 51, t. 52 (1864).

*Ster. Cortic. et rupic.* — *Am. sept.* : Prom. Breton, Carol. super.

*Sullivantii* Mac. — *A. californicum*.

*viridissimum* de Not. — *Zygodon*.

**AMPHORITHECA** Hpe. in *Ann. sc. nat. V. ter. III*, p. 340 (1866).

*Spec. omn. non indicatae sunt ENTOSTHODONTES ejusd. nom.*

*Jamesoni* Hpe. — *Physcomitrium*.

*ramulosa* Hpe. — *Funaria*.

*riparia* Aongstr. — *Funaria*.

**Amphorithecium** C. M. in *Hedwig*. 1898, p. 259 (*nom.*, sect. CUPRESSINÆ = ECTROPOTHECIUM).

**Ampullaria** C. M. *Syn. I.* p. 143 (1849) (sect. SPLACHNI).

**ANACALYPTA** Roehl. *Moosgesch. Deutschl. I.* p. 108 (1800); *C. M. Syn. I.* p. 547 (1849) et *II*, p. 621 (1851) (sect. POTTIÆ).

*Spec. omn. non indicatae sunt POTTIÆ ejusd. nom.*

*affinis* Fürnr. — *Pottia intermedia*.

*lanceolata* Roehl. var. *sibirica* E.G. et G.G. Bor. — *Pottia latifolia* ?

*pilifera* Bruch. — *Pottia latifolia* var. *pilifera*.

*recurvirostris* N. et H. — *Didymodon rubellus*.

*reflexa* Fürnr. — *Gyroweisia*.

*rubella* Hüb. — *Didymodon*.

*tophacea* Bruch. — *Trichostomum*.

**ANACAMPTODON** Brid. *Mant. M.* p. 136 t. 2, f. II (1819). — 4 spec.

*cubensis* (Sull.) *Mitt. M. austr. am.* p. 540 (1869).

*A. maynensis* Spr. *Cat.* p. 18 (1867).

*Fabronia cubensis* Sull. in *Proceed. of the Amer. Acad.* 1861.

1. *Cortic.* — *Am. sept.* : Cuba. *Am. merid.* : Amazon., And. Peruv.

*fabronioides* Welw. et Duby. in *Mem. Genève...* 1871, p. 2, t. I, f. 5.

1. *Cortic.* — *Afr.* : Angola.

Fortunei Mitt. in Journ. of the Linn. Soc. 1864, p. 152.

— As. : China or.

maynensis Spr. — A. cubensis.

pygmaeus Bryol. jav. — Schwetschkea.

splachnoides (Froehl.) Brid. Mant. M. II. cc. et Bryol. univ. II, p. 225; Bryol. eur. V, t. 453; W. P. Sch. Syn. II ed. p. 536 (1876).

Campylocladum hypnoides Schwaegr. Suppl. III, P. II, fasc. I, t. 211 (1827).

Fabronia splachnoides C. M. Syn. II, pp. 38 et 662 (1851).

Neckera splachnoides Schwaegr. Suppl. I, P. II, p. 151, t. 82 (1816).

Orthotrichum splachnoides Froehl. mss. in Brid. Sp. M. II, p. 4 (1812).

1. Ad Fagos, rar. Abiet. putresc. — Eur. : Voges., Silva nigr., Tatra, Siles., Austria, Hungar., Salisb., Stir., Carinth., Bavar., Tirol., Helvet., Ticin., Pyren. ; deest in Britann. As. : Caucas., occ. Am. sept. : Reg. or.

**Anacamptophyllum** C. M. Syn. II, p. 439, (subsect. HYPNI) (1851).

**Anacamptosphaeum** C. M. in Linn. 1875, p. 547 (nom., sect. SPHAGNI).

**ANACOLIA** W. P. Sch. Syn. I ed. p. 421 (1860). — 7 spec.

abyssinica W. P. Sch. in W. Sch. M. Abyss. et Syn. II ed. p. 514 (1876).

Bartramia abyssinica C. M. Syn. I, p. 506 (1849).

2. Rupic. — Afr. : Abyss., Choa.

Baueri (Hpe) Par. Ind. p. 27 (1893).

Bartramia Menziesii (fid. Lesq. et Jam.).

Bartramia Menziesii var. Baueri R. C. in Rev. bryol. 1892, p. 94.

Glyphocarpus Baueri Hpe in Linn. 1859 p. 457.

2. Terr. — Am. sept. : Calif.

didymocarpa (W. P. Sch.) Par. Suppl. Ind. p. 8 (1900).

Bartramia didymocarpa W. P. Sch. in Mandon Pl. Boliv. n. 1669 ; C. M. in Nuov. Giorn. bot. ital. 1897, p. 55 ; Par. Ind. p. 105 (1893).

— Am. merid. : And. Boliv.

intertexta (W. P. Sch.) Jaeg. Ad. II, p. 699 (1877-78).

Bartramia intertexta C. M. Syn. I, p. p. 503 (1849).

B. lamprocarpa W. P. Sch. in hb. Mus. Paris.

B. setifolia Mitt. M. austr. am. p. 269 p.p. (1869).

Glyphocarpa intertexta W. P. Sch. in Besch. Pr. bryol. mexic. p. 58 (1871).

2. — Am. sept. : Mexic.

Menziesii (Turn.) Par. Ind. p. 27 (1893).

Bartramia Menziesii Turn. in Kon. et Sims. Ann. bot. I, p. 525, t. II, f. I ( ) ; Brid. Bryol. univ. II, p. 48 (1827) ; C. M. Syn. I, p. 505 (1849) ; Lesq. et Jam. Man. p. 204 (1884).

Glyphocarpa Menziesii Jaeg. Ad. I, p. 525 (1873-74).

Philonotis leiophylla Kindb. mss. ; Mac. Canad. M. n. 151.

2. Rupic. — Am. sept. : Alaska, Columb. brit., ins. Vancouver, Washington, Idaho, Oregon, Calif.

setifolia (Hook.) Jaeg. Ad. II, p. 699 (1877-78).

Bartramia intertexta C. M. Syn. I, p. 503 p.p. (1849).

B. setifolia Mitt. M. austr. am. p. 268 (1869).

Glyphocarpa setifolia Jaeg. Ad. I, p. 525 (1873-74).

Gymnostomum setifolium Hook. Ic. pl. rar. II, p. 135 (1841).

2. Rupic? — Am. sept. : And. Quitens. et Peruv.

*Webbii* (Mont.) W. P. Sch. Syn. II ed. p. 513 (1876).

*Bartramia granatensis* W. P. Sch. Col. coll. p. 85, n. 7 (1856).

*B. Webbii* C. M. Syn. I, p. 504 (1849).

*Glyphocarpus Webbii* Mont. in Ann. sc. nat. 1838, p. 36, n. 97, Pl. cell. Canar. p. 28, t. 2. f. 2 (1840) et Syll. p. 29 (1856).

2. Ad terr. arid. — Eur.: Corsica, M. Granat. Afr.: Alger., Canar., Madeir.

**Anaglyphodon** Philib. in Rev. bryol. 1900, p. 19 (sect. BRYI).

**Anasmagonium** Mitt. M. austr. am. p. 615 (sect. POGONATI) (1869).

**Anastrephidium** Hpe in Linn. 1876, p. 271 (sect. POROTRICH).

**ANDREAEAE** Ehrh. in Hannov. Mag. 1778, p. 1601. — 89 spec.

*acuminata* Mitt. in Journ. of the Linn. Soc. 1859, n. 64. et in Fl. Tasman. II, p. 161, t. 7, f. 2. (1860).

*A. subappendiculata* C. M.?

Ster. Rupic. — Afr.: Ins. Kerguelen. Pac.: Tasman.

*acutifolia* Hook. f. et Wils. in Lond. Journ. of bot. 1844, p. 535, Fl. antarct. II, p. 396, t. 151, f. 2 (1847) et Handb. of the N. Zeal. Fl. p. 400 (1867); C. M. Syn. I, p. 7 (1849) et II, p. 516 (1851); Mitt. M. austr. am. p. 628 (1869).

— Afr.: Ins. Kerguelen. Am. sept.: Groenland, (steril.). Am. merid.: And Peruv? Fuegia, ins. Eremitae, Maclov. Pac.: Nov. Seeland. sept. (1), ins. Auckland, Campbell.

*alpestris* (Thed.) Bryol. eur. VI, Mon. p. 16, t. 626 (1855); W. P. Sch. Syn. II ed. p. 814 (1876).

*A. petrophila* var. *alpestris* Thed. in Nya Bot. Not. 1849, p. 79, fig. 45-47.

*A. petrophila* A heteromalla {  $\alpha$  f. intra alpina } Zeit. Mon. Andr. p. 42  
{  $\beta$  f. *alpestris* } (1855).

1. Ad saxa humid. praecip. secus rivul. glacial. — Eur.: Scot., Scandin., Boehmerwald, Riesengeb., Stir., Carinth., Albula, Voralberg., Helvet., Arvern., Longobard., Pyren. Am. sept.: Groenland, Nov. Scot.

*alpestris* var. Mitt. — *A. commutata*.

*alpina* (Dill., L.) Sm. Fl. brit. p. 179 excl. syn. nonnull. et Engl. Bot. t. 1278 (1804); Brid. Bryol. univ. II, p. 728 p.p. (1827); Bryol. eur. VI, Mon. p. 18, t. 628 (1855); C. M. Syn. I ed. p. 7 (1849) et II ed. p. 516 (1851); W. P. Sch. op. cit. p. 819.

*A. alpina* var. *compacta* Hook. in Trans. of the Linn. Soc. 1810, p. 589.

*A. alpina* var. *flavicans* ej. in op. et l. cc.

*A. petrophila* Ehrh. Beitr. I, p. 592 p. p. ( ).

*Jungermannia alpina*, surculis teretibus, foliis ovatis patentibus, calycibus

imbricatis L. Sp. pl. p. 1601 (1753).

*Lichenastrum alpinum* atro-rubens teres, calycibus squamosis Dill. Hist.

M. p. 506, t. 73, fig. 39 A-D (1741) et Hb.

1. Ad saxa irror. in rivul. — Eur.: Hibern., Alp. Scot., Cambrovall., Norveg., ins. Finmark; Arvern., summ. Alp.

1. De speciebus Brownianis Nov. Seelandicis *Andreaeaceae* 16 a nemine visis, imo bryologorum investigationis ab auctore denegatis, jam tum plus minus dubiis, cf. Rev. bryol. 1900, p. 49 et in adnot. ad gen. *Bryum*, *Campylopus*, *Dicranum*, *Grimmia*, *Orthotrichum*, etc.

- alpina* { Aongstr. — *A. obovata*.  
Brid. p.p., Hedw., W. et M. — *A. petrophila*.  
Mitt. — *A. pseudo-alpina*.
- var. *compacta* } Hook. — *A. alpina*.  
var. *flavicans* }
- var.  $\beta$  Wahlenb. — *A. obovata*.  
*ambigua* Wils. — *A. commutata*.
- amblyophylla* C. M. in sched.; Broth. in *Oefv. af Finska Vet.-Soc. Foerh.* 1895, p. 47; C. M. in *Hedwig.* 1898, p. 82.  
2. Rupic. — Pac. : Austral. or. (Nov. Vale\*), Tasman.
- angustata* Lindb. in *Limpr. Laubm. Deutschl.* ... I, p. 144 (1885).  
1. — Eur. : Stir., Arvern. (M. Dore).
- appendiculata* W. P. Sch. in *Bryol. eur. VI, Mon.* p. 5 (1855); C. M. in *Bot. Zeit.* 1864, p. 373; Mitt. *M. austr. am.* p. 629 (1869).  
*A. rupestris* H. f. et W. M. antarct.  
Ster. Rupic. — Am. merid. : Cap Horn., Fuegia, ins. Eremitae.
- aquatica* C. M. in *Hedwig.* 1898, p. 82.  
— Pac. : Nov. Seeland.
- arachnoidea* C. M. in *Linn.* 1878-79, p. 220.  
2. — Am. merid. : Alp. Argentin. subtropic.
- arctæoides* C. M. et N. Beckett in *Trans. and Proceed. of the N. Zeal. Inst.* p. 293 t. XLI et in *Hedwig.* 1898, p. 83.  
2. Rupic. — Pac. : Nov. Seeland austr.
- Arthuriana* C. M. in *Hedwig.* 1898, p. 74.  
2. — Pac. : Nov. Seeland austr.
- asperula* Mitt. in *Journ. of the Linn. Soc.* 1859, p. 65.  
*A. Mülleri* Sonder (fd. Mitt. in *Trans. and Proceed. of the roy. Soc. of Victoria*, 1883, p. 96.  
— Pac. : Alp. Austral.
- assimilis* C. M. in *Bot. Centralbl.* 1883, n. 2-4.  
2. — As. : Penins. Tschutschica.
- aterrima* C. M. in *Engler's Bot. Jahrb.* 1883, p. 96 et in *Forschungsreise...* Bot. p. 8 (1889).  
2. Ad rup. sicc. — Afr. : Ins. Kerguelen.
- attenuata* C. M. in *Hedwig.* 1898, p. 84.  
2? fr. ign. Rupic. — Pac. : Tasman. (M. Wellington).
- australis* F. v. Müll. in sched.; Mitt. in *Hooker's Journ. of bot.* 1856, p. 257.  
2. — Pac. : Alp. Austral. or.
- Blyttii* *Bryol. eur. fasc.* 52-54 (Vol. VI), *Mon.* p. 25, t. 63 (1855); W. P. Sch. *Syn. II ed.* p. 821 (1876).  
*A. perichaetialis* Zett. *Mon. Andr.* p. 26 (1855).  
2. Ad rup. juxta niv. — Eur. : Ins. Parry, Ursor., Spitzberg, Lapon., alp. Norveg. Am. sept. : Groenland.
- borbonica* Besch. *Fl. bryol. Réunion, etc.* p. 187 (1880-81).  
2? — Afr. : Ins. Borbon.

- brevipes* Spr. in *Proced. of the Linn. Soc.* 1859, p. ; Mitt.  
*M. austr. am.* p. 628 (1869).  
 Rupic. — *Am. merid.* : And. Nov. Granat., Quitens. et Peruv.
- cochlearifolia* C. M. in *Trans. and Proced. of the N. Zeal. Inst.* p. 293.  
 t. XXXIX et in *Hedwig.* 1898, p. 81.  
 Rupic. — *Pac.* : Nov. Seeland.
- commutata* C. M. in *Bot. Zeit.* 1864, p. 376.  
 A. *alpestris* var. Mitt. in *hb. Ind. or.* n. 3.  
 A. *ambigua* Wils. in *Kew Journ. bot.* IX, p. 289 (1857).  
 A. *petrophila* Mitt. *M. Ind. or.* p. 6 (1859).  
 2. — *As.* : Sikkim.
- commutata* Limpr. — A. Huntii.
- compacta* C. M. in *Bot. Centralbl.* 1883, n. 2-4.  
 1. — *As.* : Penins. Tschutschica.
- crassinervia* Bruch in *Denkschr. Akad. Münch.* 1828, p. 279, n. I, t. 10  
*Bryol. eur. fasc.* 52-54 (Vol. VI), Mon. p. 23, t. 633 (1855); C. M.  
*Syn.* I, p. 10 (1849); W. P. Sch. *op. cit.* p. 820.  
 A. *crassinervia* § *Heinemanni* C. M. *op. et l. cc.*  
 A. *Heinemanni* Hpe. et C. M. in *Bot. Zeit.* 1846, p. 364, t. 2.  
 A. *petrophila* var. *acuminata* Rab. *Bryoth. eur.* n. 52 (fid. Limpr.).  
 1. Ad rup. madid. — *Eur.* : Norveg., Scot., Alp., Pyren., S.  
 d'Estrella. *Am. sept.* : Groenland, Nov. Scot.  
 « var. *obtusifolia* Berggr. in *K. Sv. Vet.-Akad. Handl.* 1875, n. 8,  
 p. 40.  
 — *Am. sept.* : Groenland.
- crassinervia* } var. *Heinemanni* C. M. — A. *crassinervia*.  
 } var. *Huntii* Braithw. — A. *Huntii*.
- cuspidata* C. M. in *Bot. Centralbl.* 1883, n. 2.  
 2. — *As.* : Penins. Tschutschica.
- densifolia* Mitt. — *Acroschisma*.
- depressinervis* Card. in *Rev. bryol.* 1900, p. 43.  
 — *Am. merid.* : Fret. Gerlache.
- erubescens* C. M. in *Hedwig.* 1898, p. 79.  
 2? fr. ign. Rupic. — *Pac.* : Tasman. (m. Wellington).
- « var. *nigrita* C. M. in *op. et l. cc.*  
 A. *subulifolia* ej. *prius in sched.*  
 Ut et ubi typ.
- eximia* C. M. in *Hedwig.* 1898, p. 84.  
 2. — *Pac.* Tasman. (m. Wellington).
- falcata* *Bryol. eur. fasc.* 52-54 (Vol. VI), Mon. p. 24, t. 634 (1855); W.  
 P. Sch. *op. cit.* p. 821.  
 A. *Rothii* var. *falcata* (W. P. Sch.) Lindb. in *Braithw. Brit. Mossfl.* I,  
 p. 14 (1880).  
 A. *Rothii* var. *papillosa* C. M. *Syn.* I, p. 9 (1849).  
*Lichenastrum alpinum, nigricans, foliis capillaribus reflexis* Dill. *Hist. M.*  
 p. 507, t. 73, f. 40 (1741).  
 1. Ad rup. madid. — *Eur.* : Scot.? Cambrovall.? Hercyn.? Enga-  
 din., Rhaet., Harz, Thuring., Normann., Pyren.

- « *var. heteromalla* Lindb. *M. scand.* p. 31 (1879).  
 Ut typ. — *Eur.* : Scandin.
- falcata* Rab. — A. Huntii.
- Fauriei* Besch. in *Ann. sc. nat.* 1893, p. 392.  
 2. — *As.* : Japon.
- filamentosa* C. M. in *Hedwig*, 1898, p. 80.  
 Ster. — *Pac.* : Alp. Nov. Seeland. austr.
- filiformis* C. M. in *Bot. Centralbl.* 1883, n. 2-4.  
 Ster. — *As.* : Penins. Tschutschica.
- firma* C. M. in *Fl.* 1888, p. 406.  
 Ster. In graminos. — *Afr.* : Kilima N'jaro.
- flabellata* C. M. in *Engler's Bot. Jahrb.* 1883, p. 176 et in *Forschungsreise.... Bot.* p. 9 (1889).  
 2. Ad rup. sicc. — *Afr.* : Ins. Kerguelen.
- fragilis* C. M. in *Linn.* 1878-79, p. 223.  
 Fl. et fr. ign. — *Am. merid.* : Alp. Argent. subtropic.
- frigida* Hüb. *Hepat. germ.* p. 305, n. 4. (1874).  
 A. *grimsulana* Bruch mss.; de Not. *Epil.* p. 748 (1869).  
 A. *nivalis* β *frigida* Rab. *Deutsch. Kryptfl. II*, p. 3, t. 72 (1848).  
 A. *Rothii* β *frigida* Lindb. in *Braithw. Brit. Mossfl. I*, p. 13 (1880).  
 A. *Rothii* γ *grimsulana* (Br.) H. f. et W. (1844); C. M. *Syn. I*, p. 9 (1849).  
 A. *rupestris* β *grimsulana* *Bryol. eur. fasc.* 52-51 (*Vol. VI*), *Mon.* p. 22, t. 632 (1855); W. P. Sch. *op. cit.* p. 819.  
 1. Ad rup. madid. — *Eur.* : Succ., Scot., Tatra, Stir., Carinth., Rhaet., Pyren.
- « *var. cuspidata* Limpr. *Laubm. Deutschl.... I*, p. 151 (1885).  
 Ut typ. — *Eur.* : Stiria.
- « *var. sudetica* Limpr. in *LXI Jahresb. d. Schles. Gesells.* p. 222 (1883).  
 — *Eur.* : Riesengeb.
- grimsulama* Bruch. — A. *frigida*.
- Hartmani* Thed. in *Nya Bot. Not.* 1849, p. 78, t. 1-17 et 23-26; *Bryol. eur. fasc.* 52-54 (*Vol. VI*), *Mon.* p. 19, t. 269 (1855); C. M. *Syn. II*, p. 516 (1849); W. P. Sch. *op. cit.* p. 815.  
 1. Ad saxa irror. juxta niv. — *Eur.* : Lappon., Alp. Scandin.
- var. gracilis* Zett. }  
*var. Thedenii* Lindb. } — A. *Thedenii*.
- Heinmanni* Hpe et C. M. — A. *crassifolia*.
- homomalla* C. M. in *Hedwig*, 1898, p. 80.  
 Ster. Rupic. — *Pac.* : Nov. Seeland. austr.
- Huntii* Limpr. *Laubm. Deutschl.... I*, p. 145 (1885).  
 A. *commutata* Limpr. in *LXI Jahresb. d. Schles. Gesells.* 1883, p. 221 (*non* C. M.!).  
 A. *crassinervia* *var. Huntii* Braithw. *Brit. Mossfl. I*, p. 265 (1887).  
 A. *falcata* Rab. *Bryoth. eur. n.* 1301 a et b (*fid. Limpr.*).  
 1. Rupic. — *Eur.* : Norveg. occ., Britann., Harz, Bavar., Bade (Feldberg., Lusitan. *Am. sept.* : Ins. Vancouver.
- indica* Mitt. *M. Ind. or.* p. 6 (1859).  
 — *As.* : Sikkim.



- julicaulis* C. M. in *Hedwig*. 1898, p. 79.  
2. ? fr. ign. Rupic. — *Pac.* : Tasman. (m. Wellington).
- Karsteniama* C. M. in *Linn.* 1874, p. 572.  
A. petrophila Mitt. *M. austr. am.* p. 628 ?  
2. — *Am. merid.* : M. Pichincha.
- Kilima N'Djarica* Par. *Ind.* p. 30 (1893).  
A. striata C. M. in *Fl.* 1888, p. 406.  
In graminos. — *Afr.* : Kilima N'Djaro.
- Krauseana* C. M. in *Bot. Centralbl.* 1883, n. 2-4.  
2. — *As.* : Penins. Tschutschica.
- laxifolia* Hook. f. et Wils. in *Lond. Journ. of bot.* 1844, p. 536 et *Fl. antarct. II*, p. 397, t. 151, f. 4 (1847); C. M. *Syn. I*, p. 8 (1849) et *II*, p. 517 (1851); Mitt. *M. austr. am.* p. 629 (1869).  
1. — *Am. merid.* : Fuegia, ins. Eremitae.  
« var. minor C. M. op. et l. cc.  
Cum typ.
- Lorentziana* C. M. in *Linn.* 1878-79, p. 221.  
2. — *Am. merid.* : Alp. Argentin. subtropic.
- Macouni* Kindb. in *Bull. Torr. bot. Club*, April 1890, et in *Mac. Cat.* p. 10 (1892).  
2. Rupic. — *Am. sept.* : Columb. brit., Oregon.
- marginata* Hook. f. et W. op. et l. cc. p. 535, p. 396, t. 151, f. 1; C. M. *Syn. I*, p. 7 et 11, p. 516; Mitt. op. et l. cc.  
2. — *Afr.* : Ins. Kerguelen, Possessionis. *Am. merid.* : Fuegia, ins. Eremitae.
- microphylla* C. M. in *Bull. hb. Boissier*, 1898, p. 2 et in *Ule Coll. M. brasil.* n. 1730.  
Ster. Rupic. — *Am. merid.* : Brasil. or. (S. do Itatiaia).
- micro-vaginata* C. M. in *Hedwig*. 1898, p. 80.  
Ster. Rupic. — *Pac.* : Nov. Seeland. austr.
- montana* Mitt. in *Journ. of the Linn. Soc.* 1859, p. 64 et in *Fl. Tasman. II*, p. 161, t. 7, f. 1 (1860).  
Ster. Rupic. — *Pac.* : Tasman.
- Mülleri* Sonder. — A. asperula ?
- mutabilis* Hook. f. et Wils. in *Lond. Journ. of bot.* 1841, p. 536, (α *microphylla*) et *Fl. antarct. I*, p. 119, t. 57 (1844); C. M. *Syn. I*, p. 8 et 11, p. 516; *Handb. of the N. Zeal. Fl.* p. 400 (1867).  
— *Am. merid.* : Ins. Maclov. *Pac.* : Nov. Seeland., ins. Auckland, Campbell.
- « var. subsecunda } C. M. op. et l. cc.  
« var. uncinata } — *Am. merid.* : Ins. Maclov. (ambo).
- nana* C. M. in *Engler's Bot. Jahrb.* 1883, p. 176 et in *Forschungsreise...* *Bot.* p. 9 (1889).  
2. — *Afr.* : Ins. Kerguelen.
- Naumannii* C. M. in op. et l. cc.  
1. Ad rup. basalt. — *Afr.* : Ins. Kerguelen.

- nitida* Hook. f. et Wils. op. cc. p. 535, p. 118, t. 57, f. 3; C. M. Syn. I, p. 6 et II, p. 515; *Hundb. of the N. Zeal. Fl.* p. 490 (1867).  
Rup. Pac. : Nov. Seeland., Tasman., ins. Auckland.
- nivalis* Hook. in *Trans. of the Linn. Soc.* 1810, p. 395, t. 31, f. 4; *Brid. Bryol. univ.* II, p. 732 (1827); C. M. Syn. I, p. 9 (1849); *Bryol. eur. fasc.* 52-54 (Vol. VI), *Mon.* p. 25, t. 636 (1855); W. P. Sch. op. cit. p. 822.
- A. *nivalis* ♂ *Zetterstedti* Hartm. *Skand. Fl.* VII ed. p. 400 (1858).
2. Ad rup. aqua niv. irror. — *Eur.* : Scot., alp. Scand., Tatra, Salisb., Tirol., Helvet., alp. Sabaud., Pyren. *Am. sept.* : Groenland.
- var. *frigida* Rab. — A. *frigida*.
- « var. *fuscescens* Hook. op. cit. f. 4c; *Lindb. M. scand.* p. 30, n. 406 (1879).  
Ut typ. — *Eur.* : Lappon., Norveg.
- obovata* Thed. in *Nya Bot. Not.* p. 78, t. I, fig. 27-36 (1849); C. M. Syn. II, p. 515 (1851); *Bryol. eur. fasc.* 52-54 (Vol. VI), *Mon.* p. 17, t. 627 (1855); W. P. Sch. op. cit. p. 81.
- A. *alpina* Aongstr. in *Fr. Summ. veg. scand.* I, p. 97 (1846-49).  
A. *alpina* ♂ caule elongato, foliis majoribus fere obtusioribus *Wahlenb. Fl. Suec. II ed. P. II*, p. 809 (1833).  
*Jungermannia alpina*, surculis teretibus, foliis ovatis patentibus, calycibus imbricatis L. *Sp. pl.* p. 1601 p.p (1853).
1. Rupic. ad rivul. niv. — *Eur.* : Spitzberg, alp. Scand. *Am. sept.* : Groenland., Canada.
- « var. *acuminata* *Lindb. Spitzberg Mossor.* (1858).  
Ut typ. — *Eur.* : Spitzberg.
- obtusifolia* Th. Jens. in *Vid. Medd. Natur. For.* 1858, p. 55.  
Ad rup. silic. — *Eur.* : Norveg. *Am. sept.* : Groenland.
- obtusissima* C. M. in *Hedwig.* 1898, p. 83.  
— *Pac.* : Nov. Seeland. merid.
- pachyphylla* (C. M.) *Broth. Natur. Pflanzenfam.* p. 268.  
*Grimmia pachyphylla* C. M. in *Fl.* 1885, p. 422.
2. Rupic. — *Am. merid.* : Fuegia.
- papillosa* *Lindb. Spitzbergs-Mossor.* p. 557 (1866) et in *Hedwig.* 1868, p. 109.
1. Rupic. — *Eur.* : Spitzberg, Lappon. pitens. *As.* : Sibir. *Am. sept.* : Fret. Behring et Scoresby.
- « var. *brevifolia* *Berggr. M. Spetsb.* n. 155c.  
Ut typ. — *Eur.* : Ins. Ursorum.
- « var. *gracilis* *Lindb. in op. et II. cc.* : *Hedwig.* p. 110.  
Ut typ. — *Eur.* : Spitzberg.
- « var. *latifolia* *Berggr. Coll. cit.* n. 155b et in *K. Sv. Vet.-Akad. Foerh. XIII*, n. 7, p. 83.  
Ut typ. — *Eur.* : Ins. Parry, Spitzberg.
- « *subsp. sublaevis* *Kindb. Note on Canad. bryol.* 1893, (ut var.) et *Eur. a. N. Amer. Bryin.* p. 393 (1897).  
— *Am. sept.* : Canada, ins. Vancouver.
- « *subsp. subsarsifolia* *Kindb. Eur. and N. Am. Bryin.* p. 393 (1897).  
Ad rup. marit. — *Eur.* : Norveg. littor.

« *subspec. tenella* *ef. op. et l. cc.*

— *Eur.* : Pyren. (Venasque).

parallela *C. M. in Engler's Bot. Jahrb.* 1883, p. 76, *et in Forschungsreise.... Bot. p.* 8 (1889).

2. *Ad rup. sicc.* — *Afr.* : Ins. Kerguelen.

*parvifolia C. M. in Fl.* 1887, p. 219.

2. — *Am. sept.* : Alaska.

*patens C. M. in Bot. Centralbl.* 1883, n. 2-4.

2. — *As.* : Penins. Tschutschica.

*perichaetialis* } *H. f.* — *A. subulata* *var. perichaetialis.*  
                           } *Zett.* — *A. Blyttii.*

*petrophila Ehrh. in Hann. Mag.* 1784, p. 140 *et in Beitr. I, p.* 192;  
*Bryol. eur. fasc.* 52-54 (*Vol. VI*), *Mon. p.* 13, t. 623 (1855); *W. P. Sch. op. cit. p.* 812.

*A. alpina Hedw. Sp. M. t.* 7, f. 2 (1801); *Turn. Musc. Hib. p.* 13 (1801);  
*W. et M. Bot. Taschenb. t.* 2, fig. 3-4 (1807); *Brid. Bryol. univ. II,*  
*p.* 728 (1827).

*A. rupestris Hedw. op. et l. cc.*; *C. M. Syn. I, p.* 6 (1849); *H. f. et W. Fl. of the N. Zeal. II, p.* 57 (1855) *et Handb. of the N. Zeal. Fl. p.* 400 (1867).  
*Jungermannia alpina (ut supra ad A. obovatam L.)*

1. *Rupic.* — *Eur.* *E. reg. campestr. montos. usq. in alp. As.* : Japon. ?  
*Caucas. Am. sept.* : Groenland, Labrador, Terr. nov., ins. Miquelon,  
 Nov. Angl., Nov. Scot., M. Alb., Ontario, Alaska, ins. Vancouver,  
 Washington, M. Rupestr. et Selkirk, Idaho, Columb. brit.  
*Am. merid.* : M. Pichincha (an *A. Karsteniana* ?), *Fuegia. Pac.* :  
 Nov. Seeland, Tasman.

« *var. acuminata Bryol. eur. et W. P. Sch. t.* 624, *et p.* 813.

*Ut typ.* — *Eur.* : Suecia (prov. Upland).

*var. alpestris* Thed. — *A. alpestris.*

« *var. alpicola Bryol. eur. et W. P. Sch. II. cc.*

*Ut typ.* — *Eur.* : Hercyn. *Am. sept.* : Labrador.

« *var. flaccida Bryol. eur. et W. P. Sch. II. cc.*

*Ut typ.* — *Eur.* : Suecia (prov. Upland).

« *var. gracilis Bryol. eur. et W. P. Sch. II. cc.*

*Ut typ.* — *Eur.* : Thuring.

*var. homomalla* *W. P. Sch.* — *A. petrophila var. rupestris.*

« *var. pygmaea Bryol. eur. et W. P. Sch. II. cc.*

*Ut typ.* — *Eur.* : Alp. editior.

« *var. robusta Bryol. eur. t.* 625; *W. P. Sch. l. c.*

*Ut typ.* — *Eur.* : Voges.

« *var. rupestris (Hedw.) Wallr. Fl. germ. crypt. I, p.* 192 (1831).

*A. petrophila var. homomalla W. P. Sch. Syn. I ed. p.* 661 *et II ed. l. c.*  
*A. rupestris Hedw. Sp. M. p.* 47, t. 3, f. 2 (1801).

*Ut typ.* — *Eur.* : Septentr.

« *var. silvicola Bryol. eur. t.* 624; *W. P. Sch. Syn. II ed. l. c.*

*Ut typ.* — *Eur.* : In silv. subalp. *Am. sept.* : Alaska.

*var. sparsifolia* *Lindb.* — *A. sparsifolia.*

« *var. squarrosula Bryol. eur. et W. P. Sch. II. cc.*

*Ut typ.* — *Eur.* : Suecia (prov. Upland).

*petrophila* { Ehrh. — *A. alpina* p.p.  
 { Mitt. } — *A. commutata*.  
 { — *A. Karsteniana* ?

*pseudo-alpina* C. M. in *Bot. Zeit.* 1859, p. 248.

*A. alpina* Mitt. *M. austr. am.* p. 620 (1869).

2. — *Am. merid.* : Cap. Horn, ins. Eremitae.

*pseudo-subulata* C. M. — *A. subulata* var. *rigida*.

*pulvinata* C. M. in *Trans. and Proceed. of the N. Zeal. Inst.* 1892, p. 293,  
 t. XL et in *Hedwig.* 1898, p. 81.

2. fr. ign. Rupic. — *Pac.* : Nov. Seeland.

*pycnotila* Card. in *Rev. bryol.* 1900, p. 42.

— *Am. merid.* : Fret. Gerlache.

*pygmaea* Card. in *op. cit.* p. 43.

— *Am. merid.* : Fret. Gerlache.

*regularis* C. M. *Bryol. austr. Georg.* p. 10 (1890).

1. — *Am. merid.* : Austro-Georg.

*rigida* Wils. in *Kew. Journ. bot.* IX, p. 280 (1857); Mitt. *M. Ind. or.*  
 p. 6 (1859).

— *As.* : Sikkim.

*Rothii* Brid., C. M., W. et M. — *A. rupestris*.

var. *falcata* Limpr. — *A. falcata*.

var. *frigida* Lindb. — *A. frigida*.

var. *grimsulana* H. f. et W. — *A. frigida*.

var. *hamata* Lindb. — *A. rupestris* var. *hamata*.

var. *papillosa* C. M. — *A. falcata*.

*rupestris* Roth *Neue Beitr.* I, p. 234 (1802); *Bryol. eur. fasc.* 52-54  
 (vol. VI), *Mon.* p. 21, t. 631 (1855); W. P. Sch. *op. cit.* p. 819.

*A. Rothii* Web. et Mohr *Bot. Taschb.* p. 386, t. II, f. 7-9 (1807); *Brid.*  
*Bryol. univ.* II, p. 740 (1827); C. M. *Syn.* I, p. 9 (1849).

*Jungermannia rupestris*, foliis subulatis secundis L. *Fl. Suec.* p. 1045 (1716).  
*Lichenastrum alpinum*, nigricans, foliis capillaribus reflexis Dill. *Hist. M.*  
 p. 507, t. 73, f. 40<sup>a</sup> p.p. (1741).

1. Ad rup. praecip. silic. et in molib. errat. — *Eur.* : E reg. cam-  
 pestr. montos. usq. in subalp. *Am. sept.* : Groenland., sin.  
 Hudson, New-York, Georgia, Carolin.

« var. *capensis* Rehm. *M. Afr. austr. n.* 430.

Ut typ. — *Afr.* : Pr. B. Sp.

var. *grimsulana* *Bryol. eur.* — *A. frigida*.

« var. *hamata* (Lindb.) *Par. Ind.* p. 33 (1893).

*A. Rothii* var. *hamata* Lindb. *M. scand.* p. 31, nom. (1879), et in *Braithw.*  
*Brit. Mossfl.* I, p. 14 (1890).

Ut typ. — *Eur.* : Scandin.

« var. *septentrionalis* *Bryol. eur. et W. P. Sch. II. cc.*

Ad. scopul. aprie. — *Eur.* : Norveg.

*rupestris* { C. M. p.p. } — *A. petrophila*.  
 { — *A. striata*.  
 { Hedw. } — *A. petrophila* var. *rupestris*.  
 { — *A. striata*.  
 { H. f. et W. } — *A. appendiculata*.  
 { — *A. petrophila*.

semi-squarrosa *C. M. in Linn.* 1878-79, p. 222.

Fr. ign. — *Am. merid.* : Alp. Argentin. subtropic.

sparsifolia *Zett. Mon. Andr.* p. 32, n. 4 (1855); *W. P. Sch. Coroll.* p. 94 in *adn.* (1856) *et op. cit.*

*A. petrophila* var. *sparsifolia* *Lindb. Spitsb. Mossor.* p. 559 (1867) *et in Braithw. Brit. Mossfl.* p. 9 (1880).

1. Rupic. juxta niv. — *Eur.* : Alp. Norveg., Tirol. Ticin.

« var. *sublaevis* *Kindb. Note on Canad. Bryol.* 1893.

— *Am. sept.* : Ins. Vancouver.

spurio-alpina *C. M. in Bull. hb. Boissier*, 1898, p. 98 *et in Ule Coll. M. brasil.* n. 1100, 1729.

*A. striata* *Broth. in Ule Bryoth. brasil.* n. 97.

2. Rupic. — *Am. merid.* : Brasil. or. (S. do Itatiaia).

« var. *rubricalyx* *C. M. in op. et l. cc. et in Coll. cit.* n. 127.

Ut *et ubi* typ.

squamata *C. M. in Engler's Bot. Jahrb.* 1883, p. 77 *et in Forschungsreise ... Bot.* p. 10 (1889).

2. Ad rup. basalt. sicc. — *Afr.* : Ins. Kerguelen.

squarrosa *Mitt. M. austr. am.* p. 629 (1869).

— *Afr.* : Ins. Kerguelen. *Am. merid.* : And.

Quitens.

squarrosa-filiformis *C. M. in Bull. hb. Boissier*, 1898, p. 19.

Ster. Rupic. — *Am. merid.* : Brasil. or. (S. do Itatiaia).

striata *Mitt. M. austr. am.* p. 628 (1869).

*A. rupestris* *C. M. Syn. I*, p. 6 (*quoad. pl. brasil.*).

Rupic. — *Am. merid.* : And. Quitens., Brasil. austr. or.

striata } *Broth.* — *A. spurio-alpina.*

} *C. M.* — *A. Kilima N'Djarica.*

subappendiculata *C. M. in Engler's Bot. Jahrb.* 1883, p. 76 *et in Forschungsreise.... Bot.* p. 8 (1889).

*A. acuminata* *Mitt. ?*

Ster. Ad rup. sicc. — *Afr.* : Ins. Kerguelen.

subnervis *Hook. f. et Wils. in Lond. Journ. of bot.* 1847, p. 289; *C. M. Syn. I*, p. 8 (1849); *Mitt. M. austr. am.* p. 630 (1869).

— *Am. merid.* : And. Nov. Granat. et Quitens.

subfluitans *C. M. in Hedwig.* 1898, p. 81.

Ster. Ad saxa irror. — *Pac.* : Nov. Seeland. austr.

subulata *Harv. in Hook. Ic. pl. rar. III*, t. 201 (1841); *C. M. Syn. I*, p. 10.

1. Rupic. — *Afr.* : Pr. B. Sp.

« var. *perichaetialis* *H. f. et W. in Lond. Journ. of bot.* 1844, p. 538.

Ut typ. — *Am. merid.* : Ins. Eremitae, Maclov.

« var. *rigida* *Harv. et C. M. op. et ll. cc.*

*A. pseudo-subulata* *C. M. in Bot. Zeit.* 1864, p. 373.

*A. subulata* *H. f. et W. Fl. antarct. I*, p. 119, t. 57, f. 1 (1844) *et Handb. of the N. Zeal. Fl.* p. 40 (1867); *Mitt. M. austr. am.* p. 630 (1869).

Ut typ. — *Am. merid.* : Fuegia, ins. Eremitae. *Pac.* : Austral., Nov. Seeland. sept., Tasman., ins. Auckland, Campbell.

*subulata* H. f. et W. — *A. subulata* var. *rigida*.

*subulatissima* C. M. in *Hedwig*. 1898, p. 82.  
2. — *Pac.* : Tasman.

*subulifolia* C. M. — *A. erubescens* var. *nigrita*.

*tenera* C. M. in *op. cit.* p. 84.  
2. fr. ign. — *Pac.* : Austral. or. (Victoria).

*Thedenii* *Bryol. eur. fasc.* 52-54 (*Vol. VI*), *Mon.* p. 20, t. 630 (1855) ; W. P. Sch. *Syn. II* ed p. 816 (1876).

*A. Hartmani*  $\beta$  *gracilis* Zett. *Mon.* ?

*A. Hartmani* var. *Thedenii* Lindb. *M. scand.* p. 31 (1879).

1. Ad et in rivul. nival. — *Eur.* : Lappon., Norveg. (Dovre).

*turgescens* W. P. Sch. in C. M. *Syn. II*, p. 515 (1851).  
— *Am. sept.* : Summ. m. Orizaba, Nevado de Toluca.

*viridis* C. M. *Bryol. austr. Georg.* p. 10 (1890)  
2. — *Am. merid.* : Austr. Georg.

*vulcanica* Ltz. *Moosst.* p. 156 (1864).  
2. — *Am. merid.* : M. Ilinitza.

*Willii* C. M. *Bryol. austr. Georg.* p. 11 (1890).  
2. — *Am. merid.* : Austro-Georg.

*Wilsoni* C. M. — *Acroschisma*.

**Aneuron** Kindb. *Eur. and N. Am. Bryin* p. 165 (sect. FISSIDENTIS) (1897).

ANHYMENIUM Griff. *Not.* p. 471 (1849).

*polycarpus* ej. — *Rhegmatodon*.

*polysetus* ej. — *Rhegmatodon orthostegius* ?

ANICTANGIUM (*rid.* ANOECTANGIUM).

ANISODON *Bryol. eur. fasc.* 49-51 (*Vol. V*), t. 452 (1852).

<i>acutirostris</i> Bryol. eur.	} <i>Clasmatodon parvulus</i> .
<i>Bertrami</i> W. P. Sch.	
<i>perpusillus</i> Bryol. eur.	

ANISOSTICHIMUM Mitt. in *Journ. of the Linn. Soc. VII*, p. 119 (1863).

*Bakeri* ej. — *Epipterygium Wrightii*.

*pictum* ej. — *Webera*.

*Tozeri* ej. — *Epipterygium*.

ANISOTHECIUM Mitt. *M. austr. am.* p. 39 (1869).

Spec. omn. non indicatae sunt DICRANELLÆ ejusd. nom.

*convolutum* Mitt. — *Dichodontium*.

*crispum* Lindb. — *Dicranella Schreberi*.

*ferrugineum* Mitt. — *Ditrichum brachycarpum*.

*humile* Lindb. — *Dicranella varia* var. *tenella*.

*rubrum* Lindb. — *Dicranella varia*.

var. *obtusiuscula* Lindb. — *Dicranella varia* var. *obtusiuscula*.

ANODON Rat. *Deutsch. Kryptfl. II*, 3, p. 154 (1848).

*pulvinatus* ej. — *Grimmia sphaerica*.

*ventricosus* ej. — *Grimmia anodon*.

**ANODONTIUM** *Brid. Sp. M. I, p. 41* (1806).

*prorepens* ej. — *Drummondia clavellata*.

**ANODUS** *Bryol. eur. fasc. 33-36 (Vol. I) (1846)*. — 1 spec.

*Donianus (Smith) Bryol. eur. l. c. Mon. p. 3, t. 109; W. P. Sch. Syn. II ed. p. 124* (1873).

*Gymnostomum Donianum Sm. Engl. Bot. XXII, n. 1582* (1806).

*G. pusillum Drum. M. bor. am. I ed. n. 22* (1828).

*G. rigidum Wallr. in Linn. 1840, p. 681.*

*Seligeria Doniana C. M. Syn. I, p. 420* (1849 *et II, p. 603* (1851).

*S. Donii Lindb. in Oefv. af Vet.-Akad. Föerh. 1854, p. 187.*

1. *Ad rup. arenac. et calcar.* — *Eur.* : Britann., Norveg., Siles., Saxon., Gallic., Tatra, Austr. inf., Salisb., Stir., Tirol., Bavar., Helvet., Jurass., Bade., Würtemb., Westphal., Thuring., Harz, etc.  
*As.* : *Jenissei super. Am. sept.* : New-York, Ontario, M. Rupestr.

**ANOECTANGIUM** (Hedw.) *Bryol. eur. fasc. 29-30 (Vol I) (1846)*. — 56 spec.

*abyssinicum Hpe in W. P. Sch. M. Abyss. n. 1117.*

*Rupic.* — *Afr.* : Abyss. (Janka Herr, 2300 met. alt.)

*acaule* Roehl. — *Pterygoneuron subsessile.*

*aestivum Mitt. M. austr. am. p. 175* (1869).

*A. enchloron ej. in M. Jameson.*

— *Am. merid.* : And. Quitens.

*aestivum* *Bryol. eur.* — *A. compactum.*

*afro-compactum C. M. in Dusen M. Camer. n. 342; Dus. in K. Sv. Vet.-Akad. Föerh. 1895, n. 3.*

2. *Rupic.* — *Afr.* : Summ. m. Cameroon.

*andinum* Mitt. — *Molendoa.*

*angustifolium Mitt. in Journ. of the Linn. Soc. 1865, p. 2.*

*A. Knyi Jur. in Bot. Zeit. p. 20.*

*A. madeirense W. P. Sch. in Mandon Pl. Madeir.*

*A. Paivanum ej. in Coll. cit. n. 2.*

*Rupic.* — *Afr.* : Teneriff, Madeir.

*apiculatum W. P. Sch. in Besch. Prodr. bryol. mexic. p. 16* (1871).

— *Am. sept.* : Mexic.

*aquaticum* Hedw. — *Cinclidotus.*

*Balfourei Mitt. in Trans. of the roy. Soc. of Edinb. 1887, p. 445.*

*A. Mariei ej. in op. cit. p. 332.*

— *Afr.* : Ins. Socotora.

*basalticum* Dusen. — *A. Dusen.*

*birmense C. M. in Nuov. Giorn. bot. ital. XXIII, p. 600.*

— *As.* : Barman.

*borbonense Besch. Fl. bryol. Réunion, etc. p. 12* (1880-81).

*Terr.* — *Afr.* : Ins. Borbon.

*Breutelianum* *Bryol. eur.* — *A. euchloron.*

*bulbosum* Hedw. — *Cyatophorum.*

*caespitium* Schwaegr. — *Stylostegium.*

callidum *Mitt. M. austr. am. p. 177 (1869).*

In sticilid. callid. — *Am. merid.* : And. Quitens.

canadense *Kindb. in Ottawa Nat. 1900, V. p. 86.*

*Zygodon caespitosus ej. Eur. and N. Am. Bryin. p. 316 (1897).*

Ster. — *Am. sept.* : Ins. Vancouver.

*ciliatum* { C. M. — *Hedwigia nivalis.*  
Hedw. — *Hedwigia.*

*cirrosum* Hedw. — *Macromitrium.*

clarum *Mitt. M. Ind. or. p. 31 (1859).*

A. compactum var. *Wils. in Kew Journ. bot. IX, p. 326 (1857).*

2. — *As.* : Sikkim, Ceylan.

compactum *Schwaegr. Suppl. I, P. I, p. 36, t. II (1811); Bryol. eur. fasc. 29-30 (Vol. I), Mon. p. , t. 37 (1846); W. P. Sch. Syn. II ed. p. 47 (1876).*

A. aestivum *Bryol. eur. l. c.*

? *Gymnostomum aestivum Hedw. Sp. M. p. 32, t. II (1801).*

G. compactum *Schwaegr. in Schleicher's Catal. (1815); Brid. Bryol. univ. I, p. 86 (1826).*

G. compactum seu aestivum *Schleich. Crypt. helv. II, n. 7.*

G. luteolum *Engl. Bot. n. 2201.*

G. tristichum *Wahlenb. Fl. lapp. p. 303 (1812).*

*Hedwigia aestiva Hook. Engl. Fl. V, I, p. 68.*

*Pleurozygodon aestivus Lindb. M. scand. p. 29 (1879).*

*Zygodon compactus C. M. Syn. I, p. 683 (1849) et II, p. 637 (1851).*

2. In fiss. rup. madid. praecip. micac. schist. — *Eur.* : Reg. alp. tota, rarius in reg. subalp. et montana. *As.* : Caucas. *Afr.* : Ins. Canar., Madeir., Cameroen. *Am. sept.* : Groenland. *Pac.* : Nov. Seeland ?

\* var. *brevifolium Jur. in sched.; Milde Bryol. siles., p. 109 (1869).*

Ut typ. — *Eur.* : Sudet.

compactum var. *Wils. — A. clarum.*

condensatum *W. P. Sch. in Besch. Prodr. bryol. mexic. p. 16 (1871).*

— *Am. sept.* : Mexic.

? *coronatum* H. f. et W. — *Erpodium.*

*crassinervium* *Mitt. — A. pusillum.*

crispulum *Wils. in Kew Journ. bot. IX, p. 325 (1857).*

A. *Thomsoni* *Mitt. M. Ind. or. p. 31 (1859).*

*Tortula* \* *T. paludosae* allied \* *Wils. in op. cit. p. 322.*

2. — *As.* : N. W. Himalaya.

*domingense* *Schwaegr. — Erpodium.*

*Dusen* *C. M. in Dusen M. Camer. n. 157.*

A. *basalticum Dusen* in *K. Svet. Vet.-Akad. Foerh. 1895, n. 5.*

2. *Rupic.* — *Afr.* : Cameroen.

*Duthiei* *Broth. — Molendoa.*

*Eggersii* (C. M.) *Par. — Hymenostylium.*

*Erpodium domingense* *Brid. — Erpodium.*

euchloron (*Schwaegr.*) *Mitt. M. austr. am. p. 176 (1869).*

A. *Breutelium* *Bryol. eur. l. c. (Mon. p. 6, nom.) et in Besch. Prodr. bryol. mexic. p. 16 (1871).*



- A. tenellum* Mitt. in *Jameson M. exs. et in Lond. Journ. of bot.*  
*Gymnostomum euchloron* *Schwaegr. Suppl. II, P. II, p. 83, t. 176* (1827).  
*G. guadalupense* *Spreng. Syst. veg. IV, I, p. 145* ( ).  
*G. lamprocarpum* *Mont. in Ann. sc. nat. 1838, p. 49 et Syll. p. 50* (1856).  
*G. maschalocarpum* *ej. hb.*  
*Zygodon pusillus* *C. M. Syn. I, p. 684* (1849) *et II, p. 637* (1851).  
*Z. tenellus* Mitt. in *Journ. of bot. 1851, p. 56*; *C. M. Syn. II, p. 638*.  
 2. *Terr. — Am. sept. : Mexic., Guatemala, Guadalup., Martinic.,*  
*S. Doming. As. : Java. Am. merid. : Ins. Trinitat., And. Quitens.*  
*euchloron* Mitt. p.p. — *A. aestivum*.  
*eucollum* Besch. — *A. rufo-viride* var. *eucollum*.  
*excelsum* (C. M.) Par. — *Molendoa*.  
*falcatum* P. B. — *Cinclidotus aquaticus*.  
*ferrugineum* Besch. in *Ann. sc. nat. 1893, p. 329*.  
 2. *Rupic. — As. : M. Japon.*  
*flaccidum* de Not. } (fid. C. M.) — *Grimmia anodon*?  
 } (fid. Husn.) — *Grimmia sphaerica*.  
*glauescens* W. P. Sch. in *Besch. Prodr. bryol. mexic. p. 16* (1871).  
 — *Am. sept. : Mexic.*  
*Haleakalae* (C. M.) Par. *Suppl. Ind. p. 12* (1900).  
*Zygodon Haleakalae* C. M. in *Fl. 1896, p. 450*.  
 — *Pac. : Ins. Hawai.*  
*Hobsoni* Mitt. in *Trans. of the roy. Soc. of Edinb. 1887, p. 445*.  
 — *Afr. : Aden.*  
*Hornschuchianum* } *Bryol. brit. — Trichostomum hibernicum.*  
 } *Funck. — Molendoa.*  
*Humbloti* Ren. et Card. in *Bull. Soc. roy. bot. Belg. 1893, II, p. 9*; *Ren.*  
*M. Masc. Mad. n. 2.*  
*Ster.* — *Afr. : Comor. magn.*  
*Humboldtianum* Breut. } — *Rhacocarpus.*  
*Humboldti* Brid.  
*hymenodontoides* (C. M.) *Jaeg. Ad. II, p. 632* (1877-78).  
*Zygodon hymenodontoides* C. M. in *Linn. 1869-70, p. 40*.  
*Ster. Cortic. — Am. : Ceylan.*  
*ikaoense* Besch. in *Oefv. af K. Vet.-Akad. Foerh. 1900, p. 40*.  
 1. — *As. : Japon.*  
*imberbe* } *Drumm. — Hedwigia ciliata* var. *viridis.*  
 } *Hook. et Tayl. — Hedwigidium.*  
*impressum* Hpe. in *Linn. 1874, p. 208*.  
 2. — *Afr. : Madagascar.*  
*indicum* Mont. — *Braunia*.  
*jamaicense* (C. M.) Par. *Suppl. Ind. p. 13* (1900).  
*Zygodon jamaicensis* C. M. in *Bull. hb. Boissier, 1897, p. 558*.  
 — *Am. sept. : Jamaic.*  
*kilima N'Djaricum* Par. *Ind. p. 39* (1894).  
*A. pusillum* Mitt. in *Journ. of the Linn. Soc. 1886, p. 305*.  
 — *Afr. : Kilima N'Djaro.*  
*Enyi* Jur. — *A. angustifolium*.

- laetum* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1899, II, p. 715.  
Ster. — As. : Sikkim.
- lapponicum* } Hedw. — Amphoridium.  
                  } Kze. — Amphoridium curvipes.
- laxifolium* (Kze) Par. — Gymnostomum.
- laxum* C. M. in *Nuov. Giorn. bot. ital.* 1898, p. 187.  
2. — As. : China centr. (prov. Schen Si).
- Lechlerianum* W. P. Sch. in *Lechler Pl. peruv. n.* 2720; *Mitt. M. austr. am.* p. 117 (1869).  
2. Rupic. — Am. merid. : Peruv.
- \* *var. laetior* Hpe. in *Fl.* 1865, p. 37.  
Ubi typ.
- Liebmanni* W. P. Sch. in *Besch. Prodr. bryol. mexic. p.* 15 (1871).  
— Am. sept. : Mexic. (m. Orizaba).
- madeirense* W. P. Sch. — A. angustifolium.
- mafatense* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1890, I, p. 167;  
*Ren. M. Masc. Mad. n.* 101.  
— Afr. : Ins. Borbon.
- mamabolense* Rehm. *M. Afr. austr. n.* 435.  
Ster. ? — Afr. : Transvaal.
- Mandonianum* W. P. Sch. in *Mandon Pl. Boliv. n.* 1606.  
*Zygodon linearis* C. M. in *Linn.* 1878-79, p. 370.  
*Zygodon Mandonianus* ej. in *Nuov. Giorn. bot. ital.* 1897, p. 57.  
2. Rupic. ? — Am. merid. : And. Boliv., alp. Argentin. subtropic.
- Mariei* Besch. *Fl. bryol. Réunion, etc. p.* 13 (1880-81).  
2. fr. ign. Terr. — Afr. : Ins. Nossi Be, S. Maria de Madag.
- Mariei* Mitt. — A. Balfourei.
- Mougeotii* Lindb. — Amphoridium.
- obtusiuscuspis* Besch. in *Ann. sc. nat.* 1892, I, p. 49.  
2. Silv. — As. : Yunnan.
- Paivanum* W. P. Sch. — A. angustifolium.
- parasiticum* Brid. — Calymperes.
- paucidentatum* — C. M. Didymodon.
- Peckii* Sull. in *Aust. M. Appal. n.* 64 (1870) et *l. c. M. Suppl. p.* 38, t. 25 (1874); *Lesq. et Jam. Man. p.* 55 (1884).  
*Amphoridium Peckii* Sull. in *R. Rep. of Univ. of N. York*, 1869, p. 57.  
Ster. Rupic. — Am. sept. : New-York.
- peruvianum* Sull. in *Amer. Expl. Exped.* 1859, t. I; *Mitt. M. austr. am. p.* 177 (1869).  
— Am. merid. : M. Chimborazo.
- ? *planifolium* Hedw. *Sp. M. p.* 44, t. 6 (1801); *Brid. Bryol. univ. II*, p. 167 (1827); *C. M. Syn. II*, p. 173 (1851) (1).  
— Pac. : Ins. austral.
- porto-ricense* Spreng. — *Philonotis tenella* var. *humilis*.

---

(1) « Verosimiliter Neckeracea ». C. M. l. c.

- pulvinatum* Mitt. in *Trans. of the Linn. Soc.* 1891, p. 160.  
 Ster. — As. : Japon.  
*pulvinatum* Roehl. — *Grimmia sphaerica*.  
*pusillum* Wils. in *Kew Journ. bot.* IX, p. 325 (1857).  
 A. *crassinervium* Mitt. *M. Ind. or.* p. 31 (1859).  
 Ster. — As. : Tibet. occid.  
*pusillum* Mitt. — A. kilima N'Djaricum.  
*rhaphidostegium* C. M. in *Besch. Fl. bryol. Réunion, etc.* p. 13 (1880-81).  
 — Afr. : Ins. Borbon., Comor. magn.  
*repens* Hook. — *Gigaspermum*.  
*Roylei* Mitt. — *Molendoa*.  
*rufo-viride* Besch. *op. et l. cc.*  
 — Afr. : Ins. Comor. magn.  
 « var. *eucollum* *ej. op. et l. cc.*  
 Ubi typ.  
*scabrum* Broth. in *Engler's Bot. Jahrb.* 1894, p. 176.  
 2. fr. ign. — Afr. : Usambara.  
*schensianum* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 260.  
 — As. : China centr. (prov. Schen Si).  
*Schimperii* Mitt. in *Trans. of the roy. Soc. of Edimb.* 1887, p. 445 et in *W. Sch. Pl. Abyss.* n. 1147.  
 — Afr. : Abyss.  
*Schliephackei* Par. — *Pleuroweisia*.  
*Schmidii* (C. M.) Par. — *Oreoweisia*.  
*sciuroides* Bals. et de Not. — *Braunia*.  
*secundum* } Brid. — *Braunia*.  
 } Mont. — *Braunia indica*.  
*Sendtnerianum* Bryol. eur. — *Molendoa*.  
*setosum* Hedw. — *Cyrtopus*.  
*sordidum* Mitt. *M. austr. am.* p. 176 (1869) (1).  
 — Am. merid. : M. Chimborazo.  
*spathulatum* Mitt. in *Journ. of the Linn. Soc.* 1863, p. 150.  
 — Afr. : M. Cameroon.  
*sphaericum* Spreng. — *Physcomitrium*.  
*Stevensii* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1895, I, p. 57.  
 2. — As. : Sikkim.  
*Stracheyanum* Mitt. — A. *tortifolium*.  
*striatum* } Brid. — *Amphoridium lapponicum*.  
 } Wils. — *Hedwigia ciliata* var. *striata*.  
*Stuhlmanni* Broth. — *Barbula*.  
*subsessile* Spreng. — *Pterygoneuron*.  
*tapes* Besch. in *Bull. Soc. bot. Fr.* 1898, p. 53.  
 — Pac. : Taïti.  
*tasmanicum* Broth. in *Oefv. af Finska Vet.-Akad. Foerh.* 1899, p. 137.  
 2. fr. ign. — Pac. : Tasman.  
*tenellum* Mitt. — A. *euchloron*.  
*tenerrimum* C. M. — *Gymnostomum laxifolium*.

---

(1) An *Molendoa* ? *fid.* Broth.

*tenuinerve* Par. — Molendoa.

*Thomsoni* Mitt. — *A. crispulum*.

*torquatum* Broth. in *Engler's Bot. Jahrb.* 1897, p. 232.

*Zygodon torquatus* ej. in *Engler's Ost-Afrika*, V, *Pflanzenwelt*, C. p. 70 (nom.).

2. fr. ign. Rupic. — *Afr.* : Usambara.

*torquatum* Brid. — *Helicophyllum*.

*torquescens* Mitt. in *Trans. of the Linn. Soc.* 1891, p. 160.

Ster. — *As.* : Japon.

*tortifolium* Wils. in *Kew Journ. bot.* IX, p. 326 (1857).

*A. Stracheyanum* Mitt. *M. Ind. or.* p. 31 (1859).

2. — *As.* : W. Himalaya.

*trichodes* Schwaegr. — *Brachyodus*.

*viridatum* C. M., Par. — *Ceratodon*.

*Walkeri* Broth. in *Records of the bot. Survey of India*, 1899, p. 311.

2. fr. ign. Ad rup. gneiss. sicc. — *As.* : M. Ghates (Coorg.)

*weissioides* C. M. in *Fl.* 1897, p. 333.

— *Am. merid.* : And. Venezuel.

*Wilmsianum* (C. M.) Par. *Suppl. Ind.* p. 13 (1900).

*Zygodon Wilmsianus* C. M. in *Hedwig.* 1899, p. 113.

2. In fiss. rup. — *Afr.* : Transvaal.

# **ANOMOBRYUM** W. P. Sch. Syn. I, ed. p. 382 (1860). — 36 spec.

*abyssinicum* Jaeg. — *A. filiforme* var. *abyssinicum*.

? *afro-julaceum* (C. M.) Par.

*Bryum afro-julaceum* C. M. in *Dusen M. Camer.* n. 295; *Par. Ind.* p. 159 (1894).

— *Afr.* : Cameroon.

*astorense* Broth. *Natür. Pflanzenfam.* p. 561.

*Bryum astorense* ej. in *Act. Soc. sc. Fenn.* 1898, II, p. 26; *Par. Suppl. Ind.* p. 57 (1900).

2. — *As.* : Kashmir.

*auratum* (Mitt.) Jaeg. *Ad. I*, p. 604 (1873-74).

*Brachymerium filiforme* Wils. in *Kew Journ. bot.* IX, p. 363 p.p. (1857).

*Bryum auratum* Mitt. *M. Ind. or.* p. 67 (1859); *Par. Ind.* p. 161 (1894).

2. — *As.* : Nepal., Himalaya.

*brachymeniopsis* (C. M.) Broth. *op. et l. cc.*

*Bryum brachymeniopsis* C. M. in *Linn.* 1878-79, p. 301; *Par. op. cit.* p. 167.

2. — *Am. merid.* : Argentin. Tucuman.

*bullatum* (C. M.) Broth. *op. et l. cc.*

*Bryum bullatum* C. M. in *Fl.* 1887, p. 221; *Par. op. cit.* p. 168.

2. — *Am. sept.* : Alaska.

*campestre* Jaeg. — *A. filiforme*.

ceramioecarpum (C. M.) Broth. op. et l. cc.

Bryum ceramioecarpum C. M. in Linn. 1878-79, p. 476; Par. op. cit. p. 172.

2. — Am. merid. : Venezuela.

chlorosum (C. M.) Broth. op. et l. cc.

Bryum chlorosum C. M. in Bull. hb. Boissier, 1897, p. 549; Par. Suppl. Ind. p. 60 (1900).

2. — Am. sept. : Jamaïc.

clavicaule (C. M.) Broth. op. et l. cc.

Bryum clavicaule C. M. in Fl. 1875, n. 34-35; Par. Ind. p. 173 (1894).

2. fr. ign. — Am. merid. : And. Nov. Granat.

compressulum (C. M.) Broth. op. et l. cc.

Bryum compressulum C. M. in Fl. 1888, p. 409; Par. op. cit. p. 174.

Ster. — Afr. : Kilima N'Djaro.

concinatum (Spruce) Lindb. Man. M. I, p. 44 (1870).

A. julaceum var. concinatum Zett. M. Pyren. p. 34 (1866).

Bryum concinatum Spr. M. Pyren. n. 121 et in Ann. a. Mag. of nat. hist. 1849, p. 155, n. 141; C. M. Syn. II, p. 575 (1851); W. P. Sch. Syn. II ed. p. 468 (1876); Par. op. cit. l. cc.

Br. filiforme var. concinatum Boul. M. Fr. I, p. 294 (1884).

Br. julaceum var. concinatum Wils. Bryol. brit. p. 246 (1855).

Br. nimbigerum Tayl. mss.; Hartm. Skand. Fl. VII ed. p. 367 n. 48 (1861).

2. Ad rup. madid. — Eur. : Alp. Scand., Salisb., Stir., Carinth., Austr. inf., Frioul, Tirol., Bavar., Helvet., Pyren. As. : Caucas. Am. sept. : Groenland., M. Castkill.

conicum (Hsch.) Broth. op. et l. cc.

Bryum conicum Hsch. Fl. brasil I, p. 43 (1840); C. M. Syn. I, p. 301 (1849); Mitt. M. austr. am. p. 299 (1869); Par. op. cit. p. 175.

Br. Funckii Schwaegr. Sp. M. p. 69?

2. — Am. merid. : Brasil. austr. or.

cygnicollum (C. M.) Jaeg. Ad. I, p. 603 (1873-74).

Bryum Climacodontium Par. Ind. p. 173 (1894).

Br. cygnicollum C. M. in Bot. Zeit. 1848 (1).

Climacodontium cygnicollum Hpe. in Jaeg. Ad. II, p. 713 (1877-78).

— Afr. : Abyss.

cymbifolium (Lindb.) Broth. op. et l. cc.

A. filiforme Jaeg. op. cit. p. 604.

Brachymenium filiforme Griff. Not. p. 443 et Ic. pl. asiat. II, t. 102, f. 2 (1849).

Bryum cymbifolium Lindb. Man. M. I, p. 45 (1870); Par. op. cit. p. 177.

Br. filiforme Mitt. M. Ind. or. p. 67 (1859).

2. — As. : Himalaya, Nepal., Khasia, Java.

fabroniopsis (C. M.) Broth. op. et l. cc.

Bryum fabroniopsis C. M. in Hedwig. 1895, p. 127; Par. Suppl. Ind. p. 63 (1900).

2. Rupic. — Am. merid. : Brasil. or (Goyaz).

filiforme (Dicks.) Husn. Musc. gall. p. 222 (1888).

A. campestre Jaeg. op. cit. p. 602.

---

(1) Sec. Jaeg. l. c.; in C. M. Syn. deest.

- A. julaceum* W. P. Sch. *Syn. I* ed. p. 382 (1860); *Rab. Bryoth. eur.* n. 243, 363, 974 et 1219.  
*Bryum* filiforme *Dicks. Pl. crypt. fasc. IV*, p. 16 (*excl. syn.* (1802); W. P. Sch. *Syn. II* ed. p. 465 (1876); *Par. Ind.* p. 182.  
*Br. julaceum* Sm. *Fl. brit. III*, p. 1357, n. 13 p.p. (1804); *Brid. Bryol. univ. I*, p. 659 (1826); *Bryol. eur. fasc. 6-9* (Vol. IV), *Mon.* p. 79, t. 382 (1839); *C. M. Syn. I*, p. 315 (1849) et *II*, p. 577 (1851).  
*Webera julacea* de Not. *Epil.* p. 416 (1869).
2. *Ad rup. irrig.* — *Eur.* : Norveg. (ster.!), Britann., alp. Salisb., Stir., Carinth., Pinzgov., Tirol., Helvet., Vall. Tellin., Delphinat., Pyren. As. : China centr. (prov. Schen-Si). *Afr.* : Ins. Canar., Madeir., Azor.; Cameroon, Ruvenzori. *Am. sept.* : And. Nov. Granat. et Quitens., Brasil, Monte-Video.
- « *var. abyssinicum* (W. P. Sch.) *Broth. op. et l. cc.*  
*A. abyssinicum* Jaeg. *op. cit.* p. 602.  
*A. julaceum var. ? abyssinicum* W. P. Sch. *Syn. I* ed. p. 303, *nom.* (1860).  
*Bryum* filiforme *var. abyssinicum* *Par. Ind.* p. 182 (1894).  
— *Afr.* : M. Abyss.
- « *var. americanum* Ren. et Card. in *Bot. Gaz.* 1896, p. 51.  
*Bryum* filiforme *var. americanum* *Par. Suppl. Ind.* p. 63 (1900).  
— *Am. sept.* : Wisconsin.
- « *var. madagassum* eor. in *Bull. Soc. roy. bot. Belg.* 1894, *II*, p. 123; *Ren. M. Masc. Mad.* n. 223.  
*Bryum* filiforme *var. madagassum* *Par. op. et l. cc.*  
*Ad terr. arenac.* — *Afr.* : Madagascar.
- « *var. mexicanum* (W. P. Sch.) *Par.*  
*A. julaceum var. ? mexicanum* W. P. Sch. *op. et l. cc.*  
*A. mexicanum* Jaeg. *op. cit.* p. 603.  
*Bryum* filiforme *var. mexicanum* *Par. op. et l. cc.*  
— *Am. sept.* : Mexic. or.
- filiforme* Jaeg. — *A. cymbifolium*.  
*var. juliforme* Husn. — *A. juliforme*.  
*filinerve* (Mitt.) *Broth. op. et l. cc.*  
*Bryum* filinerve Mitt. *M. austr. am.* p. 305 (1869); *Par. op. cit.* p. 182.  
2. *Lignic.* — *Am. merid.* : And Quitens.
- humillimum* (C. M.) *Broth. op. et l. cc.*  
*Bryum* humillimum C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 26; *Par. Suppl. Ind.* p. 64 (1900).  
2. — *Am. merid.* : And Boliv.
- julaceum* W. P. Sch. — *A. filiforme*.  
*var. ? abyssinicum* W. P. Sch. — *A. filiforme var. abyssinicum*.  
*var. ? algerianum* W. P. Sch. — *A. juliforme*.  
*var. concinnatum* Zett. — *A. concinnatum*.  
*var. ? mexicanum* W. P. Sch. — *A. filiforme var. mexicanum*.
- juliforme* Solms Laub. *Tentam. bryogeogr. Algarv...* p. 38 (1867).  
*A. filiforme var. juliforme* Husn. *Musc. gall.* p. 222 (1888).  
*A. julaceum var. ? algerianum* W. P. Sch. *op. et l. cc.*  
*Bryum* campestre Lindb. *Man. M. I*, p. 41 (1870).  
*Br. filiforme var. neapolitanum* Boul. *M. Fr. I*, p. 293 (1884).  
*Br. juliforme* W. P. Sch. *Syn. II* ed. p. 465 (1876); *Par. Ind.* p. 189 (1894).  
*Webera neapolitana* de Not. *Epil.* p. 417 (1869).

2. Ad terr. humid. et subturfac. — *Eur.* : W. Norveg., Armor. marit., Gall. merid., Helvet., Etrur., Neapol., Corsica, Algarv. *As.* : Japon. *Afr.* : Alger, Madeir.

kashmirenses *Broth. op. et l. cc.*

*Bryum kashmirenses* *ej. in Act. Soc. sc. Fenn.* 1898, *II*, p. 24; *Par. Suppl. Ind.* p. 65 (1900).

2. — *As.* : Kashmir.

laceratum (*Besch.*) *Broth. op. et l. cc.*

*Bryum laceratum* *Besch. Fl. bryol. Réunion, etc.* p. 99 (1880-81); *Par. op. cit.* p. 190.

2. Terr. — *Afr.* : Ins. Borbon.

“ var. comorense (*Besch.*) *Par.*

*Bryum laceratum* var. comorense *Besch. op. et l. cc.*

Ut typ. — *Afr.* : Comor. magna.

leptoclados (*Sull.*) *Broth. op. et l. cc.*

*Bryum leptoclados* *Sull. in Proced. of the Amer. Acad.* 1861.  
*Plagiobryum leptoclados* *Par. Ind.* p. 957 (1896).

2. Terr. — *Am. sept.* : Cuba.

mexicanum *Jaeg.* — *A.* filiforme var. mexicanum.

nitidum (*Mitt.*) *Jaeg. op. cit.* p. 604.

*Bryum nitidum* *Mitt. M. Ind. or* p. 67 (1859); *Par. op. cit.* p. 198 (1894).

2. — *As.* : Himalaya, Ceylan.

obtusatissimum (*C. M.*) *Broth. op. et l. cc.*

*Bryum obtusatissimum* *C. M. in Nuov. Giorn. bot. ital.* 1897, p. 26; *Par. Suppl. Ind.* p. 68 (1900).

Ster. In aquos. — *Am. sept.* : Costa Rica.

orbiculatum (*Mitt.*) *Jaeg. op. cit.* p. 604.

*Bryum orbiculatum* *Mitt. M. austr. am.* p. 306 (1869); *Par. Ind.* p. 200 (1894).

Ster. Terr. — *Am. merid.* : M. Pichincha.

perappressum (*C. M.*) *Broth. op. et l. cc.*

*Bryum perappressum* *C. M. in Bull. hb. Boissier*, 1897, p. 182; *Par. Suppl. Ind.* p. 69 (1900).

Ster. — *Am. sept.* : Guatemala.

perimbricatum (*C. M.*) *Broth. op. et l. cc.*

*Bryum perimbricatum* *C. M. in Dusen M. Camer. n.* 362; *Broth. in Engler's Bot. Jahrb.* 1897, p. 245; *Par. Ind.* p. 204 (1894).

2. fr. ign. Muric. — *Afr.* : Cameroon.

procerrimum *Rehm. M. Afr. austr. n.* 540.

*Bryum procerrimum* *Par. op. cit.* p. 206.  
*Miehllichhoferia procerrima* *Rehm. in Coll. cit. n.* 219.

Ster. — *Afr.* : Prom. B. Sp., Transvaal.

prostratum (*C. M.*) *Jaeg. op. cit.* p. 603.

*Bryum prostratum* *C. M. Syn. I.* p. 317 (1849); *Mitt. M. austr. am.* p. 306 (1869); *Par. op. et l. cc.*

2. Saxic. — *Am. merid.* : And. Nov. Granat. et Quitens.

*prostratum* W. P. Sch. — A. soquense.

*pycnobaseum* (C. M.) Broth. op. et l. cc.

*Bryum pycnobaseum* C. M. in *Fl.* 1897, p. 329; *Par. Suppl. Ind.* p. 71 (1900).

Ster. — Am. merid. : Venezuela.

*quitense* W. P. Sch. in *Mandon Pl. Boliv. n.* 1664 bis.

*Bryum quitense* Par. Ind. p. 209 (1891).

— Am. merid. : And. Boliv.

*Schmidii* (C. M.) Jaeg. op. cit. p. 602.

*Bryum Schmidii* C. M. in *Bot. Zeit.* 1853, p. 24; *Mitt. M. Ind. or.* p. 68 (1859); *Par. op. cit.* p. 213.

2. — As. : M. Neilgherr.

*semi-ovatum* (Brid.) Jaeg. op. et l. cc.

*Bryum julaceum* Kth. *Syn. pl. Æquinoct.* p. 56 (1822-28).

Br. *semi-ovatum* Brid. *Bryol. univ. I.* p. 846 (1826); C. M. *Syn. I.* p. 316 (1849); *Mitt. M. austr. am.* p. 305 (1869); *Par. op. cit.* p. 214.

2. Saxic. et terr. — Afr. : Ins. Tristan da Cunha. Am. sept. : Costa Rica. Am. merid. : And. Quitens. et Peruv.

*semi-reticulatum* (C. M.) Broth. op. et l. cc.

*Bryum semi-reticulatum* C. M. in *Linn.* 1878-79, p. 301 et 1880-83, p. 377; *Par. op. et l. cc.*

Ster. — Am. merid. : Argentin. Saltens.

*sericeum* (de Lacroix) Husn. *Muscol. gall.* p. 223 (1838).

A. *leptostomoides* W. P. Sch. in *lit. et in Rab. Bryoth. eur. n.* 1218 (1873).

*Bryum filiforme* γ *sericeum* Boul. *M. Fr. I.* p. 293 (1884).

Br. *leptostomum* W. P. Sch. *Syn. II ed.* p. 469 (1876); *Par. op. cit.* p. 191.

2. Ad rup. humid. — Eur. : Alp. Rhaet., Gall. centr. (m. Dore Arvernicae).

*soquense* (Par.) Broth. op. et l. cc.

A. *prostratum* W. P. Sch. in *Mandon. Pl. Boliv. n.* 1667.

*Bryum cymbifolium* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 26.

Br. *soquense* Par. *Suppl. Ind.* p. 72 (1900).

— Am. merid. : Boliv.

*spongiosum* Jaeg. — Philenotis.

*terminale* (C. M.) Broth. op. et l. cc.

*Bryum terminale* C. M. in *Linn.* 1878-79, p. 303; *Par. Ind.* p. 218 (1894).

2. — Am. merid. : Argentin. Tucuman.

**Anomocladus** Mitt. *M. austr. am.* p. 567 (sect. ЛЕСКЕÆ) (1869). — 41 spec.

**ANOMODON** Hook. et Tayl. *Muscol. brit.* I ed. (1818).

*abbreviatus* Mitt. in *Trans. of the Linn. Soc.* 1891, p. 187.

A. *brevisetus* Broth. mss.

2. — As. : Japon.

*acutifolius* Mitt. *M. Ind. or.* p. 126 (1859).

2. — As. : W. Himalaya, Japon.



*angustatus* (Tayl.) Jaeg. *Ad. II*, p. 309 (1876-77).

*Hypnum angustatum* C. M. *Syn. II*, p. 472 (1851).

*Leskea angustata* Tayl. in *Lond. Journ. of bot.* 1846, p. 62.

*Porotrichum insularum* Mitt. *M. austr. am.* p. 461 ?

Ad arb. — *Am. merid.* : Nov. Granat., Quito.

*apiculatus* *Bryol. eur. fasc. 44-45* (Vol. V), *Mon.* p. 6, *nom.* (1850); *Sull. Ic. M.* p. 120, t. 75 (1864); *W. P. Sch. Syn. II ed.* p. 601 (1876).

*Hypnum Rügellii* C. M. *Syn. II*, p. 472 (1851).

*Leskea apiculata* W. P. Sch. *mss.*; C. M. *op. et l. cc.*

2. Ad arb. et rup. in montos. — *Eur.* : Sept. et med. : Suecia, Fennia, Siles., Rhoengeb., Thuring., Hassia, Bohem., Tatra, Stir., Bavar., Baden, Westphal., Helvet., Pyren. *As* : Saghalien, Japon., Caucas. occ. et centr. *Am. sept.* : Ottawa, Ontario, reg. sept. et or,

*armatus* *Broth. in Hedwig.* 1899, p. 243.

*A. rostratus* Besch. in *Faurie Pl. Japon. n.* 121.

2. — *As.* : Japon.

*asperifolius* C. M. in *Nuov. Giorn. bot. ital.* 1896, p. 117.

Ster. Inter *Lichenes*. — *As.* : China centr. (prov. Schen-Si).

*attenuatus* (Schreb.) Hüb. *Muscol. germ.* p. 562 (1833); *Bryol. eur. l. c.* p. 475; *W. P. Sch. op. cit.* p. 600.

*Hypnum arboreum erectum*, fruticuli specie, ramulis compressis *Dill. Cat. Giss.* p. 220 (1718).

*H. attenuatum* Schreb. *Spic. fl. Lips* p. 100, n. 1073 (1771); C. M. *Syn. II.* pp. 473 et 671 (1851).

*H. clavatum* Bell. *Append. ad Fl. pedemont.* p. 51.

*H. filicinum sericeum*, molle et pallidum mucronibus aduncis *var.* non splendens *Dill. Hist. M.* p. 287, t. 36, f. 22<sup>c</sup> (1741).

*H. fulvum* Brid. *Sp. M. II.* p. 151 (1812).

*H. heterophyllum aquaticum*, polycephalum, repens *var.* rigidior et ramosior *Dill. op. cit.* p. 293, t. 37, f. 27<sup>a</sup> (1741).

*H. repens filicinum ramosum*, ramulis surrectis et minus complanatis *ej. Cat. Giss.* p. 218 (1718).

*H. repens*, trichodes arboreum majus, caulibus ramosis *ej. Hist. M.* p. 331, t. 42, f. f. 66<sup>a-c</sup> (fid. C. M.).

*Leskia attenuata* Hedie. *Descript. I.* p. 23, t. 12 (1787).

*L. radicans* Schrank *Baiers. Fl. II.* p. 460 (1792).

*Neckera attenuata* Myr. *Coroll. fl. Upsal.* (1834).

1. Ad arb. et viar. cavar. marg. — *Eur.* : Reg. campestr. et montos. except. Britann. *As.* : Japon., Caucas. *Am. sept.* : S. E. Canada, Columb. brit., reg. sept., centr. or., et merid. *Pac.* ? (hb. de Poli).

« *var. brevifolius* Ren. et Card. in *Centralbl.* 1890, p. 51.

*Rupic.* ? — *Am. sept.* : Indiana, Illinois, Wisconsin, Ohio.

« *var. immersa* Ryan in D. K. D. *Vid. Saelsk. Skr.* 1896, n. I, p. 125.

— *Eur.* : Norveg.

*attenuatus* Sull. — *A. Wrightii.*

*brasiliensis* Hpe. — *Myrinia.*

*brevinervis* *Broth. in Oefv. af Finsha Vet.-Akad. Foerh.* 1890, p. 107.

Ster. Cortic. — *Pac.* : Austral. or. (Queensland).

*brevisetus* *Broth.* — *A. abbreviatus.*

*californicus* Lesq. in *Mem. Calif. Acad. I*, p. 30 ( ); Lesq. et Jam.  
*Man.* p. 306 (1884).  
 Ster. — *Am. sept.* : Calif.

*cladorhizans* Hüb. — *Eutodon Schleicheri*.  
*clavirameus* Jaeg. — *Pseudo-Leskea*.  
*compressus* Bruch. — *Entodon*.  
*consanguineus* Jaeg. — *Leskea*.  
*curtipendulus* H. et T. — *Antitrichia*.  
*devolutus* Mitt. — *A. Toccoae*.

*exilis* Mitt. in *Journ. of the Linn. Soc.* 1872, p. 309.  
 Ster. — *Afr.* : Natal., ins. Rodriguez.

*filiformis* Thw. et Mitt. in *Journ. of the Linn. Soc.* 1872, p. 308.  
 Ster. — *As.* : Ceylan.

*filiformis* Garov. — *Pterigynandrum*.

*filivagus* C. M. in *Fl.* 1890, p. 495.  
 Ster. — *Afr.* : M. Aberdare.

*flagelliferus* C. M. — *A. Toccoae*.  
*flexilis* Hpe. — *Papillaria illecebra*.  
*fragilis* Wils. — *Anomodon tristis*.

*Giraldii* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 273.  
 Ster. Inter *Lichenes*. — *As.* : China centr. (prov. Schen-Si).

*gracilis* Garov. — *Peterogonium*.  
*gracillimus* Jaeg. — *Schwetschkea*.  
*Grateloupitii* Mont. — *Schwetschkea*.

*heteroideus* Kindb. in *Mac. Cat.* p. 172 (1892).

*Leskea nervosa* var. *flagellifera* Kindb. in *Ott. Nat. IV*, p. 62 (1889).  
*L. nigrescens* ej. in *Bull. Torr. bot. Club, XVI*, p. 97 (1889); *Mac.*  
*Canad. M. n.* 395, et n. 252 p.p.

2. Ster. Ad saxa calc. et rad. *Acer.* -- *Am. sept.* : Ottawa, Ontario,  
 M. Rupestr.

*Huttonii* Mitt. in *Trans. and Proced. of the N. Zeal. Inst.* 1871,  
 pp. 229-232 et in *Journ. of the Linn. Soc.* 1872, p. 309.  
 Ster. — *Pac.* : Nov. Seeland.

*integerrimus* Mitt. *M. Ind. or.* p. 126 (1859).  
 Ster. — *As.* : Nepal., Himalaya.

*intricatus* Hpe. — *Pylaisia*.  
*janeirensis* C. M. — *A. Toccoae*.

*lagoënsis* Hpe. *Symb.* 1870, p. 283.  
 1. — *Am. merid.* : Brasil. austr. or.

*Leikipiae* C. M. — *Pseudo Leskea*.

*leptodontoides* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 274.  
 Ster. — *As.* : China centr. (prov. Schen-Si).

*leskeoides* (W. P. Sch.) Par. *Ind.* p. 44 (1893).

*Hypnum leskeoides* W. P. Sch. in *Drege Pl. capens. n.* 8158; C. M. in  
*Bot. Zeit.* 1855, p. 787.

1. — *Afr.* : Pr. B. Sp.

*longifolius* (Schleich.) Bruch in litt.; Rab. Deutsch. Kryptfl. II, 3, p. 255 (1848); Bryol. eur. fasc. 44-45 (Vol. VI), Mon. p. 3, t. 474 (1850); W. P. Sch. Syn. II ed. p. 599 (1876).

*Grimmia cylindracea* W. et M. Bot. Taschb. p. 157, fide Brid. (1807).

*Hypnum longifolium* C. M. Syn. II, pp. 474 et 691 (1851).

*H. repens trichodes arboreum majus*, cauliculis ramosis Dill. Hist. M. p. 331, t. 42, f. 66 (1741) et Hb. (fid. Limpr.).

*Leskea incurvata* Myr. Coroll. fl. Upsal (1834).

*L. longifolia* Spr. M. Pyr. exs. n. 87 (1847) et in Ann. and Mag. Nat. hist. 1849.

*Pterigynandrum longifolium* Schleich. Cent. IV pl. helv. n. 8 et Cat. 1807.

*Pt. nervosum*  $\beta$  *longifolium* Brid. Bryol. univ. II, p. 190 (1827).

*Pt. plumulosum*  $\beta$  *longifolium* ej. Mant. M. p. 129 (1819).

2. Ad arb., sax. et terr. lapidos. — Eur. : Centr. et sept., in zon. intermed. et montos. ; rar. fruct. As. : Jenissei super., Caucas. occ.

α var. *pumila* Milde in XLVII Jahresb. d. Schles. Gesells. 1870, p. 222. Ster. Rupic. — Eur. : Silesia.

*longinervis* Broth. in Hedwig. 1899, p. 243.

2. — As. : Japon.

*mexicanus* Jaeg. — *Leskea*.

*minor* (P. B.) Fürnrohr in Fl. XII, P. II, Ergaenz. p. 49 (1829); Sull. Ic. M. p. 119, t. 74 (1864).

*A. obtusifolius* Br. et Sch. in Drumm. M. bor. am. I ed. n. 163 et in Lond. Journ. of Bot. 1843, p. 668; Bryol. eur. fasc. 44-45 (Vol. V), Mon. pp. 3 et 6 (1850); Lesq. et Jam. Man. pp. 305 et 319 (1881).

*Hypnum stoloniferum* P. B. Prodr. p. 70 (1805).

*H. viticulosum*  $\beta$  *brachycarpum* C. M. Syn. II, p. 473 (1851).

*Isothecium minus* Sull. in Asa Gray Man. bot. U. St. I ed. p. 667 (1848).

*Neckera flaccida* (Haud C. M.) Brid. Sp. M. II, p. 137 (1812) et Mant. M. p. 139, n. 18 (1819); Schwaegr. Suppl. I, II, p. 151 (1816).

*N. minor* P. B. Prodr. p. 78 (1805).

*N. viticulosa*  $\beta$  *minor* Hedw. Sp. M. p. 210, t. 48, f. 6-8 (1801).

2. Cortic. — As. : Jenissei super., Japon. Am. sept. : New Brunswick, Gaspes., Ontario, Niagara, reg. sept., centr. et occid.

*Moseri* Kindb. — *Leskea*.

*mutabilis* Mont. — *Lescuraea striata*.

*nervosus* Hüb. — *Leskea*.

*obscurus* Jaeg. — *Leskea*.

*obtusifolius* } Bryol. eur. — *A. minor*.  
                      } Mac. — *A. platyphyllus*.

*ovicarpus* Besch. in Ann. sc. nat. 1893, p. 366.

2. Cortic. — As. : Japon.

*pellicola* C. M. in Hedwig. 1897, p. 127.

1. Saxic. — Am. merid. : And. Argentin. subtropic. Tucuman.

*platyphyllus* Kindb. Note on Canad. Bryol. 1893.

*A. obtusifolius* Mac. Canad. M. n. 256 et Cat. p. 171 (jure ! fid. Card.).

2. Ad rad. arb. in silv. humid. — Am. sept. : Ottawa, Ontario, Alabama.

*pellucinervis* Jaeg. — *Thuidium*.

*perpusillus* Garov. — *Habrodon*.

*planatus* Mitt. M. Ind. or. p. 126 (1859).

Ster. Filicic. — As. : Himalaya, Japon.

*pseudo-attenuatus* (C. M.) Jaeg. *Ad. II*, p. 304 (1876-77).

Hypnum *pseudo-attenuatum* C. M. in *Bot. Zeit.* 1855, p. 786.

1. — *Afr.* : Pr. B. Sp.

*pseudo-tristis* (C. M.) Par.

Hypnum *pseudo-triste* C. M. in *Bot. Zeit.* 1855, p. 786.

*Leskea pseudo-tristis* Jaeg. *Ad. II*, p. 298 (1875-76; ; *Par. Ind.* p. 745 (1896).

Ster. — *Afr.* : Prom. B. Sp.

*ramulosus* Mitt. in *Trans. of the Linn. Soc.* 1891, p. 187.

Ster. — *As.* : China, Japon.

*repens* Hüb.

var. *major* de Not. { — *Platygyrium repens*.

*rigidulus* Kindb. — *Leskea nervosa*.

*robustus* Rehm. — *A. Toccoae*.

*rostratus* (Hedw.) W. P. Sch. *Syn. I ed.* p. 488 (1860) *et II ed.* p. 598 (1876).

Hypnum *Caturum* P. B. *Prodr.* p. 62 (1805).

H. *desmiphorum* ej. *op. cit.* p. 63.

H. *flavens* Rich. in *Mich. Fl. bor. Amer.* (1803).

H. *imbricatum* P. B. *op. cit.* p. 65.

H. *julaceum, erectum sericeum repens, capsulis cylindraceutis* Dill. *Hist. M.* p. 322, t. 41, f. 58 c (1741).

H. *rostratum* P. B. *op. cit.* p. 69; C. M. *Syn. II*, pp. 475 et 691 (1851).

H. *sabinaefolium* Brid. *Mant. M.* p. 166 (1819).

*Leskea marginata* ej. *op. cit.* p. 133 *et Bryol. univ. II*, p. 203 (1827).

L. *imbricatula* Hedw. *Sp. M.* p. 224, t. 52 (1801).

L. *rostrata* ej. *op. cit.* p. 226, t. 55 (1801); *Brid. Bryol. univ. II*, p. 303; *Bryol. eur. fasc.* 41-45 (Vol. V), *Mon.* p. 4, t. 473 (1850).

*Pterigynandrum crinitum* Brid. *op. cit.* p. 192.

Pt. *marginatum* Rich. in *Mich. Fl. bor. Amer. II*, p. 292.

*Pterogonium marginatum* Schwaegr. *Suppl. I, P. I*, p. 168 (1811).

2. Ad arb. rad. et saxa calcar. — *Eur.* : ♂ ign.; reg. infer. jugi alp. : Austria, Salisb., Stir., Carinth., Carniole, Bavar., Tirol. Helvet., Ital. super., Pyren. *As.* : Caucas. occ. et centr., N. W. Himalaya, Panjab. *Am. sept.* : Uberr. fr. ! S. E. Canada, reg. sept., centr., or. et merid.; ins. Vancouver.

*rostratus* Besch. — *A. armatus*.

*rupestris* Kindb. — *Leskea*.

*scaberrimus* Broth. in *Bihang till K. Sv. Vet.-Akad. Foerh.* 1895, n. 3.

2. Cortic. — *Am. merid.* : Brasil. or.

*sciuroides* Broth. — *A. Toccoae*.

*sinensis* C. M. in *Nuov. Giorn. bot. ital.* 1896, p. 118.

Ster. — *As.* : China centr. (prov. Schen-Si).

*sinensi-tristis* C. M. in *op. et l. cc.*

Ster. Inter *Lichenes*. — *As.* : China centr. (prov. Schen-Si).

*stratosus* Jaeg. — *Pseudo-Leskea*.

*striatus* Hüb. — *Lescuraea*.

var. ♂ *saxicola* Hartm. — *Lescuraea striata* var. *saxicola*.

*subintegerimus* Broth. et Par. in *Rev. bryol.* 1900, p. 77.

Ster. Cortic. — *As.* : Tonkin. super.

*subpilifer* Lindb. et Arn. *M. As. bor.* 1890, II, p. 111.

2. fr. ign. Cortic. — *As.* : Jenissei.

*subrigidulus* Kindb. — Pseudo Leskea.

*tectorum* Kindb. — Pseudo Leskea.

*thranstus* C. M. in *Nuov. Giorn. bot. ital.* 1898, p. 207 (« an var. *As var. A. leptodontoides* ? » (C. M. l. c.)

Ster. — *As.* : China centr. (prov. Schen-Si).

*Toccoae* Sull. *M. of U. St.* p. 58 (1856); *Sull. et Lesq. in Sull. Ic. M.* p. 121, t. 76 A (1864); *Lesq. et Jam. Man.* p. 306 (1884).

*A. devolutus* Mitt. *M. Ind. or.* p. 127 (1859) *fid. Salmon*; *Par. Ind.* p. 43 (1894).

*A. flagelliferus* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 273; *Par. Suppl. Ind.* p. 14 (1900).

*A. jancirensis* C. M. in *Fl.* 1890, p. 496; *Par. Suppl. Ind.* p. 14.

*A. robustus* Rehm. *M. Afr. n.* 639 et 639\*; *Par. Ind.* p. 45.

*A. sciuroides* Broth. in *Bihang till K. Sv. Vet.-Akad. Föerh.* 1900.

? *Neckera sciuroides* Hpe. *Enum. M. brasil.* p. 43 (1879); *Par. Ind.* p. 858 (1896).

*Thamnium Toccoae* Kindb. *Eur. a. N. Amer. Bryin.* p. 44 (1897).

2. *Cortic. et rupic.* — *As.* : China centr. (prov. Schen-Si), Tonkin, N. W. Himalaya; Celebes, m. Javae, Sumatra. *Afr.* : Prom. B. Sp. *Am. sept.* : Georg. super., Ludovic. *Am. merid.* : Brasil. or.

*tonkinensis* Besch. in *Bull. Soc. bot. Fr.* 1894, p. 84.

*Rupic.* — *As.* : Tonkin.

« var. *leptoclados* ej. in *Rev. bryol.* 1898, p. 75.

*Ubi typ.*

*tristichus* W. P. Sch. — Triquetrella.

*tristis* (Cesati) Sull. *M. and Hep. of the U. St.* p. 58 (1856); *Milde in Bot. Zeit.* 1864, Beil. p. 15.

*Anomodon fragilis* Wils. in *Lond. Journ. of bot.* 1848, p. 275, t. 10.

*Haplohymenium Sieboldi* Dz. et Mh. in *Ann. sc. nat.* 1844, p. 310 et *M. frond. ined. Archip. ind.* p. 127, t. 40 (1851).

*Heterocladium triste* Kindb. *Eur. a. N. Amer. Bryin.* p. 46 (1897).

*Hypnum emarginatum* Schleich. in sched.

*H. fragile* C. M. *Syn. II.* p. 471 (1851).

*H. triste* ej. *op. cit.* p. 478.

*Leskea fragilis* H. f. et Wils. in *Drumm. M. bor. am. II ed. n.* 101.

*L. tristis* Cesati in *de Not. Syll. M.* p. 67 (1838); *W. P. Sch. Syn. II ed.* p. 596 (1876).

*Neckera Sieboldi* C. M. *op. cit.* p. 81.

2. *Cortic. et rupic.* — *Eur.* : Rhaet., Tirol. merid., Ital. super. *As.* : Japon., China or., Népal., Himalaya et Tibet. occid. *Am. sept.* : A Montreal usq. ad Ludoviciam.

*viticulosus* (L.) Hook. et Tayl. *Muscol. brit. I ed.* p. 79, t. 22 (1818); *Bryol. eur. fasc.* 44-45 (Vol. V.), *Mon.* p. 5, t. 47<sup>3</sup> (1850); *W. P. Sch. op. cit.* p. 601.

*Hypnum repens trichodes arboreum majus*, capitulis et surculis erectis, minus ramosis *Dill. Cat. Gissen.* p. 216 (1718).

*H. reticulatum* L. *hb. teste W. P. Sch.*

*H. subhirsutum*, viticulis gracilibus erectis, capsulis teretibus *Dill. Hist. M.* p. 347, t. 39, f. 43 (1711).

*H. viticulosum* L. *Sp. pl. I ed.* p. 1127 (1753); *C. M. Syn. II.* pp. 472 et 690 (1851).

*Leskea viticulosa* Spr. *M. Pyr. exs. n.* 100 (1847) et in *Ann. and Mag. of nat. Hist. 2. Ser. III.* p. 289 (1849).

*L. vivipara* Willd.

*Neckera Schlechtendalii* Piper de N. Schlechtendalii, 1838, p. 9, t. I.

*N. viticulaeformis* *ej. op. cit.* p. 8, t. I.  
*N. viticulosa* *Hedw. Fund. M.* p. 209, t. 48 (1782) et *Sp. M.* p. 209, t. 48 (1801).

2. Ad saxa, mur. praecip. antiq., arb. annosior. — *Eur.* : Tota, arctic. except. *As.* : China, Jenissei super., Nepal., W. Himalaya, Kashmir, Caucas., Minor. *Afr.* : Alger. *Am. sept.* : Canada merid. or., Wisconsin.

« *var. mexicanus* *Ren. et Card. in Bull. Soc. roy. bot. Belg.* 1899, II p. 238.

— *Am. sept.* : Mexic.

« *var. microphyllus* *Kindb. Laubm. Schoed. u. Norwieg.* p. 12 (1883).

*A. viticulosus f. depauperata* *Limpr. Laubm. Deutschl..... II*, p. 774 (1895).

— *Eur.* : Scand. in.

*Wichurae* *Broth. in Hedwig.* 1899, p. 245.

2. fr. ign. Cortic. — *As.* : Japon.

*Wrightii* *C. M. in Hedwig.* 1898, p. 262.

*A. attenuatus* *Sull. in Proceed. of the Amer. Acad...* 1861, p. 284.

2. Rupic. — *Am. sept.* : Cuba.

*xanthophyllus* *Hpe. et Ltz.* — *Lescuraea*.

**Anetis** *Kindb. Eur. and N. Am. Bryin.* p. 202 (sect. *CAMPYLOPODIS*) (1897).

**Antennidens** *C. M. in Linn.* 1875, p. 354. (sect. *FISSIDENTIS*, *CONOMITRII sensu Müller.*)

**ANTITRICHIA** *Brid. Mant. M.* p. 136 (1819). — 4 spec.

*brasiliensis* *Hsch.* — *Pilotrichum*.

*californica* *Sull. in Trans. of the Amer. Acad. . XIII*, p. 11 ( ) et *1c. M. Suppl.* p. 79, t. 59 (1874); *Lesq. et Jam. Man.* p. 291 (1884); *W. P. Sch. Syn. II ed.* p. 577 (1876).

*A. curtipendula* *Sull. in Pac. R. Rep. IV*, p. 189

*A. curtipendula var. hispanica* *W. P. Sch. in sched. et Coroll.* p. 109 (1856); *C. M. Syn. II*, p. 116 (1851).

*Macouniella californica* *Kindb. Check List. of Eur. and N. Amer. M.* (1894).

2. Ad arb., mur. et rup. — *Eur.* : Tirol. ital., Hispan. merid. *Afr.* : Alger. *Am. sept.* : Alaska, ins. Vancouver, Columb. brit., Washington, Idaho, Oregon, Calif.

« *var. ambigua* *Ren. et Card. in Bot. Gaz.* 1890, p. 59.

— *Am. sept.* : Oregon.

*curtipendula* (*L.*) *Brid. Mant. M.* p. 136 (1819) et *Bryol. univ. II*, p. 222 (1827); *Bryol. eur. fasc.* 44-45 (Vol. V), t. 469 (1850); *W. P. Sch. op. cit.* p. 576.

*Anomodon curtipendulus* *H. et T. Muscol. brit. II*, p. 79, t. 22 (1827).

*Cyrtopus curtipendulus* *Spr. M. Pyr. n.* 113 et in *Ann. and Mag. of nat. hist.* 1849, p. 151.

*Hypnum arboreum repens, capitulis reflexis, brevibus pediculis infidentibus* *Dill. Cat. Giss.* p. 220 (1718).

*H. curtipendulum* *L. Sp. pl.* p. 1128 (1753) et *Herb.*

*H. deniatum, curtipendulum, viticulis rigidis* *Dill. Hist. M.* p. 33, t. 43 f. 69 (1741).

*H. montanum* *Lam.*

*Neckera curtispindula* Timm. *Prodr. fl. megap.* p. 221, n. 801 (1788).  
*N. hamulosa* Vill. *Cat. meth.* p. 42 (1786).

2. Ad arb. silv. et rup. umbros. — *Eur.* : E reg. campestr. ad mont. zonae intermed. *As.* : Caucas. occ. *Afr.* : Ins. Canar., Madeir., alp. Abyss., prom. B. Sp. *Am. sept.* : Groenland, Terra nov., ins. Miquelon, lac. Super, ins. Vancouver, Washington, Oregon, Calif. *Am. merid.* : Patagon.

« *var. gigantea* Sull. et Lesq. *M. bor. am. II ed. n.* 356; *Lesq. et Jam. op. cit.* p. 291; *W. P. Sch. op. cit.* p. 577.

Ad arbor. — *Am. sept.* : Fret. Behring, Alaska, Sitka, Columb. brit., ins. Vancouver, Washington, Idaho, Calif.

*var. hispanica* W. P. Sch. — *A. californica*.

« *var. spinosa* C. Jens. *Consp. fl. Groenl.* p. 342 (1887).

— *Am. sept.* : Groenland.

*curtispindula* Sull. — *A. californica*.

*cuspidata* Nees. — *Papillaria illecebra*.

*kilimandscharica* Broth. in *Engler's Bot. Jahrb.* 1897, p. 254.

Ster. — *Afr.* : Kilima N'Djaro, reg. lacustr.

*oligoclada* Kindb. — Pseudo Leskea.

*pseudo-californica* Kindb. — *Alsia californica*.

*tenella* Kindb. in *Bull. Torr. bot. club. XVII*, p. 27 (1890) et in *Mac. Cat.* p. 165 (1892).

2 ♀ ign. Rupic. — *Am. sept.* : Ins. Vancouver.

**AONGSTRÆMIA** Bryol. eur. fasc. 33-36 (Vol. I) (1846). — 16 spec.

Spec. omn. non indicatae sunt DICRANELLÆ ejusd. nom.

*acerosa* Hpe. — *A. vaginata*.

*andicola* C. M. — *A. julacea*.

*apophysatula* C. M. — *Microdus*.

*austro-exigua* C. M. — *Microdus*.

*Baileyana* C. M. — *Microdus*.

*banatica* Hpe. — *Dicranella heteromalla*.

*bicolor* C. M. — *Oncophorus*.

*brevifolia* Hpe et Ltz. — *Rhamphidium*.

*brevipes* Hpe. mss.; *Besch. Prodr. bryol. mexic.* p. 19, in obs. (1871).

*Schlumbergeria vulcanica* W. P. Sch. mss.

*Weisia vulcanica* Hook. mss.

— *Am. sept.* : Mexic. (summi. M. Orizaba).

*Buchanani* C. M. — *Campylopodium*.

*capituligera* C. M. in *Linn.* 1876-79, p. 308.

*Dichodontium capituligerum* Par. *Suppl. Ind.* p. 114 (1900).

*Dicranella capituligera* ej. *Ind.* p. 327 (1895).

2.

— *Am. merid.* : Argentin. Cordob.

*convoluta* Hpe. in *Ann. sc. nat.* 1866, III p. 336.

*Anisothecium convolutum* Mitt. *M. austr. am.* p. 40 (1869).

*Dichodontium convolutum* Par. *Ind.* p. 322 (1895).

*Dicranella convoluta* Jaeg. *Ad. I*, p. 85 (1870-71).

1.

— *Am. merid.* : And. Nov. Granat.

*curvicaulis* C. M. — *Oncophorus*.  
*curviseta* Hpe. — *Campylopodium*.  
*cylindrica* C. M. — *Trichodon*.

*elegans* (Duby) Broth. *Natür. Pflanzenfam.* p. 308.

*Dichodontium elegans* Jaeg. *Ad. II*, p. 631 (1877-78); *Par. op. et l. cc.*  
*Dicranum elegans* Duby in *Mém... Genre*, 1869, t. I, f. I.

2. — *Am. merid.* : Chile.

*euchlora* C. M. — *Microdus*.  
*euphoroclada* C. M. — *Campylopodium*.  
*exigua* } C. M. — *Microdus*.  
           } Wils. — *Microdus pomiformis*.

*falcicaulis* C. M. in *Ren. Prodr. fl. bryol. Madag. etc.* p. 57 (1897).  
 — *Afr.* : Madagascar.

*Fendleri* C. M. — *Campylopodium*.  
*flaccidula* C. M. — *Microdus*.

*Gayana* (Mont.) C. M. *Syn. I*, p. 427 (1849); *W. P. Sch. Syn. I ed.*  
 p. 69 (1860); *Mitt. M. austr. am.* p. 27 (1869).

*Dicranum Gayanum* Mont. in *Ann. sc. nat.* 1845, *IV*, p. 112, in *Gay Hist. Chil. Bot. crypt.* t. 2, f. 3 (1850) et *Syll.* p. 45 (1856).

2. Terr. — *Am. merid.* : Chile austr.

*gracillima* C. M. — *Dicranella Bellii*.  
*Guilleminiana* Sull. — *Dicranella reticulata*.  
*gymna* C. M. — *Microdus*.  
*Harrisii* C. M. — *Microdus*.  
*hawaica* C. M. — *Microdus*.  
*Hilariana* C. M. — *Microdus*.  
*Hildebrandti* C. M. — *Microdus*.  
*Hookeri* C. M. — *Dicranella Jamesoni*.  
*hydrophila* C. M. — *Microdus*.  
*integra* C. M. — *Campylopodium*.  
*italiaiensis* C. M. — *Campylopodium*.

*jamaicensis* C. M. in *Bull. hb. Boissier*, 1897, p. 554.

*A. vulcanica* Mitt. *M. austr. am.* p. 28 (*quoad pl. jamaic.*).

Ster. — *Am. sept.* : Jamaic.

*julacea* (Hook.) Mitt. *M. austr. am.* p. 27 (1869).

*A. andicola* C. M. in *Bot. Zeit.* 1817, p. 189 et *Syn. I*, p. 428 (1849).

*Gymnostomum julaceum* Hook. *M. ex.* t. 42 (1820); *Brid. Bryol. univ. I*,  
 p. 92 (1826).

*Illecebraria julacea* Hpe *Prodr. Fl. Nov. Granat.* p. 354 (1865).

2. — *Am. merid.* : And. Nov. Granat. et Quitens.

*julaceo-divaricata* C. M. in *Ule Bryoth. brasil.* n. 60.

— *Am. merid.* : Brasil. or.

*lagunaria* C. M. — *Microdus*.

*Lamyi* Boul. — *Leptotrichum vaginans* var. *brevifolium*.

*longipes* (Somm.) *Bryol. eur. fasc.* 33-36 (*Vol. I*), *Mon.* p. 3, t. 94 (1846);  
*C. M. Syn. I*, p. 426 (1849); *W. P. Sch. Syn. II ed.* p. 69 (1876).

*Dicranum julaceum* H. f. et W. in *Drumm. M. bor. am.* n. 100.

*D. varium* eor. in *Coll. cit.* n. 99 p.p.

*Weisia longipes* Sommerf. *Suppl. ad Wahlenb. Fl. lapp.* p. 52, t. I,  
 f. 1-10 (1826).





2. Ad rup. basalt. — *Afr.* : Ins. Bourbon., Madagascar. *Am. sept.* : And. Quitens.

*vulcanica* Mitt. p. p. — *A. jamaicensis*.

*Wahlenbergii* C. M. — *Cynodontium virens* var. *Wahlenbergii*.

*Wrightii* C. M. — *Microdus*.

**Apalodictyon** C. M. *Syn. I*, p. 291 (sect. *BRYI*) (1849).

APALODIUM Mitt. *M. austr. am.* p. 15 (= *ORTHODONTIUM*) (1869).

APHANORHEGMA Sull. in *Asa Gray Man. bot. of U. St. I ed.* p. 647 (1856).

Spec. non indicatae sunt *PHYSCOMITRIA* ejusd. nom.

*Hampei* Lindb. — *Physcomitrella*.

*patens* Lindb. — *Physcomitrella*.

var. *anomalum* Hpe. — *Physcomitrella Hampei*.

APIOCARPA Hüb *Muscol. germ.* p. 154 (1833).

*elongata* ej. — *Mielichhoferia nitida* var. *elongata*.

*Mielichhoferi* ej. — *Mielichhoferia nitida*.

APLONDON R. Br. in *Neu Bot. Zeit. VII, Suppl. ad it. Parray.* p. 299.

*Wormskioldii* ej. — *Sphachnum*.

APODANTHUS la Pyl. in *Journ. bot. Desv. V.* p. 20, t. 33, f. 1 (1813).

*aphyllus* ej. — *Splachnum heteum*.

**Aporodictyon** Lindb. *M. scand.* p. 23 (sect. *DICRANI*) (1879).

**Aporotheca** Limpr. *Laubm. Deutschl.... II*, p. 615 (sect. *POLYTRICHI*) (1893).

APTYPHUS C. M. *Syn. II*, p. 325 (subsect. *HYPNI*) (1851), *postea* gen. propr.

Spec. omni. non indicatae sunt. *RHAPHIDOSTRIGIA* ejusd. nom.

*leucodontaceus* C. M. — *Pterogoniella*.

*molliculus* C. M. — *Isopterygium*.

*sarcophyllus* C. M. — *Pterogoniella*.

*tener* C. M. — *Isopterygium*.

**ARCHIDIUM** Brid. *Bryol. univ. I*, p. 747 (1826). — 26 spec.

*africanum* Mitt. in *Journ. of the Linn. Soc.* 1885, p. 299.

1. — *Afr.* : M. Usagara.

*alternifolium* W. P. Sch. — *A. phascoides*.

*amplexicaule* C. M. in *Linn.* 1880-82, p. 346.

— *Am. merid.* : Uruguay.

♂ var. *brevi-stolonaceum* C. M. in *op. cit.* p. 347.

Ut et ubi typ.

*Archavaletae* C. M. in *Fl.* 1888, n. 1.

— *Am. merid.* : Monte-Video.

*brisbanicum* Broth. in *Bailey Contr. Queensl. Fl. Bull. n. 7*, p. 3 (nom.

1891) et in *Oefv. af Finska Sv. Vet.-Akad. Foerh.* 1893, p. 35.

1. Ad terr. humid. — *Pac.* : Austral. or. (Queensland).

*Campylopodium* C. M. in *Hedw.* 1899, p. 52.

*A. compactum* ej. in *Rehm. M. Afr. austr. (forma tenerior)*.

— *Afr.* : Pr. B. Sp.

- capense *Hsch. in Linn.* 1841, p. 135; *C. M. Syn.* I, p. 13 (1849).  
— *Afr.* : Pr. B. Sp.
- chrysosporum *W. P. Sch. in Breut. M. capens.*; *Jaeg. M. cleistoc.* p. 9.  
— *Afr.* : Prom. B. Sp.
- chrysosporum *Rehm.* — *A. julicaule.*  
compactum *C. M.* — *A. Campylopodium.*  
Donnellii *Aust.* — *A. ohioëns* var. *Donnellii.*
- Durienanum *W. P. Sch. in sched.*; *Jaeg. Ad. I*, p. 6 (1869-70); *Besch. Cat. M. Alg.* p. 3 (1882).  
— *Afr.* : Alger.
- Ecklonianum *Hpe. in sched.*; *Jaeg. M. cleistoc.* p. 9; *C. M. in Hedwig.* 1889, p. 53.  
— *Afr.* : Pr. B. Sp.
- ephemeroides *C. M. in Linn.* 1880-82, p. 345.  
— *Am. merid.* : Uruguay.
- falcatulum *C. M.* — *A. Rehmanni.*
- Giberti *Mitt. in Journ. of the Linn. Soc.* 1886, p. 300.  
1. — *Am. merid.* : Monte-Video.
- Hallii *Aust. in Bull. Torr. bot. Club, VI*, p. 145 (1879).  
1. — *Am. sep.* : Kansas, Texas.  
« var. minus *Ren. et Card. in Bot. Gaz.* 191, p. 237.  
— *Am. sept.* : Ludovic.
- indicum *C. M. in Fl.* 1888, n. 1.  
— *As.* : Barman.
- julaceum *C. M. in Linn.* 1880-82, p. 347.  
— *Am. merid.* : Uruguay.
- julicaule *C. M. in Hedwig.* 1899, p. 53.  
*A. chrysosporum Rehm. M. Afr. austr.*  
— *Afr.* : Pr. B. Sp.
- laterale *Bruch in Fl.* 1846, p. 132; *C. M. Syn. I* p. 14 (1849).  
— *Afr.* : Pr. B. Sp., Natal.
- Lescurii *Aust.* — *Pleuridium alternifolium* var. *robustum.*
- longifolium *Lesq. et Jam. Man.* p. 50 (1884).  
*Syn.* — *Am. sept.* : Florid.
- Lorentzii *C. M. in Linn.* 1880-82, p. 343.  
— *Am. merid.* : Uruguay.
- ohioëns *Bryol. eur. I, Mon.* p. 3, nom. (1837); *Sull. Ic. M.* p. 16, t. 7 (1864); *C. M. Syn. II.* p. 517 (1851); *Lesq. et Jam. Man.* p. 50 (1884).  
*A. phascoides Sull. M. Allegh. exs. n.* 213; *C. M. Syn. I*, p. 13 p.p.; *H. f. et W. in Drumm. M. bor. am. II ed. n.* 12 et 13.  
1. Ad terr. — *Am. sept.* : New-York, New-Jersey, Pensilv., Ohio, Minnesota, Kansas, Alabama sept., Ludovic.  
« var. *Donnellii (Aust.) Lesq. et Jam. op. et t. cc.*  
*A. Donnellii Aust. in Bull. Torr. bot. Club, VI*, p. 190 (1889).  
— *Am. sept.* : Virgin., Florid.

*phascoides* *Brid. Bryol. univ. p. 747, t. suppl. 3* (1826); *Schwaegr. Suppl. III, t. 205* (1827); *Bryol. eur. (Vol. I), t. 8* (1837) *et t. 637* (1855); *C. M. Syn. I, p. 13 p.p.* (1849) *et II, p. 517* (1851) (1).

*A. alternifolium* *W. P. Sch. Syn. I ed. p. 28* (1860) *et II ed. pp. 23 et 810* (1876).

*Phascum alternifolium* *H. et T. Muscol. brit. p. 6, t. 5* (1818).

? *Ph. alternifolium*, caulibus capsuliferis nanis, sterilibus altioribus erectis, foliis lanceolatis alternis *Dick. Pl. crypt. fasc. I, p. I, t. I, fig. 2* (1785).

*Ph. Bruchii* *Spreng. in L. Syst. veg. IV, p. 142* (1827).

*Ph. globiferum* *Br. in Fl. 1825, p. 281, t. I.*

1. In ericet humid. — *Eur.* : Med. (in vall. Rhen. freq.) *et merid. rar. septentr.*

*phascoides* { *Drumm.* } — *A. ohioënsæ.*  
                  { *Sull.* — *A. ohioënsæ.*  
                  { — *A. tenerimum.*

*Ravenelii* *Aust. in Bull. Torr. bot. Club. VI, p. 145* (1879).

*Syn.* — *Am. sept.* : Carolin. (ambo), Florid.

*Rehmanni* *Mitt. in Journ. of the Linn. Soc. 1886, p. 300.*

*A. falciculatum* *C. M. in Rehm. M. Afr. austr. n. 429, b et d.*

1. — *Afr.* : Pr. B. Sp.

*sinense* *DR. in Bull. Soc. bot. Fr. 1862, p. 61.*

1. — *As.* : China or.

*stolonaceum* *C. M. in Fl. 1888, n. 1.*

*Syn. ?* — *Pac.* : Austral.

*subulatum* *C. M. in op. et l. cc.*

— *Afr.* : Pr. B. Sp.

*tenerimum* *Mitt. in Journ. of the Linn. Soc. 1864, p. 17.*

*A. phascoides* *Drumm. M. bor. am. II ed. n. 11.*

1. — *Am. sept.* : Ludovic.

**ARCTOA** *Bryol. eur. I (= DICRANUM).*

**Arctobryum** *Hagen in Tromsø Mus. Aarsheft. 1898-99, p. 222.* (subsect. *BRYI*).

**Areodictyon** *C. M. Syn. I, p. 46* (sect. *BRYI*) (1849).

**AREOFISSIDENS** *C. M. Syn. I, p. 46* (subsect. *FISSIDENTIS*); *Fleisch. gen. propr. in sched.*

*Naumannii* *Fl. — F. multiflorus.*

**Argyro-Barbula** *C. M. Syn. I, p. 597* (sect. *BARBULÆ*).

**ARGYRO-BRYUM** *Hpe. in Linn. 1876, p. 312.*

*subrotundum* *ej. — Bryum subrotundifolium.*

**Argyrothrix** *C. M. in Linn. 1871-73, p. 41* (sect. *MACROMITRII*).

**ARRHENOPTERUM** *Hedw. Sp. M. p. 198, t. 43, f. 1-9 (= AULACOMNIUM).*

---

(1) De nom. specif. cf. *Limpr. Laubm. Deutschl... I, p. 157.*

**ARTHROCORMUS** Dz. et Mk. M. frond. ined. Archip. ind. p. 75 (1854)  
— 2 spec.

Spec. omn. non indicatae sunt EXODICTYA ejusd. nom.

*africanus* Broth. — Octoblepharum.

*asper* Jaeg. — Leucophanes.

*Hookeri* Hpe. — Leucophanes glaucum.

*pulvinatus* Dz. et Mk. — Octoblepharum.

Schimperi (Dz. et Mk.) Card. in Rev. bryol. 1899, p. 7.

Miehlichhoferia Schimperi Dz. Mk. in Ann. sc. nat. 1844, II, p. 312.

M. trifaria eor. in Hb. Lugdun. Batav.

Octoblepharum Schimperi Mitt. in Voy. H. S. M. Challenger, III, p. 258 (1885).

2. Cortic. et terr. — As. : Ins. Philippin., Amboine, Borneo, Java. Pac. : Ins. Admiral.

thraustus C. M. in Micholitz M. Philipp. n. 182.

— As. : Mindanao. Pac. : Nov. Guinea.

**ASCHISMA** Lindb. Utkast naturl. grupper. p. 28 (1878). — 2 spec.

aethiopicum (Weho. et Duby) Lindb. op. et l. cc.

Ephemerum aethiopicum W. et Dy. in Mém. . . . Genève, 1871, p. 19, t. 3, f. 8; Par. Ind. p. 428 (1895).

2. Ad ter. humid. — Afr. : Angola.

carniolicum (Web. et Mohr) Lindb. op. et l. cc.

Acaulon carniolicum C. M. in Bot. Zeit. 1847, p. 100, Syn. I, p. 23 (1849) et II, p. 519 (1851); Bryol. eur. I, Mon. p. 12, t. 5.

Phascum carniolicum W. et M. Bot. Taschenb. pp. 69 et 450 (1807); Brid. Bryol. univ. I, p. 26 (1827); W. P. Sch. Syn. II ed. p. 19 (1876).

Systegium carniolicum de Not. Epil. p. 741 (1869).

1. rariss. syn. Ad terr. limos. — Eur. : Gall. merid. (Monspel.), Carniol. ; Neuwied? Helvet? Afr. : Tingit. Am. sept. : Texas occ.

« var. speciosum (Mor.) Limpr. Laubm. Deutschl. . . I, p. 196 (1886).

A. speciosum Fleisch. in Malpighia, III (1896).

Phascum speciosum Moris in sched.

— Eur. : Sardin.

speciosum Fleisch. — A. carniolicum var. speciosum.

**ASCHISTODON** Mont. in Ann. sc. nat. 1845, IV, p. 109.

Spec. omn. sunt DITRICHIA ejusd. nom.

**Asteriscium** C. M. in Linn. 1877-78 (sect. BARBULÆ).

**ASTOMIOPSIS** C. M. in Linn. 1880-82, p. 391 (= PLEURIDIOPSIS).

**ASTOMUM** Hpe in Fl. 1837, p. 285, p. p.

Spec. omn. non indicatae sunt PLEURIDIA ejusd. nom.

*axillare* Hpe. — Pleuridium nitidum.

*crispum* Hpe. — Systegium.

*cryptocarpum* Broth. — Systegium.

*cylindricum* Mitt. — Phascum tetragonum var. cylindricum.

*Drummondii* Kindb. — Hymenostomum rostellatum.

*exsertum* Broth. — Hymenostomum.

*Lecieri* Limpr. — Systegium.

- ludovicianum* Sull. — Systegium.  
*Mittenii* Bryol. eur. — Systegium.  
*multicapsulare* Bryol. eur. — Systegium.  
*nervosum* C. M. } — Pleuridium.  
                           } — Pleuridium Robinsoni.  
*nitidulum* Bryol. eur. — Systegium.  
   var. *pygmaeum* Lesq. — Systegium nitidulum var. *pygmaeum*.  
*polycarpum* Hpe. — Phascum cuspidatum.  
*rostellatum* Bryol. eur. — Hymenostomum.  
 ? *rostratum* Hpe. — Hymenostomum rostellatum.  
*subacaule* Jaeg. — Hymenostomum.  
*Sullivantii* Bryol. eur. — Systegium.

**ASTRODONTIUM** *Schwaegr. Suppl. I, p. 128 (1823).*

Spec. omn. non indicatae sunt LEUCODONTES ejusd. nom.

*indicum* Dz. et Mk. — Clastobryum.

**ASTROPHYLLUM** *Lindb. M. scand. p. 13 (1879).*

Spec. omn. non indicatae sunt MNIA ejusd. nom.

- Blyttii* Lindb. — Mn. stellare.  
*cuspidatum* Lindb. — Mn. affine.  
*marginatum* Lindb. — Mn. serratum.  
*medium* subsp. *curvatulum* Kindb. — Mn. curvatum.  
*orthorhynchum* var. *inclinatum* Kindb. — Mn. inclinatum.  
*pseudo-punctatum* Lindb. — Mn. subglobosum.  
*Seligeri* Lindb. — Mn. affine var. *elatum*.  
*silvaticum* Lindb. — Mn. cuspidatum.

**Ateleobryum** *Mitt. in Godman's Hist. nat. of I. Azor. p. 61 (sect. Bryi = Br. Notarisii) (1870).*

**ATRACTYLOCARPUS** *Mitt. M. austr. am. p. 13 (1869).*

Spec. omn. non indicatae sunt DITRICHIA ejusd. nom.

- alpinus* Lindb. — Metzleria.  
*brasiliensis* Broth. — Metzleria.

**ATRICHUM** *P. B. Prodr. p. 42 (1805). — 42 spec.*

*androgynum* (C. M.) Jaeg. *Ad. I, p. 703 (1873-74).*

- A. *polyphyllum* Rehm. *M. Afr. austr. n. 266 et 572.*  
*Catharinea androgyna* C. M. *Syn. I, p. 193 (1849).*

l. vel *syn.* *Rupic.* — *Afr. : Transvaal, Natal, Pr. B. sp.*

*angustatum* (Brid.) *Bryol. eur. fasc 21-22 (Vol. IV), Mon. p. 9 t. 411 (1844); W. P. Sch. Syn. II ed. p. 528 (1876).*

- A. *controversum* P. B. *Prodr. p. 42 excl. syn. (1805).*  
*Bryum Juniperi* foliis rugosis capsulis rectoribus *Dill. Hist. M. p. 302, t. 46, f. 19 (1741) et Hb.*  
*Br. juniperifolium Hoffm. Deutsch. Fl. II, p. 40, n. 37 in obs. ? (1795).*  
*Catharinea angustata* *Brid. Mant. M. p. 204 (1819) et Bryol. univ. II, p. 105 (1827); C. M. Syn. I, p. 193 (1849).*  
*C. controversa* *Roehl. Moosg. Deutschl. I, p. 206, n. 54 b (1800).*  
*Polytrichum angustatum* *Brid. Sp. M. I, p. 78 (1806); W. Arn. in Mém. Soc. Hist. nat. Paris, II, p. 320 (1825).*  
*P. controversum* *Brid. Muscol. rec. II, p. 78 (1798).*  
*P. cylindricum* *Sw. in Mühlb. Cat. pl. Amer. sept. p. 99 (1813).*  
*P. undulatum*  $\beta$  minus *Michx. Fl. bor. am. II, p. 295 (1803).*

2. Ad terr. aren. in ericet. — *Eur.* : Rar. : Scand. ? (*non ! fid. Limpr. II, p. 597*), Britann., Sili., Carinth., Rhaet., Pyren. *As.* : Japon., Caucas. occ. *Am. sept.* : Terra nov., New Brunswick, Ontario, reg. sept., centr., or. et merid.
- anomalum *Milde in Hedwig. 1869, p. 161.*  
*Polytrichum formosum (fid. Limpr. op. cit. II, p. 619).*  
*Ster. Rupic. — Eur. : Sudetes.*
- bogotense (*Hpe.*) *Besch. in Bull. hb. Boissier, 1894, p.*  
*Catharinea bogotensis Hpe. mss.*  
*— Am. merid. : And. Nov. Granat.*
- borbonicum *Besch. Fl. bryol. Réunion, etc. p. 108 (1880-81).*  
*— Afr. : Ins. Borbon.*
- controversum *P. B. — A. angustatum.*
- crispulum *W. P. Sch. in Savat. M. Jap. n. 630; Besch. in Ann. sc. nat. 1893, p. 351.*  
 2. ♂ et fr. ign. — *As.* : Japon.
- crispum (*Jam.*) *Sull. M. of U. St. p. 41 (1856), et Ic. M. p. 73, t. 46 (1864); W. P. Sch. Syn. II ed. p. 530 (1876); Lesq. et Jam. Man. p. 257 (1884).*  
*A. laxifolium Wils. mss.*  
*A. tortifolium Sull. mss.*  
*A. xanthopelma Mac. Canad. M. n. 463.*  
*Catharinea crispa Jam. in Proceed. of the Amer. Acad... 1855, p. 445.*  
 2. Ad rip. gramin. rivul. inter saxa et rup. — *Eur.* : Britann. (*ster.*)  
*Am. sept.* : New Jersey, Ontario, Columb. brit.
- var. densifolium (*Lindb.*) *Par. Ind. p. 52 (1894).*  
*Catharinea crispa var. densifolia Lindb. in Not. 1867, p. 150.*  
 ♀ et fr. ign. — *Eur.* : Anglia (Cheshire).
- fertile *Naw. — A. Hausknechtii.*
- flavisetum (*Wils.*) *Mitt. M. Ind. or. p. 150 (1859).*  
*Catharinea flaviseta Wils. in sched. p.p.*  
*— As. : Japon., Nepal., N. W. Himalaya.*
- gracile (*C. M.*) *Par. Suppl. Ind. p. 17 (1900).*  
*Catharinea gracilis C. M. in Nuov. Giorn. bot. ital. 1896, p. 92.*  
 2. — *As.* : China centr. (prov. Schen-Si).
- Hausknechtii *Jur. et Milde in Verh. d. K. K. zool. bot. Gesells. in Wien, 1870, p. 598.*  
*A. fertile Nawaschin in Hedwig. 1889, p. 359.*  
 ? *A. undulatum β attenuatum Bryol. eur. fasc. 21-22 (Vol. IV), Mon. p. 8 (1844).*  
*Catharinea anomala Bryhn in Bot. Not. 1886, p. 157.*  
*C. Hausknechtii Broth. Etud. distrib. M. Caucas. p. 4 (1884); Hagen in Rev. bryol. 1891, p. 3.*  
*C. lateralis Vaiz. Ann. of bot. II, p. 69 (1889).*  
*C. undulata var. Hausknechtii Div. et Jam. Stud. Handb. p. 500 (1896).*  
 1. Terr. — *Eur.* : Norveg. austr. or., Fennia, Rossia, Austria, Bavar., Helvet., Anglia. *As.* : Jenissei, Talysch, Japon., Caucas. occ. *Am. sept.* : Ins. Princip. Edouard, Cap Breton.

*hercynicum* P. B. — *Oligotrichum*.

*hirtellum* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1892, I, p. 168.

2. In silv. — *Am. sept.* : Costa Rica.

*laevifolium* (Lindb. et Arn.) Par. Ind. p. 52 (1894).

*Catharinea laevifolia* L. et A. M. As. bor. 1890, II, p. 11.

Syn. — As. : Jenissei.

*laxifolium* Wils. — *A. crispum*.

*leiophyllum* Kindb. in *Bull. Torr. Bot. Club. XVII*, p. 275 (1890) et in *Mac. Cat.* p. 148 (1892).

*Oligotrichum leiophyllum* ej. Eur. and N. Am. Bryin. p. 156 (1897).

2. Ad terr. in silv. humid. — *Am. sept.* : M. Selkirk, ins. Vancouver.

*leptocylindricum* (C. M.) Par. Suppl. Ind. p. 17 (1900).

*Catharinea leptocylindrica* C. M. in *Hedwig*. 1897, p. 338.

2. Ad terr. limos. — *Pac.* : Nov. Seeland., Austral. or. (Victoria).

*Lescurii* Jam. — *Bartramiopsis*.

*ligulatum* Mitt. — *A. Mülleri*.

*minutum* Par. — *Oligotrichum*.

*Moseni* (Broth.) Par. op. et l. cc.

*Catharinea Moseni* Broth. in *Bihang till K. Sv. Vet.-Akad. Foerh.* 1895 n. 3.

2. Rupic. — *Am. merid.* : Brasil. or.

*Mülleri* (C. M. et Hpe.) Jaeg. Ad. II, p. 243 (1874-75).

*A. ligulatum* Mitt. in *Kew Journ. bot.* 1856, p. 262 et in *Fl. Tasman. II*, p. 200 (1860) *ex ips!*

*Catharinea Mulleri* C. M. et Hpe. in *Linn.* 1853, p. 500.

2. — *Pac.* : Austral. or., ins. lord Howe, Tasman.

*Mülleri* W. P. Sch. — *A. Schimperii*.

*obtusulum* (C. M.) Jaeg. op. cit. p. 717 (1877-78).

*Catharinea obtusula* C. M. in *Fl.* 1878, n. 6.

Syn. Ad terr. — As. : China centr., N. W. Himalaya, Kashmir, Tchitral.

*Erstedtianum* (C. M.) Mitt. M. austr. am. p. 605 (1869).

*Catharinea Erstedtiana* C. M. Syn. II, p. 558 (1851).

2. — *Am. sept.* : And. Costa-Ricens.

« *var. longifolium* (Lindb.) Par. Ind. p. 53 (1893).

*Catharinea Erstedtiana var. longifolia* Lindb. in *Not....* 1867, p. 104, n. 155.

— *Am. sept.* : Honduras.

*pallidum* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1895, I, p. 63.

2. ♂ ign. — As. : Sikkim.

*parallelum* Mitt. in *Journ. of the Linn. Soc.* 1864, p. 48, t. 8; *Sull. Ic. M. Suppl.* p. 54 t. 38 (1874); *Lesq. et Jam. Man.* p. 258 (1884).

*Oligotrichum parallelum* Kindb. Eur. and N. Am. Bryin. p. 156 (1897).

*Polytrichum undulatum* Drumm. M. bor. am. 1 ed. p. 286.

2. Ad rup. madid. — *Am. sept.* : Alaska, M. Rupestr. et Selkirk.



- parvirosula* (C. M.) Par. *Suppl. Ind.* p. 17 (1900).  
*Catharinea parvirosula* C. M. in *Nuov. Giorn. bot. ital.* 1898, p. 163.  
 Ster. — As. : China centr. (prov. Schen-Si).
- pastasanum* Mitt. *M. austr. am.* p. 605.  
 2. — Am. merid. : And. Quitens.
- piriforme* Jaeg. — *Psilopilum*.  
*planifolium* (C. M.) Jaeg. *Ad. I*, p. 703 (1873-74).  
*Catharinea planifolia* C. M. in *Bot. Zeit.* 1868, n. 47-48.  
 Terr. — Am. merid. : Æquator.
- polycarpum* (C. M.) W. P. Sch. *mss.*  
*Catharinea Callibryon-brevilamellosa* C. M. *Syn. I*, p. 193 (1819).  
*C. polycarpa* C. M. *op. cit. II*, p. 558 (1851).  
 1. — Am. sept. : Mexic. Am. merid. : And. Nov. Granat.  
 et Quitens.
- polyphyllum* Rehm. — *A. androgynum*.
- pusillum* (C. M.) Par. *Suppl. Ind.* p. 17 (1900).  
*Catharinea pusilla* C. M. in *Hedwig.* 1897, p. 338.  
 2. — Pac. : Tasman.
- rigidum* Ltz. *Moosst.* p. 157 (1864).  
 2 ? Ad terr. arenac. — Am. merid. : Chile austr.
- rio-grandense* (Broth.) Par.  
*Catharinea rio-grandensis* Broth. in *Bihang till K. Sv. Vet.-Akad. Föerh.* 1900, p. .  
 1. Terr. — Am. merid. : Brasil. austr. or. (Rio Grande do Sul.)
- rhystophyllum* (C. M.) Par. *Suppl. Ind.* p. 17 (1900).  
*Catharinea rhystophylla* C. M. in *Nuov. Giorn. bot. ital.* 1896, p. 93.  
 2. — As. : China centr. (prov. Schen-Si).
- rosulatum* C. M. et Kindb. in *Mac. Cat.* p. 148 (1892).  
*Catharinea rosulata* Kindb. *Eur. and N. Amer. Bryin.* p. 154 (1897).  
 Ster. Terr. — Am. sept. : Columb. brit.
- runcinatum* (C. M.) Par. *op. et l. cc.*  
*Catharinea runcinata* C. M. in *Bull. hb. Boissier*, 1897, p. 174.  
 2. — Am. sept. : Guatemala.
- Schimperi* Jaeg. *Ad. I*, p. 703 (1873-74).  
 A. Mülleri W. P. Sch. in *Besch. Prodr. bryol. mexic.* p. 62 (1871).  
 2. — Am. sept. : Mexic. (Orizaba).
- Selwynii* Aust. in *Coult. Bot. Gaz. II*, p. 95 ( ).  
*Catharinea Selwynii* Kindb. *op. cit.* p. 153.  
 2. In fissur. rup. — Am. sept. : M. Selkirk, Columb. brit., Washington, Idaho, Oregon.
- semi-lamellosum* (C. M.) Par. *op. et l. cc.*  
*Catharinea semi-lamellosa* C. M. in *Hedwig.* 1897, p. 337.  
 2. — Pac. : Ins. Lord Howe.

*sideroloma* (C. M.) Par. op. et l. cc.

Catharinea *sideroloma* C. M. in op. et l. cc.

2. — Pac. : Austral. or. (Victoria).

*sithkanum* Kindb. — Bartramiopsis.

*subseriatum* Mitt. — A. undulatum var. subseriatum.

*subulirostrum* W. P. Sch. in Besch. Prodr. bryol. mexic. p. 61 (1871).

2. — Am. sept. : Mexic. (Orizaba).

*synoicum* (C. M.) Par. op. et l. cc.

Catharinea *synoica* C. M. in Bull. hb. Boissier, 1897, p. 549.

Syn. Terr. — Am. sept. : Jamaic.

*tenellum* (Roehl.) Bryol. eur. fasc. 21-22 (Vol. IV), Mon. p. 2, t. 412 (1844); W. P. Sch. Syn. II ed. p. 529 (1876) (1).

Catharinella *tenella* Roehl. in Ann. Wett. III, p. 234 (1814); Brid. Bryol. univ. II, p. 104 (1827); C. M. Syn. I, p. 191 (1849).

Polytrichum undulatum var. Sw. M. Succ. p. 78 (1798).

P. undulatum  $\beta$  minus Funck Moostaschb. p. 70 (1820); Wahlenb. Fl. lapp. p. 349 (1812).

P. undulatum  $\beta$  *tenellum* Wahlenb. Fl. Suec. II ed., II, p. 768 (1832).

2. In terr. arenos. limos. et stagn. exsicc. — Eur. : Britann., Scandin., German. sept., Batav., vall. Rhen. med., Salisb., Stir., Carinth., Tirol., Helvet.

*torquescens* W. P. Sch. in Besch. Prodr. bryol. mexic. p. 62 (1871).

— Am. sept. : Mexic. (Orizaba).

*tortifolium* Sull. — A. crispum.

*Ulei* (C. M.) Par. Suppl. Ind. p. 18 (1900).

Catharinea *Ulei* C. M. in Ule Bryoth. brasil. n. 38.

Terr. — Am. merid. : Brasil. subtropic. or.

*undulatiforme* Ren. et Card. in Bull. Soc. roy. bot. Belg. 1892, I, p. 168.

2. Ad terr. — Am. sept. : Costa-Rica.

*undulatum* (L.) P. B. Prodr. p. 42 (1805); Bryol. eur. l. c. Mon. p. 8, t. 410; W. P. Sch. op. cit. p. 528.

Bryum erectum, capitulis oblongis, rubentibus foliis, oblongis, angustis pellucidis rugosis Dill. Cat. Giss. p. 222 (1718).

Br. phyllitidifolium Neck. Meth. M. p. 203 (1771).

Br. Phylliditis folio rugoso acuto, capsulis incurvis Dill. Hist. M. p. 360, n. 18, t. 46, f. 18 (1741) et Hb.

Br. undulatum L. Sp. pl. II, p. 117, n. 10 (1753).

Callibryum polytrichoides Weib. Prim. fl. Werth. p. 290 (1799).

C. undulatum Zenk. et Dietr. M. Thur. n. 41 (1822).

Catharinea *Callibryon* Ehrh. in Hann. Mag. 1780, p. 934; C. M. Syn. I, p. 192 (1849).

C. Ehrhardti Voit, M. herbip. p. 17 (1812).

C. undulata Web. et Mohr, Ind. pl. crypt. (1803); Brid. Bryol. univ. II, p. 102 (1827).

Mnium undulatum Sw. Meth. M. p. 27 (1781).

Oligotrichum undulatum Lam. et DC. Fl. fr. III et I, II, p. 492 (1803).

Polytrichum undulatum Hedw. Fund. II, p. 90 (1782).

(1) Cl. Limpricht (Laubm. Deutschl. etc., II, pp. 455 et 598) eadem pr. max. p. synonyma *Mnio orthorhyncho* et *Atricho tenello* assignat.

1. In silv., virgult, mur. delaps. etc. — *Eur.* : Fere tota. *As.* : Japon., China. Cochinch., Pers., Caucas., Minor. *Afr.* : Alger., Madeir., ins. Azor. *Am. sept.* : Terra nov., ins. Miquelon, M. Rupestr., Columb. brit., tota temper.
- var. *abbreviatum* Bryol. eur. — *A. undulatum* var. minor.
- « var. *alte-cristatum* Ren. et Card. in Bot. Gaz. 1890, p. 58.  
— *Am. sept.* : Pensilv., Ohio, Missouri, Michigan, Idaho, Oregon, Kansas.
- « var. *attenuatum* Bryol. eur. l. c.  
Catharinea Callibryon  $\beta$  attenuata C. M. op. et l. cc.  
Polytrichum undulatum var. I H. f. et Wils. in Drumm. M. bor. am. I ed. n. 66.  
Ut et ubi typ. in *Eur.* et *Am. sept.*
- « var. *gracilisetum* Besch. in Ann. sc. nat. 1893, p. 351.  
— *As.* : Japon.
- « var. minor (Hedw.) Web. et Mohr, Bot. Taschb. p. 217 (1807); Brid. Bryol. univ. II, p. 104 (1827).  
*A. undulatum*  $\gamma$  abbreviatum Bryol. eur. l. c. p. 8 (1844).  
Bryum juniperifolium Hoffm. Deutsch. Fl. II, p. 40, n. 37 in obs. ?  
Catharinea Callibryon  $\beta$  abbreviata C. M. l. c. p. 193.  
C. undulata  $\beta$  abbreviata Rab. Deutsch. Kryptfl. II, P. III, p. 233 (1848).  
C. undulata  $\beta$  corsica de Not. Syll. n. 214 (1838).  
Oligotrichum undulatum  $\beta$  minus Lam. et DC. op. et l. cc.  
Polytrichum undulatum  $\beta$  minus Hedw. Desor. I, p. 43, t. 17, f. 14-21 (1787).  
Ut et ubi typ. in *Eur.*; *As.* : Japon. *Am. sept.* : Kansas.
- « var. *rivularis* (Bryhn) Par.  
Catharinea undulata var. rivularis Bryhn in Kgl. Norske Vid. Selsk. Skrift. n. 3, p. 43 (1899).  
— *Eur.* : Norveg. bor.
- « var. *subserratum* (Hook.) Par. Ind. p. 55 (1894).  
*A. subserratum* Mitt. M. Ind. or. p. 150 (1859).  
Catharinea Callibryon  $\epsilon$  subserrata C. M. op. et l. cc.  
C. serrata Lindb. in Not... 1867, p. 101, n. 51.  
Polytrichum angustatum Griff. Not. p. 486, p.p. ? (1849).  
P. undulatum var. subserratum Hook. in Lond. Journ. of bot. 1840, p. 3.  
— *As.* : Nepal., N. W. Himalaya.
- xanthopelma Jaeg. Ad. I, p. 705 (1873-74); Lesq. et Jam. Man. p. 237 (1884).  
Catharinea xanthopelma C. M. in Fl. 1873, p. 482.  
2. — *Am. sept.* : Arkansas, Texas, Kansas, Ludovic.
- xanthopelma Mac. — *A. crispum*.

**AULACOMITRIUM** Mitt. in Trans. of the Linn. Soc. 1891, p. 161. (3 spec.)

evanescens Broth. — *A. humillimum*.

humillimum Mitt. in op. et l. cc.

*A. evanescens* Broth. in hb. Berol.

Macromitrium humillimum Par. Ind. p. 777 (1895).

Cortic. et rupic. — *As.* : Japon.

parvulum (Wils.) Par. Suppl. Ind. p. 18 (1900).

A. calycinum Mitt. in op. et l. cc.

Macromitrium calycinum ej. M. Ind. or. p. 49 (1859); Par. Ind. p. 773.

M. parvulum Wils. in Kew Journ. bot. IX, p. 327 (1857).

1. — As. : Japon., Ceylan.

Warburgii Broth. in Hedwig. 1899, p. 215.

1. Ad Theam. — As. : China.

**AULACOMNIUM** Schwaegr. Suppl. III, P. 1, fasc. I. t. 215 (1827).  
(9 spec.)

acuminatum (Lindb. et Arn.) Par. Ind. p. 55 (1894).

Sphaerocephalus acuminatus L. et A. M. As. bor. 1890, II, p. 27.

2. — As. : Jenissei med.

androgynum (L.) Schwaegr. op. cit. Mon. p. 2, n. 1; Bryol. eur. fasc. 10 (Vol. IV), Mon. p. t. 406 (1841); W. P. Sch. Syn. II ed. p. 503 (1876).

Bryum androgynum Web. Spic. Fl. Goett. p. 122, n. 191 (1778).

Fusiconia androgyna P. B. in Mem. Soc. Linn. Paris, I, t. VII, f. 5 (1822).

Gymnocephalus androgynus Rich. in Schwaegr. Suppl. I, P. II, p. 87, n. 1 (1816).

Gymnocybe androgyna Fries, St. agr. fensj. p. 27 (1825).

Hypnum androgynum Schrank Bayer. Fl. II, p. 469 (1789).

Mnium androgynum L. Sp. pl. I ed. II, p. 1110, n. 1 (1753); Brid. Bryol. univ. II, p. 5 (1827); C. M. Syn. I, p. 170 (1849); Lindb. M. scand. p. 14 (1879).

Mn. angustifolium Neck. Meth. M. p. 234, n. 2 (1771).

Mn. perangustis et brevibus foliis Dill. Cat. Giss. p. 214 (1718) et App.

p. 84, t. 1, f. A.-C. (1719); Hist. M. p. 230, t. 31, f. 1 (1741) et Hb.

Orthopyxis androgyna P. B. Prodr. pp. 32 et 78 (1805).

2. Ad rup. quarz., terr., rad. arb. et trunc. putrec. in palud. Eur. : Tota campestr. et montos.; fruct. frequentior in zon. sept. Am. sept. : Terr. nov., Nov. Scot.; Alaska, ins. Vancouver, Columb. brit., Washington, Oregon, Idaho, Montana, Tennessee, Ludovic., Calif.

chilense C. M. — Rhizogonium mnioides.

flexifolium H. f. et W. — Zygodon squarrosus.

Gaudichaudi Mitt. — Leptotheca.

heterostichum (Hedw.) Bryol. eur. fasc. 10 (Vol. IV), Mon. p. 8, t. 403.

Arrhenopterum heterostichum Hedw. Sp. M. p. 198, t. 46, f. 1-9 (1801).

Bryum heteropterum pellucidum Dill. Hist. M. p. 362, t. 45, f. II (1741).

Br. heterostichum W. Arn. in Mem. Soc. hist. nat. Paris II, p. 292.

Hypnum heterostichum W. et M. Ind. pl. crypt. (1803).

Mnium Arrhenopterum Sm. in Trans. of the Linn. Soc. 1814, p. 263 p.p. (fid. C. M.).

Mn. heterostichum Sw. in Mühl. Cat. pl. Am. sept. p. 94; Brid. Bryol. univ. II, p. 12 (1827); C. M. Syn. I, p. 169.

Orthopyxis heterosticha P. B. Prodr. p. 79 (1805).

1. Ad terr. humid. — As. : Japon. Am. sept. : Ontario, reg. sept., centr., or. et merid.

“ var. humilis (C. M.) Par. Ind. p. 56 (1894).

Bryum membranaceum foliis membranaceis obtusis Dill. Hist. M. p. 551, t. 85, f. 2 (1741).

Mnium heterostichum var. humile C. M. l. c. inter. syn.

Mn. membranaceum Brid. Musc. rec. II, P. III, p. 85 (1803) et Bryol. univ. II, p. 13 (1827).

Orthopyxis membranacea var. humilis Lindb. de Mn. europ. p. 78 (1868).

Ut et ubi typ.

*marginatum* (Aongstr.) Jaeg. Ad. II, p. 715 (1877-78).

*Gymnocybe marginata* Aongstr. in Oefv. af K. Vet.-Akad. Foerh. 1876, n. 4, p. 20.

Ster. — Am. merid. : Brasil. austr. or.

*palustre* (L.) Schwaegr. Suppl. III, P. I, fasc. I, Mon. p. 4, n. 2 (1827); *Bryol. eur. fasc. 10* (Vol. IV), Mon. p. 9, t. 405 (1841); W. P. Sch. Syn. II ed. p. 504 (1876).

*Bryum inordinatum* Brid. Mant. M. p. 117 (1819).

Br. *palustre* Neck. Meth. M. p. 210, n. 22 (1771).

Br. *ventricosum* (haud Dicks.) Sibth. Fl. oxon. p. 289 (1794).

*Gymnocephalus inordinatus* Brid. Sp. M. III, p. 3 (1817).

G. *palustris* Schwaegr. Suppl. I, P. II, p. 87 (1816).

*Gymnocybe palustris* Fries, St. agr. femsj. p. 27 (1825).

G. *palustris* var. *inordinata* Lindb. de Mn. europ. p. 87 (1868).

*Hypnum elodes* (haud Spr.) W. et M. Bot. Saschb. p. 272, n. 2 (1807).

H. *palustre* (haud Huds.) cor. Ind. mus. pl. crypt. (1803).

H. *palustre erectum* brevius, ramis longioribus, foliis angustioribus Dill. Cat. Giss. p. 219 (1718).

*Linnobryum palustre* Rab. Kryptfl. Tachs. I, p. 502 (1863).

Mnium *inordinatum* Brid. Musc. rec. II, P. III, p. 76 (1803) et *Bryol. univ.* II, p. 7 (1827).

Mn. majus, ramis longioribus bifurcatis Dill. Hist. M. p. 233 bis, 239, t. 31 f. 3, 4 et 8 (1741).

Mn. *palustre* L. Sp. pl. I ed. II, p. 1110, n. 3 (1753) et Hb. p.p.

Mn. ramis brevibus inordinate progredientibus Dill. op. cit. p. 239, t. 31, f. 8 (1741) (fide C. M.).

Mn. *reclinatum* Sic. in Act. Soc. Linn. VII, p. 262.

*Orthopyxis palustris* P. B. Prodr. pp. 32 et 79 (1805).

*Sphaerocephalus palustris* Lindb. M. scand. p. 14 (1879).

2. In pratis paludos. et silvat. uligin. — Eur. : E reg. campestr. usq. in subalp. ; Spitzberg, ins. Ursor., Septem Ins. As. : Penins. Samo-jed., Jenissei, Obi, Kamschatka, Okhotsk, Amour, Kouril., Japon., Tian Shan, Alatau, Kashmir, Baltistan, Caucas. occ., Syr. Afr. : Alger., alp. Abyss. Am. sept. : Groenland, Labrador, fret. Hudson et Kotzebue, Terra nov., ins. Miquelon, Canada merid. ; Alaska, Sitka, M. Rupestr., Columb. brit., ins. Vancouver, Washington; reg. sept., centr., occ. et or. Pac. : Tasman.

« var. *alpestris* W. P. Sch. Coroll. p. 81 (1856) et Syn. II ed. p. 505.

*Gymnocybe palustris*  $\xi$  *alpestris* Lindb. op. et l. cc.

In turfis. uligin. — Eur. : Helvet. (Grimsel). Am. sept. : Reg. montana.

« var. *brentelioides* Mol. Moosst. Alg. p. 82 (1865).

Ad rup. aren. madid. — Eur. : Alp. Algov.

« var. *congestum* Boulay, M. Fr. p. 224 (1884).

— Eur. : Alp. Saband. (m. Blanc), Pyren. Am. sept. : Ins. Miquelon.

« var. *fasciculare* (Funck) Bryol. eur. et W. P. Sch. II. cc.

*Gymnocybe palustris*  $\delta$  *fascicularis* Lindb. op. et l. cc.

Mnium *fasciculare* Funck in Brid. Bryol. univ. II, p. 733.

Mn. *palustre* var. *fasciculare* C. M. op. et l. cc.

In uligin. — Eur. : Alp. editor. Am. sept. : Groenland, reg. montana.

« var. *imbricatum* Bryol. eur. et W. P. Sch. II. cc.

*Gymnocybe palustris*  $\gamma$  *imbricata* Lindb. op. et l. cc.

Mnium *palustre* var. *imbricatum* C. M. op. et l. cc.

Ubi praeced. in Eur. ; Am. sept. : Terra nov., Ontario, Columb. brit. ins. Vancouver.

« *var. laxifolium* Kindb. in *Mac. Cat.* p. 145 (1892); et in *Dix. et Jam. Stud. Handb.* p. 282 (1896)?

Ad terr. humid. — *Eur.* : Anglia (Bedfordshire)? *Am. sept.* : M. Selkirk.

« *var. polycephalum* (Brid.) Hüb. *Muscol. germ.* p. 313 (1833); *Bryol. eur. et W. P. Sch.* II. cc.

*Bryum androgynum* β *prolixum* Roth, *Fl. germ.* III, p. 229 (1800).

*Br. palustre* β *ramosum* Turn. *Muscol. hibern.* p. 113 (1804).

*Gymnocybe palustris* ε *ramosa* Lindb. *op. et l. cc.*

*Limnobryum palustre var. polycephalum* Rab. *op. cit.* p. 303.

*Mnium majus, minus ramosum, capitulis pulverulentis crebrioribus* Dill. *Hist. M.* p. 285, t. 31, f. 4 (1741).

*Mn. palustre* β *flagelliferum* Mart. *Fl. crypt. Erlang.* p. 65 (1817).

*Mn. palustre* β *majus* Schwaegr. *Suppl. I, P. II.* p. 122 (1816).

*Mn. palustre* δ *polycephalum* C. M. *op. et l. cc.*

*Mn. polycephalum* Brid. *Musc. rec. II, P. III,* p. 76 (1803) et *Bryol. univ. II,* p. 10 (1827).

*Mn. prolixum* Neck. *Meth. M.* p. 135 (1771).

*Mn. ramosum* Huds. *Fl. angl.* p. 403 (1762).

*Orthopyxis ramosa* P. B. *Prodr.* p. 79 (1805).

In turf. depress. — *Eur.* : Praecip. sept. *As.* : Caucas. occ. *Am. sept.* : Groenland, Labrador, Terra nov., New Brunswick, Ontario,

M. Rupestr., ins. Vancouver, reg. montana.

« *var. submersum* Sanio in sched. et in *Limpr. Laubm. Deutschl...* II, p. 529 (1893).

In stagn. — *Eur.* : Borussia. or.

*papillosum* (C. M.) Jaeg. *Ad. II,* p. 715 (1877-78); *Lesq. et Jam. Mun.* p. 253 (1884).

*Mnium papillosum* C. M. in *Fl.* 1875, p. 93.

2. — *Am. sept.* : Colorado.

*pentastichum* Mont. — *Stenomitrium.*

*stolonaceum* C. M. in *Abhandl. nat. Ver. Brem.* 1900, p. 503.

Ster. — *Pac.* : Nov. Seeland.

*turgidum* (Wahlenb.) Schwaegr. *Suppl. III, P. I, fasc. I, Mon.* p. 7, n. 3 (1827); *Bryol. eur. fasc. 10, Mon.* p. , t. 404 (1841); *W. P. Sch. op. cit.* p. 506.

*Arrhenopterum turgidum* Wahlenb. *ms.*; W. et M. *Bot. Taschb.* p. 318 in obs. (1807).

*Bryum turgidum* Sw. *Summ. veg. scand.* p. 41 (1814).

*Gymnocybe turgida* Lindb. in *Not. ur Saellsk. p. Faun. et Fl. fennica Foerh.* IX, p. 85 (1867).

*Hypnum illecebrum* L. *Fl. Suec. II ed.* p. 399, n. 1032 excl. syn. (1755).

*H. turgidum* Wahlenb. *ms.*; W. et M. *Ind. mus. pl. crypt., nom. solum* (1803).

*Mnium Arrhenopterum* Sm. in *Trans. of the Linn. Soc.* 1801, p. 263 (id. *Limpr.*).

*Mnium turgidum* Wahlenb. *Fl. lapp.* p. 351, n. 619, t. 23 (1812); *Schwaegr. Suppl. I, P. II,* p. 123, t. 77; *Brid. Bryol. univ. II,* p. II (1827); *C. M. Syn. I,* p. 171 (1849).

*Sphaerocephalus turgidus* Lindb. *M. scand.* p. 14 (1879).

2. Ad terr. turf. vel ad rup. irrig. — *Eur.* : Alp. reg. sept., rar. med. : Spitzberg. Septem Ins., ins. Ursor., Lappon., Norveg., Scot., Tatra, Salisb., Tirol.; Anglia et Stir. (ster!). *As.* : Penins. Samojed. et Tschutschica, Taimyr, Jenissei, Obi, Ochotsk, Japon., Turkestan sept. *Am. sept.* : Groenland, Labrador, ins. Melville, fret. Raffen, Hudson, Kotzebue, Smith et Behring, lac. Super., M. Alb. et Adirondack, Nov. Anglia.

α *var. elongatum* (Lindb. et Arn.) Par. Ind. p. 58 (1894).

*Sphaerocephalus turgidus var. elongatus* L. et A. M. As. bor. 1890, II, p. 27.

— As. : Jenissei infer.

α *var. proliferum* Geheeb in Fl. 1879, p. 476.

— As. : Alatau.

*venezuelanum* Dz. et Mh. in Mitt. M. austr. am. p. 317 (1869).

Ster.

— Am. merid. : Venezuela.

**AULACOPILUM** Wils. in Lond. Journ. of bot. 1848, p. 90. — 6 spec.

*abbreviatum* Mitt. in Journ. of the Linn. Soc. 1872, p. 308, t. VD.

1. Filic. — As. : Ind. or.

*Balansae* C. M. — A. paraguense.

*glaucum* Wils. in op. cit. p. 9, I, t. 4 ; C. M. Syn. II, p. 186 (1851).

Ad arbor. — Afr. : Austral. (fid. Mitt.). Pac. : Nov. Seeland.

*incanum* Mitt. in op. et l. cc.

Ad arbor. — Afr. : Austral.

*paraguense* Besch. in Rev. bryol. 1885, p. 18.

A. Balansae C. M. in Fl. 1887, p. 447.

1. Ad Aurant. — Am. merid. : Paraguay.

*trichophyllum* Aongstr. mss. ; C. M. in Bot. Zeit. 1862, p. 393.

1. — Afr. : Pr. B. Sp.

*tumidulum* Thw. et Mitt. in Journ. of the Linn. Soc. 1872, p. 307, t. VC.

1. Ad arbor. — As. : Ceylan.

*Wildii* Broth. — Wildia solmsiellacea.

**Aureolaria** Kindb. Eur. and N. Am. Bryin. p. 86 (sect. CAMPTOTHECII).

**AUSTINIA** C. M. in Linn. 1875, p. 439. — 1 spec.

*tenuinervis* (Mitt.) C. M. in op. et l. cc.

*Clasmatodon parvulus* Sull. in Proceed. of the Amer. Acad... 1861, p. 283  
et in Wright M. cubens. n. 65.

Cl. trichelyma Sull. mss.

*Hypnum tenuinerve* Mitt. M. austr. am. p. 549 (1869).

1 ? Cortic. — Am. sept. : Cuba.

**Barbella** C. M. in Fl. 1896, p. 464 (sect. PILOTRICHLLÆ, nom.).

**BARBULA** Hedw. Fund. M. II, p. 92 p.p. (1782), reduc. Vent. in Comm.

Fauna, Fl. etc. n. 3, Venezia, 1868. — 232 spec.

Spec. omn. non indicatae sunt TORTULÆ ejusd. nom.

*abbreviatifolia* H. Müll. — B. icmadophila.

*acrophylla* C. M. in Hedwig. 1898, p. 127.

2. — Pac. : Austral. or. (Nov. Vales.).

- acuminata* } Brid. — *Tortula agraria*.  
                   } Hedw. — *B. fallax*.  
*acuta* Brid. — *B. gracilis*.  
*acutata* C. M. in *Hedwig*. 1899, p. 109.  
     Ster. — *Afr.* : Transvaal.  
*aestiva* Schultz. — *Tortula muralis* var. *aestiva*.  
 \* *ticinensis* Kindb. — *Tortula muralis* var. *ticinensis*.  
*afro-caespitosa* C. M. — *Tortella*.  
*afro-unguiculata* C. M. in *Hedwig*. 1899, p. 105.  
     2. Ad *Mangiferam*. — *Afr.* : Zambes. med.  
*albicuspis* Mitt. — *Didymodon*.  
*alexandrina* Ltz. — *Trichostomum nitidum*.  
*aloides* Fürnr. et Br. — *Aloina*.  
*alpigena* Lindb. et Arn. — *Didymodon*.  
*alpina* Kindb. — *Tortula brevipes*.  
     var. *mutica* Mol. — *Tortula alpina* var. *inermis*.  
*altipes* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 254.  
     2. — *As.* : China centr. (prov. Schen-Si).  
*ambigua* Bryol. eur. — *Aloina*.  
*amblyacra* C. M. in *Ule Bryoth. brasil.* n. 216.  
     2. Ad aqueduct. — *Am. merid.* : Rio de Janeiro.  
*amblyophylla* Jaeg. — *Didymodon*.  
*amœna* C. M. in *Hedwig*. 1898, p. 128.  
     2. — *Pac.* : Austral. or. (Victoria).  
*amœna* Schum. — *B. unguiculata*.  
*amphidiacea* C. M. — *Tortula* ?  
*amplexa* Lesq. in *Trans. of the Amer. Phil. Soc. XIII*, p. 5; *Lesq. et Jam. Man.* p. 118 (1884).  
     *Tortula humilis* Drumm. *M. bor. am. I ed. n.* 137.  
     2. Ad rup. madid. — *Am. sept.* : Canada, Oregon, Calif.  
*amplexifolia* (Mitt.) Jaeg. *Ad. I*, p. 272 (1871-72).  
     *Tortula amplexifolia* Mitt. *M. Ind. or. p.* 29 (1859).  
     2. — *As.* : N. W. Himalaya.  
*anastomosans* C. M. in *Linn.* 1878-79, p. 337.  
     2. — *Am. merid.* : Argentin. subtropic.  
*Anderssonii* (Aongstr.) Card. — *Tortula fuegiana*.  
*androgyna* C. M. — *Tortula* ?  
*angustata* } Braithw. — *Tortula subulata* var. *angustata*.  
                   } Mitt. (Bryol. jav.). — *Trichostomum*.  
*angustifolia* (Hook. et Grev.) C. M. *Syn. I*, p. 603.  
     *B. tenuirostris* Brid. *Bryol. univ. I*, p. 826 (1826).  
     *Tortula angustifolia* H. et G. in *Brewster Edinb. Journ. I*, p. 299 (1824);  
     Mitt. *op. cit.* p. 28.  
     2. — *As.* : Ceylan, Java.  
*angustifolia* Brid. — *Tortella inclinata*.  
*anoëctangiacea* C. M. in *Hedwig*. 1899, p. 105.  
     Ster. — *Afr.* : Prom. R. Sp.



*anomala* Mitt. — Timmiella.

*anostomosans* C. M. in *Linn.* 1878-79, p. 337.

2. — *Am. merid.* : Argentin. subtropic.

*anthropophila* C. M. — Aloina.

*apiculata* Hpe. in *Linn.* 1861-62, p. 519.

*Tortula apiculata* Mitt. *M. austr. am.* p. 153 (1869).

2. — *Am. merid.* : And. Nov. Granat.

*apiculata* Hedw. — *B. unguiculata* var. *apiculata*.

*spoclada* Par. *Suppl. Ind.* p. 20 (1900).

*B. subrevoluta* C. M. in *Linn.* 1878-79, p. 134.

1. — *Am. merid.* : Argentin. Cordob.

*appressifolia* (Mitt.) Jaeg. *op. cit.* p. 281.

*Tortula appressifolia* Mitt. *op. cit.* p. 159.

2. — *Am. merid.* : And. Peruv.

*aprica* C. M. — *Tortella*.

*arcuata* } C. M. — *Tortula*.

Griff. — *B. comosa*.

*arenacea* Jaeg. — *Tortula obtusifolia*.

*arenaria* Griff. — *Crossidium squamigerum*.

*argentinica* Par. — *Didymodon*.

*aristata* Br. et Sch. in *W. Sch. M. Abyss.*

— *Afr.* : M. Abyss.

*aristata* Brid. — *B. unguiculata*.

*aristatula* C. M. in *Jaeg. Ad. II*, p. 674 (1877-78).

*B. calycina* ? (*fid. Broth.*).

*Encalypta aristata* Hpe. in *Linn.* 1871-73, p. 513.

*Pyramitrium aristatum* Mitt. in *Trans. a. Proceed. of the roy. Soc. of Victoria*, 1882, p.

— *Pac.* : Austral. or. (Nov. Vales.).

*artocarpa* Lesq. in *Trans. of the Amer. Phil. Soc. XIII*, p. 4; *Lesq. et Jam. Man.* p. 126 (1884).

2. Terr. — *Am. sept.* : Oregon, Calif.

*asperifolia* Mitt. *M. Ind. or.* p. 34 (1859).

Ster. — *As.* : M. Alatau, Sikkim.

*asperifolia* C. M. — *Tortula*.

*astoriensis* Broth. in *Roell M. Am. bor. exs.*

— *Am. sept.* : Washington.

*atlantica* { Resch. — *B. graminicolor*.

Brid. — *Eucladium verticillatum*.

W. P. Sch. — *B. orizabensis*.

*atro-virens* (Sm.) W. P. Sch. var. *robusta* W. P. Sch. — *B. revolvens*.

*Australasiae* (Hook. et Grev.) Brid. *Bryol. univ. I*, p. 828 (1826); C. M. *Syn. I*, p. 619 (1849).

*B. rufseta* Tayl. in *Lond. Journ. of bot.* 1846, p. 51.

*Tortula Australasiae* H. et Gr. in *Brewster Edinb. Journ. I*, p. 301, t. 12 (1824); *Fl. of the N. Zeal. II*, p. 70 (1855).

- T. fuscescens* Fl. of the N. Zeal. II, p. 73, t. 85, f. I.  
*Trichostomum Australasiae* Jaeg. op. cit. p. 245.
2. — *Pac.* : Fret. reg. Georges, Austral., Nov.  
 Seeland. sept., Tasman.
- australis* Par. — *Tortula fuegiana*.
- austro-revoluta* Besch. in Bull. Torr. bot. Club, 1896, p. 80.  
 2. — *Am. merid.* : And. Boliv.
- basoutensis* Rehm. — *Hyophila*.  
*Beccarii* (Vent.) Jaeg. — *Tortella*.  
*Beecheyi* Lesq. — *B. vinealis* var. *cylindrica*.
- Bescherellei* Sb. in Jaeg. Ad. II, p. 673 (1877-78).  
*B. rigidula* Besch. Prodr. bryol. mexic. p. 36 (1871).  
 2. — *Am. sept.* : Mexic.
- bicolor* (Bryol. eur.) Lindb. in Oefv. af K. Vet.-Akad. Foerh. XX, p. 386 (1863).  
*Gymnostomum bicolor* Bryol. eur. fasc. 33-36 (Vol. I), Mon. p. 4, t. 29 (1846).  
*Pottia bicolor* C. M. Syn. I, p. 563 (1849).  
*Tortula bicolor* Lindb. de Tortul. p. 247 (1864).  
 2. Ad terr. humid. — *Eur.* : Salisb., Stir., Bavar., Tirol., Rhaet., Helvet. *Pac.* : Tasman.?
- Blyttii* W. P. Sch. — *Amphoridium Mougeotii*.
- bokkepoldiana* Rehm. M. Afr. austr. n. 108.  
 — *Afr.* : Prom. B. Sp.
- boliviana* Par. Ind. p. 63 (1894).  
*B. glaucescens* W. P. Sch. in Mand. Pl. Poliv. n. 1622.  
*B. subglaucescens* C. M. in Nuov. Giorn. bot. ital. 1897, p. 52.  
 — *Am. merid.* : And. Boliv.
- Bourgaeana* Besch. Prodr. bryol. mexic. p. 35 (1871).  
 2. — *Am. sept.* : Mexic.
- brachypelma* (C. M.) Par. — *Ptychomitrium sinense*.
- brachyphylla* Sull. in Bot. Pac. R. R. Survey (Whipple), p. 186, t. 2 (1856); Lesq. et Jam. Man. p. 123 (1884).  
 2. — *Am. sept.* : Calif.
- « *var. angustifolia* C. M. et Kindb. in Mac. Cat. pp. 55 et 264 (1892).  
*B. lateritia* Kindb. Eur. a. N. Am. Bryin. p. 264 (1897).  
 Ster. — *Am. merid.* : Columb. brit.
- brachyphylla* } Hpe. — *B. Hampeana*.  
 } Sull. et Lesq. — *B. purpurea*.  
*brachypila* (C. M.) Par. — *Tortula alpina*.  
*brachypus* Brid. — *B. unguiculata* var. *brachypus*.  
*Brebissoni* Brid. — *Dialytrichia*.  
 var. *fragilis* Card. — *Dialytrichia Brebissoni* var. *fragilis*.  
*brevicaulis* } Kindb. — *Desmatodon latifolius* var. *brevicaulis*.  
 } Schwaegr. — *B. fallax* var. *brevicaulis*.  
*brevifolia* } Brid. — *B. fallax* var. *brevifolia*.  
 } Lindb. — *Didymodon tophaceus*.

*brevisetata* Mont. in *Ann. sc. nat.* 1845, p. 107; *C. M. Syn. I*, p. 644 (1849).

*Tortula brevisetata* Mont. *Syll.* p. 39 (1856); *Mitt. M. austr. am.* p. 172 (1869).

2. Terr. — *Am. merid.* : Chile.

*brevisetata* Hpe. et C. M. — *Tortula*,

*brevisetacea* Jaeg. — *Tortula brevisetata*.

*Brotheri* (Lindb.) Par. — *Tortella*.

*Bruchiana* Hpe. — *Tortula canescens*.

*brunneola* C. M. in *Bull. hb. Boissier*, 1897, p. 113.

2. — *Am. sept.* : Guatemala.

*caespitosa* } Er. — *Tortula marginata*.

(Schwaegr.) Schultz. — *Tortella*.

var. *abbreviata* W. P. Sch.

var. *angustifolia* Rehm.

var. *longisetata* (Hpe.) Par.

*calceolifolia* Spr. — *Aloina*.

*calicicola* (Hpe.) Broth. *Natür. Pflanzenfam.* p. 408.

*Tortula calicicola* Mitt. in *Trans. and Proceed. of the roy. Soc. of Victoria*, 1882, p. 60.

*Trichostomum calicolum* Hpe.  *Ic. M. t.* 29 (1844); *C. M. Syn. I*, p. 587 (1849); *Par. Ind.* p. 1319 (1898).

2. Rupic. — *Pac.* : Austral. or. (Victoria).

*calycina* Schwaegr. *Suppl. II, P. I*, p. 63, t. 119 (1823); *Brid. Bryol. univ. I*, pp. 541 et 828 (1826); *C. M. Syn. I*, p. 599 (1849).

*Tortula calycina* Hook. et Grev. in *Brewster Edinb. Journ. I*, p. 291 (1824).

*T. flexuosa* var. *Hook. M. ex. ?*

2. Cortic. et terr. — *As.* : Ind. or. (fid. Hook. et Wils. *Fl. Tasman. II*, p. 174). *Am. merid.* : Chile. *Pac.* : Fret. reg. Georges, Austral., Nov. Seeland, King's Isl., Tasman.

• var. *brevicaulis* Hpe. in *Linn.* 1856, p. 204.

— *Pac.* : Austral. or.

*calyculosa* (Mitt.) Jaeg. *Ad. I*, p. 1, p. 266 (1871-72).

*Tortula calyculosa* Mitt. *M. austr. am.* p. 156 (1869).

2. — *Am. merid.* : And. Quitens.

*calymperifolia* C. M. — *Tortula javanica*.

*campylocarpa* Tayl. — *Didymodon*.

*cancellata* C. M. — *Dialytrichia*.

*capillipes* Broth. in *Hedwig.* 1895, p. 125.

2. Terr. — *Am. merid.* : Brasil. or. (Goyaz).

*carnifolia* C. M. et Kindb. — *Tortula Bolanderi*.

*catillum* C. M. — *Aloina*.

*cavifolia* W. P. Sch. — *Pterygoneuron lamellatum*.

*chimboraensis* (Mitt.) Jaeg. — *Didymodon*.

*chloronotos* } Brid. — *Crossidium squamigerum*.

Bruch. — *Crossidium*.

f. *nivea* Besch. — *Crossidium chloronotos f. nivea*.

*chloropus* Kindb. — *Ceratodon*.

- chlorotricha* (*Broth. et Geh.*) *Par. Ind.* p. 67 (1894).  
*Tortula chlorotricha* *Br. et G. in Oefv. af K. Sv. Vet.-Akad. Foer* 1893.  
 2. — *Pac.* : Austral. or. (Nov. Vales., Victoria).
- chrysochacte* *C. M. in Hedwig.* 1898, p. 126.  
 2. *Terr.* — *Pac.* : Austral. or. (Victoria).
- chrysopila* *C. M.* — *Tortula*?
- chrysopoda* *C. M. et Kindb. in Mac. Cat.* p. 57 (1892).  
 2. *Terr.* — *Am. sept.* : Columb. brit., ins. Vancouver.
- chrysopus* *C. M. in Hedwig.* 1898, p. 127.  
 2. — *Pac.* : Tasman.
- circinalis* *C. M.* — *Tortula atro-virens.*
- circinacula* *C. M. et Kindb. in op. cit.* p. 56.  
 2. *Terr.* — *Am. sept.* : Columb. brit., ins. Vancouver.
- cirrata* { *Brid. p.p.* — *Dicranoweisia.*  
 { *Brid., Bryol. eur.* — *Tortella caespitosa.*
- cirrifolia* *W. P. Sch.* — *Trichostomum hibernicum.*
- Closteri Aust. in Coult. Bot. Gaz. I, p. 29 et IV, p. 30; Lesq. et Jam.*  
*Man. p. 127 (1884).*  
*Ster.* — *Am. sept.* : New Jersey, Florid.
- commutata* *Jur.* — *B. convoluta var. commutata.*
- comosa* *Dz. et Mk. M. frond. ined. Archip. ind. p. 46, t. 17 (1854);*  
*Bryol. jav. I, p. 103 (1855-61).*  
*B. arcuata Griff. Not. p. 411 et Ic. pl. asiat. II, t. 79, f. I (1849).*  
*Trichostomum tortile Wils. in Kew Journ. bot. IX, p. 321 (1857).*  
 2. *Terr.* — *As.* : Nepal., Sikkim, Khasia, Java, Singapore, Amboine.
- concava* *W. P. Sch.* — *Pterygoneuron lamellatum.*
- confertifolia* *Mitt.* — *Didymodon*?
- conica* { *Brid.* — *Tortula agraria.*  
 { *Kindb.* — *Ceratodon.*  
 { *Spreng.* — *Ceratodon stenocarpus.*
- consanguinea* (*Thw. et Mitt.*) *Sb. in Jaeg. Ad. II, p. 673 (1877-78).*  
*Tortula consanguinea T. et M. in Journ. of the Linn. Soc.* 1872, p. 300.  
 2. *Terr.* — *As.* : Ceylan.
- constricta* *Mitt. M. Ind. or. p. 33 (1859).*  
*Tortula vinealis var. Wils. in Kew Journ. bot. IX, p. 322 (1857).*  
 2. — *As.* : N. W. Himalaya, Nepal., Sikkim.
- convoluta* *Hedw. Descr. I, p. 86, t. 32 (1787); Brid. Bryol. univ. I,*  
*p. 560 (1828); Bryol. eur. fasc. 13-15 (Vol. II), Mon. p. 30,*  
*t. 154 (1842); C. M. Syn. I, p. 614 (1849); W. P. Sch. Syn.*  
*II ed. p. 214 (1876).*  
*B. deusta Brid. Mant. M. p. 96 (1819).*  
*B. purpurea Agh. mss.; Schultz Rec. p. 22 (1823).*  
*B. setacea Hedw. Fund. II, p. 92 (1782).*  
*Bryum convolutum Dicks. Pl. crypt. fasc. II, p. 6 (1790).*  
*Br. setaceum Huds. Fl. angl. p. 409 (1762).*  
*Br. stellata nitidum, pallidum, capsulis tenuissimis Dill. Hist. M. p. 381,*  
*t. 48, f. 41 fide C. M. (1741).*  
*Hypnum setaceum Weis Fl. Gott. p. 206 (1770).*

- Mnium setaceum* Poll. *Pl. palat.* III, p. 54 (1777); *Ehrh. in Hannov. Mag.* 1780.  
*Streblotrichum convolutum* P. B. *Prodr.* pp. 27 et 89 (1805).  
*Tortula convoluta* Schrad. *Spic. fl. germ.* p. 66 (1794).  
*T. setosa* P. B. *op. cit.* p. 93.
2. Ad terr. nud. vel herbos. — *Eur.* : Tota. As. : Japon., Jenissei med., Caucas. *Afr.* : Alger, Tunet., ins. Azor. *Am. sept.* : Canada, Pensilv., Wisconsin, M. Rupestr., Columb. brit., ins. Vancouver, Kansas, Calif., Alabama, etc.
- « *var. commutata* (Jur.) Husn. *Muscol. gall.* p. 108 (1886).  
*B. commutata* Jur. in *Verh. d. K. K. zool. bot. Ges. in Wien*, 1874, p. 377 et *Laubmfl. v. Oesterr. Ung.* p. 119 (1882); *Par. Ind.* p. 67 (1894).  
*B. convoluta* Jur. et Ung. in Ky. « *die Insel Cypern.* »  
*B. convoluta var. densa* Milde *Bryol. siles.* p. 116 p.p. (1869).  
*B. convoluta var. fragilifolia* Spr.  
*B. convoluta var. sardoa* C. M. *Syn. I*, p. 615 (1849) ?  
*Tortula commutata* Vent. et Bott. *Enum. critt.* p. 49 (1884).  
*Trichostomum undatum* W. P. Sch. *op. cit.* p. 180.
- Ad. mur. et rup. calcar. — *Eur.* : Belgia, Arduenn., Westphal., Sardin., ins. Zante, Pyren. occid., Catalaun., ins. Balear. As. : Ins. Cypr.
- var. densa* Milde p.p. } — *B. convoluta var. commutata.*  
 — *B. convoluta var. uliginosa.*
- « *var. filiformis* Hagen in Tromsø Mus. *Aarsheft.* 1898-99, p. 51.  
 — *Eur.* : Norveg. bor.
- var. fragilifolia* Spr. — *B. convoluta var. commutata.*
- « *var. obtusata* C. M. et Kindb. in *Mac. Cat.* p. 265 (1892).  
*Rupic.* — *Am. sept.* : Ontario.
- var. sardoa* C. M. — *B. convoluta var. commutata* ?
- « *var. Stockumi* Warnst. in *Allgem. bot. Zeitschr.* 1899, *Beiheft.*  
 — *Eur.* : Vall. Rhen. med.
- « *var. uliginosa* Limpr. *Kryptfl. v. Schles. I*, p. 172 (1876) et *Laubm. Deutschl. II*, p. 630 (1888).  
*B. convoluta var. densa* Mitt. *op. et l. cc.* p.p.  
 Ad tumul. terr. — *Eur.* : Boruss. or., Siles.
- convoluta* Jur. et Ung. — *B. convoluta var. commutata.*  
*cordata* Lindb. — *Didymodon.*  
*corticicola* R. C. — *Tortella* ?
- costa-ricensis* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1892, I, p. 154.  
 2. Ad. lapid. — *Am. sept.* : Costa-Rica.
- costata* (Mitt.) Jaeg. *Ad. I*, p. 281 (1871-72).  
*Tortula costata* Mitt. *M. austr. am.* p. 159 (1869).  
 2. — *Am. merid.* : And. Quitens.
- Craufordi* Par. — *Tortula asperifolia.*  
*crinita* Schultz. — *B. pilifera.*  
*crispa* Hedw. — *Dicranella Schreberi.*  
*crispata* Hpe. — *Calyptopogon.*  
*crispatula* C. M. — *Calyptopogon.*
- crispifolia* (Mitt.) Jaeg. *op. cit.* p. 276.  
*Tortula crispifolia* Mitt. in *Handb. of the N. Zeal. Fl.* p. 419 (1867).  
 2. — *Pac.* : Nov. Seeland. merid.

- crispula* Hpe. — *B. subulifolia*.  
*crocea* Brid. } — *B. paludosa*.  
     *var. paludosa* Roehl. }
- Crügeri* Sonder in *C. M. Syn. I*, p. 618 (1849) et *II*, p. 630 (1851); *Lesq. et Jam. Man.* p. 122 (1884).  
*Tortula Crügeri* Mitt. *M. austr. am.* p. 150 (1869).  
 2. Terr. — *Am. sept.*: Florid., Ludovic., Cuba, ins. S. Vincent, Barbades. *Am. merid.*: Venezuela, And. Quitens. et Peruv.
- cucullatifolia* C. M. — *Aloina*.  
*cucullatula* C. M. — *Tortula navicularis*.  
*cucullifera* (Mitt.) Jaeg. *op. cit.* p. 273.  
*Tortula cucullifera* Mitt. *op. cit.* p. 155.  
 2. — *Am. merid.*: And. Quitens. et Peruv.
- cuneifolia* } (Dicks.) Brid. — *Tortula*.  
                   } W. et M. — *Tortula muralis var. aestiva*.  
*curta* Hedw. — *Ditrichum tortile var. pusillum*.  
*curvipes* C. M. in *Linn.* 1878-79, p. 344.  
 2. — *Am. merid.*: And. Argentin. subtropic.
- curvirostris* Lindb. — *Hymenostylium*.  
     *var. commutata* Lindb. — *H. curvirostre var. commutatum*.  
     *var. laeviuscula* L. et A. — *H. curvirostre var. laeviusculum*.  
*cuspidata* } (H. f. et W.) Jaeg. — *Tortula*.  
                   } Schultz. — *B. unguiculata var. cuspidata*.
- cylindrangia* C. M. in *Hedwig.* 1898, p. 126.  
 2. — *Pac.*: Austral. or. (Victoria).
- cylindrica* } Mac. p.p. — *B. tortellifolia*.  
                   } (Tayl.) W. P. Sch. — *B. vinealis var. cylindrica*.  
                   } Wils. — *Dialytrichia Brebissoni*.  
     *var. sinuosa* Lindb. — *B. sinuosa*.  
*cylindrotheca* (Mitt.) Jaeg. — *Trichostomum*.  
*clanica* Lange. — *Tortula pulvinata*.
- decidua* C. M. in *Linn.* 1880-82, p. 427.  
     *B. uruguensis* Par. *Ind.* p. 101 (1894).  
 2. — *Am. merid.*: Uruguay.
- decidua* (Mitt.) Jaeg. — *Tortula*.
- declivium* C. M. in *Linn.* 1880-82, p. 425.  
     Fl. et fr. ign. Arenic. — *Am. merid.*: Argentin. Buenos-Ayr.
- decursivula* Kindb. in *Mac. Cat.* p. 264 (1892).  
     Ster. Rupic. et terr. — *Am. sept.*: M. Rupestr.
- defossa* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 236.  
 2. — *As.*: China centr. (prov. Schen-Si).
- depressa* Sull. — *B. flagellaris*.  
*deusta* Brid. } (Mant. M.) — *B. convoluta var. commutata*.  
                   } (Bryol. univ.) — *Didymodon luridus*.  
*Dicksoniana* (1) Schultz. — *Tortula cuneifolia*.

(1) *B. diaphana* Brid. *Bryol. univ.* I, p. 577. — « An *Grimmia*? » C. M. *Syn. I*, p. 645. — Deleatur.

- Dieckei* Broth. — Didymodon.  
*dimorpha* C. M. — Didymodon.
- dioritica* C. M. in *Linn.* 1875, p. 402.  
 Ster. Rupic. — *As.* : M. orient. mar. Erythraei.
- dissita* C. M. in *Hedwig.* 1898, p. 1:9.  
 Ster. — *Pac.* : Nov. Caledon.
- domestica* Brid. — Tortula agraria.  
*Dorrii* R. C. — Hyophila.  
*Drummondii* (Mitt.) Jaeg. — Tortella fragilis.  
 var. *setacea* R. Farn. — Tortella fragilis var. *setacea*.  
*dubia* Brid. — B. unguiculata.  
*duriuscula* (Wils.) Jaeg. — Trichostomum.
- Duseni* C. M. in *Dusen M. Camer.* n. 120; *Broth. in Engler's Bot. Jahrb.* 1897, p. 237.  
 2. Rupic. — *Afr.* : Cameroon.
- elata* D. R. et Mont. *Fl. Alger. ined.*; C. M. *Syn.* I, p. 620 (1849).  
 Tortula *elata* Mont. *Syll.* p. 41 (1856).  
 2. Rupic. — *Afr.* : Alger. *Am. sept.* : Montana, Washington, Oregon, Calif.
- Elliottii* Broth. in *Engler's Bot. Jahrb.* 1897, p. 257.  
 2. fr. ign. — *Afr.* : Reg. lacustr.
- elliptithecica* C. M. in *Nuov. Giorn. Bot. ital.* 1897, p. 238.  
 2. — *As.* : China centr. (prov. Schen-Si).
- Endressii* Garov. { — B. flavipes (*fid. Jur.*)  
 { — Tortula aciphylla ? (*fid. W. P. Sch.*)  
*ericetorum* Brid. — B. unguiculata var. *patagonica*.
- erosa* Hpe. *miss.*; C. M. in *Bot. Zeit.* 1862, p. 348.  
 2. — *Am. merid.* : Venezuela.
- eroso-denticulata* C. M. — B. trachyphylla.  
*erythrodonta* Tayl. — Streptopogon.
- erythropoda* W. P. Sch. in *Besch. Prodr. bryol. mexic.* p. 36 (1871).  
 2. — *Am. sept.* : Mexic.
- erythrotricha* C. M. — Tortula alpina.
- eubryum* C. M. in *Fl.* 1879, n. 24.  
 Ster. — *Afr.* : M. Kenia.
- eucalyx* W. P. Sch. in *Breutel M. capens.*; *Jaeg. Ad. I*, p. 305 (1871-72).  
 — *Afr.* : Prom. B. Sp.
- eustegia* Card. et Ther. in *Bot. Gaz.* 1900, p. 17, t. IV.  
 2 ? Terr. — *Am. sept.* : Idaho.
- eutrichostomum* C. M. — Tortella.
- excavata* (Mitt.) Jaeg. *op. cit.* p. 275.  
 Tortula *excavata* Mitt. *M. austr. am.* p. 151 (1869).  
 2. — *Am. merid.* : And. Quitens.
- excurrents* Broth. — Didymodon  
*exesa* C. M. — Tortula ?
- falcifolia* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 237.  
 2. — *As.* : China centr. (prov. Schen-Si).

*fallax* Hedw. *Descr. I*, p. 62, t. 24 (1787); *Brid. Bryol. univ. I*, p. 556 (1826); *Bryol. eur. fasc.* 13-15 (*Vol. II*), *Mon. p.* 23, t. 147 (1842); *C. M. Syn. I*, p. 616 (1849); *W. P. Sch. Syn. II ed.* p. 205 (1876).

*B. acuminata* Hedw. *Sp. M.* p. 117, t. 25, f. 5-7 (1801); *Schwaeagr. Suppl. I, P. I*, p. 123 (1811).

*B. fallax* ♂ *flavescens* *Brid. Bryol. univ. l. c.*

*B. flavescens* *ej. Mant. M.* p. 92 (1819).

*B. imberbis* *Brockm. Laubm. Mecklenb.* p. 77 (1869).

*B. nervosa* *Schultz Fl. Starg.* p. 305 *excl. syn. Brid.* (1806).

*Bryum fallax* *Dicks. Pl. crypt. fasc. III.* p. 5 (1793).

*Br. flavescens* *ej. op. cit. fasc. II*, p. 4, t. 4, f. 5 (1790).

*Br. imberbe* *Huds. Fl. angl. I ed.* p. 409 (1762).

*Br. perangustis foliis et cauliculis, foliis rarioribus cinctis, capitulis erectis et surculis annotinis egredientibus* *Dill. Cat. Giss.* p. 225 (1718).

*Br. tenue barbatum, foliis angustioribus et rarioribus* *ej. op. cit.* p. 385, t. 48, f. 49 *fid. Limpr.*

*Br. tenue, imberbe et pallidum, foliis crebrioribus* *Dill. Hist. M.* p. 382, t. 48, f. 46 (1741), *fid. C. M.*

*Dicranum pellucidum* *Drumm. M. bor. am. n.* 108 *fid. Kindb.*

*Mollia fallax* *Schrank Bayer. Fl. II*, p. 458 (1789).

*Tortula fallax* *Schrad. Samml. Krypt. Gew. I. n.* 53 (1796).

*T. flavescens* *Brid. Sp. M. I*, p. 263 (1806).

*T. imberbis* *Sm. Fl. brit. III*, p. 1261 (1804); *Lindb. de Tortul.* p. 250 (1864).

*T. Stokesii* *Turn. Musc. hib.* p. 48 (1804).

2. Ad mur. et terr. humidiusc. fere ubique. — *Eur.* : In reg. campest. vulgariss. *As.* : *Jenissei med. et infer.*, *Caucas. Afr.* : *Alger.*, *Tunet.*, *Madeir.* *Am. sept.* : *Groenland.*, *Canada tot.*, *M. Rupestr.*, *Columb. brit.*, *ins. Vancouver, reg. sept.*, *occid. et orient.*

“ *var. brevicaulis* (*Schleich.*) *Bryol. eur., C. M. et W. P. Sch. II. cc.*

*B. brevicaulis* *Schwaeagr. Suppl. I, P. I*, p. 126, t. 32 (1811); *Brid. Bryol. univ. I*, p. 551 (1826).

*Tortula brevicaulis* *Schleich. Cat.* 1807.

*T. brevicaulis* γ *Hook. et Grev. in Brewster Edinb. Journ. I*, p. 300 (1824).

Ut typ. — *Eur. et Am. sept.* : Cum typ.; *Vermont.*

“ *var. brevifolia* *Schultz Rec.* p. 20, t. 33, f. 22 B (1823); *Bryol. eur., C. M. et W. P. Sch. II. cc.*

*B. brevifolia* *Brid. Bryol. univ. I*, p. 555 (1826).

*Bryum brevifolium* *Dicks. Pl. crypt. fasc. II*, p. 4 (1790).

*Br. palustre, brevifolium, capsulis nigricantibus* *Dill. Hist. M.* p. 377, t. 47, f. 39 (1741).

*Tortula brevifolia* *Sm. Fl. brit. III*, p. 1259 (1804).

Ut typ. — *Eur.* : *Septentr.*; *Sardin.*, *Am. sept.* : *Passim.*

*var. flavescens* *Brid.* — *B. fallax.*

“ *var. longifolia* *Fleisch. et Warnst. Bryoth. Eur. merid. n.* 36.

Ad rup. calcar. — *Eur.* : *Sicilia* (*Panorm.*)

*var. recurvifolia* *Husn.* — { *B. reflexa.*

*var. reflexa* *Brid.* —

*var. Turneri* *Brid.* — *Didymodon rubellus.*

*var. vinealis* *Hüb.* — *B. vinealis.*

*fallax* *Sull.* — *B. subfallax.*

*fastigiata* *Schultz.* — *B. unguiculata var. fastigiata.*

*Fendleri* *C. M. in Linn.* 1878-79, p. 483.

2. — *Am. merid.* *Venezuela.*



*ferrinervis* C. M. in *Bull. hb. Boissier*, 1897 p. 557.

2. — *Am. sept.* : Jamaic.

« var. *Eggersiana* C. M. in *Hedwig*, 1898, p. 232.

Cortic. — *Am. sept.* : S. Doming.

*ferrinervis* C. M. (China). — B. *ferrugineo-nervis*.

*ferruginea* W. P. Sch. in *Besch. Prodr. bryol. mexic.* p. 37.

2. — *Am. sept.* : Mexic.

*ferrugineo-nervis* *Broth. op. et l. cc.*

B. *ferrinervis* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 255.

B. *serrinervis* (errat. !) *Par. Suppl. Ind.* p. 31 (1900).

Ster. — *As.* : China centr. (prov. Schen-Si).

Fiori Vent. in *Rev. bryol.* 1885, p. 66.

2. Ad coll. arenac. — *Eur.* : Ital. super. (Modena).

*flaccidiseta* Ltz. *Moosst.* p. 161 (1864).

2. — *Am. sept.* : Mexic.

*flagellaris* W. P. Sch. in *Ann. sc. nat. II ser., VI*, p. 146, t. 10; C. M. *Syn. I*, p. 613 (1849).

B. *depressa* Sull. in *U. St. Exped. (Wilkes)* 1859, p. 5.

*Tortula flagellaris* Mitt. *M. austr. am.* p. 150 (1869).

2. — *Am. merid.* : Chile, fret. Magellan.

*flavescens* (Hook. et Grev.) Brid. *Bryol. univ. I*, p. 613 (1826); C. M. *Syn. I*, p. 613 (1849).

B. *fuscescens* Wall. *Cat. n.* 7567.

*Tortula flavescens* H. et Gr. in *Preioster Edinb. Journ. I*, p. 297, t. 12 (1824); Mitt. *M. Ind. or.* p. 29 (1859).

2. Ad terr. argill. — *As.* : N. W. Himalaya, Nepal.

*flavescens* Brid. — B. *fallax*.

*flavicaulis* C. M. in *Hedwig*, 1897, p. 258.

2. — *As.* : China centr. (prov. Schen-Si).

*flavipes* *Bryol. eur. fasc. 13-15 (Vol. II)*, *Mon.* p. 30, t. 155 (1842); C. M. *Syn. I*, p. 621 (1826); W. P. Sch. *Syn. II*, ed. p. 214 (1846).

B. *Enderessii* Garov. *Bryol. austr.* p. 37 (1840).

B. *lutescens* Laurer in sched.

*Tortula flavipes* Wils. *Bryol. brit.* p. 128 (1855).

2. Ad terr. humos. glareos. — *Eur.* : Austr. infer., Salisb., Stir., Carinth., alp. Bavar., Francon., Tirol., Voralberg, Helvet., alp. Comens.; deest in Britann. et in Scandin.

*flavipila* W. P. Sch. — B. *pilifera*.

*flavo-virens* Lindb. — *Trichostomum*.

*flexicaulis* C. M. — *Grimmia suspecta*.

*flexifolia* Hpe. in *Linn.* 1859-60, p. 496.

2. Terr. — *Am. sept.* : Oregon, S. Nevada Calif.

*flexifolia* } Brid. — ? (1)  
                  } Mac. — B. *tortellifolia*.

(1) *flexifolia* Brid. *Bryol. univ. I*, p. 560.

B. *flexuosa* ej. *Mant. M.* p. 93.

B. *scoparia* Schultz in *Nov. Act. Phys.*

medic. Acad. Caesar. XII, II, p. 718.

*Tortula flexuosa* Brid. *Sp. M. I*, p. 256.

« An *Macromitrium* ? » C. M.  
*Syn. I*, p. 645.

*flexiset* Kindb. — Timmiella.

*flexuosa* (Hook.) Schultz Rec. p. 16, t. 33 (1823); Brid. Bryol. univ. I, p. 573 (1826); C. M. Syn. I, p. 609 (1849).

B. Hookeri Steud. Nom. crypt. p. 72.

*Tortula flexuosa* Hook. M. ex. t. 125 (1818-20).

2. — Afr. : Prom. B. Sp.

*flexuosu* Brid. — Cf. *B. flexifolia* ej.

*fragilis* } Tayl. — *Tortula*.  
              } Wils. — *Tortella fragilis*.

var. *furcata* Hpe. — *Tortula fragilis* var. *furcata*.

var. *setacea* Rob. Farn. — *Tortella Drummondii* var. *setacea*.

*frigida* C. M. in Bot. Zeit. 1858, p. 163.

2. In turf. — Am. merid. : Summ. And. Peruv.

*Fristedti* C. M. — *Tortella*.

*fulvo-fusca* C. M. in Hedwig. 1898, p. 128.

Ster. — Pac. : Nov. Caledon.

*Funckiana* Schultz. — *B. paludosa* var. *Funckiana*.

*fusca* C. M. Syn. I, p. 610 (1849).

*Tortula fusca* Mitt. M. austr. am. p. 160 (1869).

2. — Am. merid. : Chile.

*fusca* W. P. Sch. — *B. Mandoni*.

*fuscescens* Wall. — *B. flavescens*.

*fuscinervia* (Mitt.) Jaeg. Ad. I, p. 275 (1871-72).

*Tortula fuscinervia* Mitt. op. cit. p. 154.

2. — Am. merid. : Ins. Chiloë.

*fuscula* C. M. in Linn. 1878-79, p. 343.

Fl. et fr. ign. — Am. merid. : Argentin. Cordob. et subtropic.

*galeata* C. M. — *Aloina*.

*gangetica* C. M. in Linn. 1871-73, p. 177.

2. — As. : Bengal.

*Geheebii* Broth. — *Crossidium*.

*geheebiopsis* C. M. — *Geheebia* Mülleri.

*geniculata* Mont. in Ann. sc. nat. 1845, p. 107; C. M. Syn. I, p. 629 (1849).

*Tortula geniculata* Mont. Syll. p. 39 (1856); Mitt. op. cit. p. 161.

2. — Am. merid. : Chile.

*Germainii* C. M. — *Tortella*.

*gigantea* Funck. — *Geheebia gigantea*.

*glabriuscula* C. M. — *Didymodon rubellus*.

*glacialis* { Br. et Sch. — *Tortula subspathulata* ?  
              { Kindb. (ut subsp.) — *Desmatodon latifolius* var. *glacialis*.  
              { Kze. — *Tortula*.  
              { Besch. — *Tortula amphidiacea* ?

*glauescens* Hpe. in Linn. 1863, p. 128.

*Tortula glauescens* Mitt. op. cit. p. 152.

2. Rupic. — Am. merid. : And. Nov. Granat.

*glaucescens* W. P. Sch. — *B. boliviana*.  
*Glaziovii* Sb. — *B. lurida*.

*Godmanniana* C. M. in *Bull. hb. Boissier*, 1897, p. 193.  
*Tortula campylocarpa* Mitt. *M. austr. am.* p. 159 p.p. (1869).  
 2. — *Am. sept.* : Guatemala.

*goniospora* C. M. in *Rev. bryol.* 1887, p. 57 (nom.) et in *Hedwig.* 1898, p. 131.  
 2. — *Pac.* : Nov. Caledon.

*gracilentia* Mitt. *M. Ind. or.* p. 35 (1859).  
*Tortula flavescens* var. *Wils.* in *Kew Journ. bot.* IX, p. 322 (1857).  
 2. — *As.* : W. Himalaya.

*gracilentia* Hpe. — *B. lurida*.

*gracilescens* W. P. Sch. in *Besch. Prodr. bryol. mexic.* p. 34 (1871).  
 2. — *Am. sept.* : Mexic.

*graciliformis* W. P. Sch. in *op. cit.* p. 35.  
 2. — *Am. sept.* : Mexic.

*gracilis* (Schleich.) *Schwaeagr. Suppl. I, P. I*, p. 125, t. 34 (1811); *Brid. Bryol. univ. I*, p. 53 (1826); *Bryol. eur. fasc.* 13-15 (*Vol. II*), *Mon.* p. 22, t. 45 (1842); *C. M. Syn. I*, p. 609 (1849); *W. P. Sch. op. cit.* p. 210.

*B. acuta* *Brid. Mant. M.* p. 96 (1819).  
*Tortula acuta* *ej. Sp. M. I*, p. 265 (1806).  
*T. gracilis* *Schleich. Cat.* 1807.

2. In glareos. calcar. et aren. argill. — *Eur.* : Media et praecip. merid. *As.* : Caucas. *Afr.* : Alger. Tunet., *Am. sept.* Canada, Vermont., *M. Rupestr.*, Calif.

• *var. irrigata* H. Müll. *Westfal. Laubm. n.* 404; *Limpr. Laubm. Deutschl. . . II*, p. 625 (1888).  
*Ster. Aquat.* — *Eur.* : Westphal.

• *var. multiseta* *Limpr. op. et l. cc. (forsan spec. propria !)*  
 — *Eur.* : Istria (Tergest.)

*var. propagulifera* Schiffn. — *Didymodon rigidulus f. propagulifera*.

• *var. rufescens* W. P. Sch. in *sched.*; *Limpr. op. et l. cc.*  
 In argill. — *Eur.* : Alsatia.

*var. squarrulosa* Kindb. — *B. subicmadophila*.

• *var. viridis* *Bryol. eur., C. M. et W. P. Sch. op. et l. cc.*  
*B. viridis* W. P. Sch. in *Rab. Bryoth. eur. n.* 818 (1865).  
*Tortula fallax var. stricta* de Not. *Spec. de Tortul. ital. n.* 26.  
*T. viridis fallax* H. et Grev. in *Brewster Edinb. Journ. I*, p. 300 (1824).  
 Ut typ. — *Eur.* : Belgia, Helvet., Istr., merid. *Afr.* : Alger.

*gracilis* Sull. — *B. subulifolia*.

*graminicolor* C. M. *Syn. I*, p. 611 (1849).  
*B. atlantica* W. P. Sch. in *sched. (fid. C. M.)*.  
*Tortula graminicolor* Mitt. *M. austr. am.* p. 160 (1869).  
 2. Terr. — *Am. merid.* : Chile.

*graminicolor* Resch. — *B. orizabensis*.

*grandiretis* *Broth. M. Turkestan. n.* 29 et 30.  
 Ad *Salic.* et terr. — *As.* : Turkestan.

*gregaria* (Mitt.) Jaeg. *Ad. I*, p. 272 (1871-72).

*Tortula apiculata* Wils. in *Kew Journ. bot. IX*, p. 322 (1857).

*T. gregaria* Mitt. *M. Ind. or.* p. 29 (1859).

*Trichostomum brachyphyllum* Wils. in *op. cit.* p. 321.

2. — *As.* : Tibet., W. Himalaya, Nepal. *or.*

*grimmiacea* C. M. in *Linn.* 1880-82, p. 427.

2. — *Am. merid.* : Argentin, Buenos-Ayr.

*grisea* (Jur.) Boul. — *Crossidium*.

*Hampeana* Par. *Ind.* p. 76 (1894).

*B. brachyphylla* Hpe. in *Linn.* 1859, p. 625.

2. — *Pac.* : Austral. *or.*

*hamulus* C. M. — *Aloina*.

*Harristii* Par. — *B. recurvicauspis*.

*hastata* Mitt. *M. Ind. or.* p. 34 (1859).

*Tortula comoa*? Wils. in *op. et l. cc.*

2. — *As.* : Sikkim.

*Haussknechtii* Jur. et Milde. — *Didymodon*.

*helvetica* Kindb. — *Ceratodon purpureus*.

*Henrici* Rau. — *Crossidium*.

*himantina* Besch. — *Tortella*.

*Hookeri* Stend. — *B. flexuosa*.

*Hornschuchiana* Schultz *Rec. in Nov. Act. Acad. Leop. XI*, p. 217,

t. 33, f. 25 (1823); *Bryol. eur. fasc.* 13-15 (*Vol. II*), *Mon.* p. 28, t. 148;

*C. M. Syn. I*, p. 608 (1849); *W. R. Sch. op. cit.* p. 211.

*B. revoluta* W. et M. *Bot. Tschb.* p. 210 (1807).

*B. revoluta*  $\beta$  *Hornschuchiana* Brid. *Bryol. univ. I*, p. 572 (1826).

*Tortula Hornschuchiana* de Not. *Syll.* p. 179 (1838).

*T. revoluta* H. et T. *Musc. brit. Fed.* p. 33, t. 12 (1818).

*T. revoluta*  $\beta$  *Hornschuchiana* de Not. in *Mem. Acad. Turin. XL*, p. 314 (1858).

2. In loc. glareos. argill. — *Eur.* : Med. et merid.; Norvegia, Dania.

*Afr.* : Alger., Choa. *Pac.* : Austral. *or.*

*horridifolia* C. M. et Kindb. in *Mac. Cat.* p. 57.

Ster. Ad rup. humid. — *Am. sept.* : Columb. brit., ins. Vancouver.

*humilis* Hedw. — *Tortella coespitosa*.

*hyalinoblata* (Broth.) Par. — *Tortella*.

*hyperborea* ) R. Br.? Brid. — *Tortula mucronifolia*.

Montb. — *Tortula magellanica*.

*icmadophila* *Bryol. eur. fasc.* 43 (*Vol. II*), *Mon. Suppl.* p. 1, t. 146 (1850);

*C. M. Syn. I*, p. 614 (1849) et *II*, p. 630 (1851); *W. P. Sch. op. cit.* p. 211.

*B. abbreviatifolia* H. Müll. in *Verh. d. bot. Ver. d. Prov. Brandenb.* 1866, p. 58.

*Tortula icmadophila* Lindb. de *Tortul.* p. 249 (1861)

2. Ad rup. cataract. — *Eur.* : Norveg. (Dovre), Westphal., Pinzgov., Salisb., Stir., Carinth., alp. Bavar., Tirol., Helvet., Arvernia. *As.* : Caucas.

*imberbis* Brockm. — *B. fallax*.

*imperfecta* (C. M.) Broth. *op. et l. cc.*

*Pottia imperfecta* Par. *Ind.* p. 1023 (1897).

*Trichostomum imperfectum* C. M. in *Linn.* 1878-79, p. 31.

2. — *Am. merid.* : Bolivia.

*inaequalifolia* Tayl. in *Lond. Journ. of bot.* 1846, p. 49, t. XVD; C. M. *Syn. I*, p. 616 (1849).

*Tortula inaequalifolia* Mitt. *M. austr. am.* 153 p. (1869).

2. — *Am. merid.* : And. Quitens.

*incerta* (Mitt.) Par. — *Tortula*.

{ Schum. — *B. unguiculata*.

*inclinans* W. P. Sch. in *Besch. Fl. bryol. Réunion, etc.* p. 55 (1890-81)

2. — *Afr.* : Ins. Franciae.

*inclinata* Schwaegr. — *Tortella*.

f. *acuminata* R. Farn. —

var. *brachypoda* Besch. —

var. *densa* l.tz. —

var. *leptotheca* Brid. —

var. *nitida* Mol. — *Trichostomum*.

*inclinatula* C. M. et Kindb. — *Tortella*.

*incrassata* Lindb. — *Didymodon*.

*incurvidens* (Z. Stirt.) Par. — *Tortula*.

*indica* Brid. — *Tortula orientalis*

*indigens* (Ren. et Card.) Par.

*B. unguiculata* (fide Kindb.).

*Trichostomum indigens* R. C. in *Bot. Gaz.* 1896, p. 50, t. IV, f. A; Par.

*Ind.* p. 1324 (1898).

2? ♂ ign. — *Am. sept.* : Terra nov.

*insidiosa* Jur. et M. — *Didymodon spadicens*.

*insulana* Husn. — *B. vinealis* var. *cylindrica*.

*intermedia* { de Not. — *Tortella caespitosa*.

W. P. Sch. — *Tortula montana*.

var. *calva* W. P. Sch. — *Tortula montana* var. *calva*.

var. *planifolia* Fl. et W. — *Tortula montana* var. *montana*.

var. *pulvinata* Husn. — *Tortula pulvinata*.

*inundata* (Mitt.) Jaeg. — *Didymodon*.

*Itzigohnii* Hpe. — *Tortula muralis*.

*Jamesoni* Tayl. — *Didymodon*.

*Jooriana* C. M. in *Bull. Torr. bot. Club.* V, p. 49 (1878) et in *Fl.* 1888, p. 77.

*B. Ravenelii* Aust. ?

2. Ad terr. argill. — *Am. sept.* : Carol. infer ?, Florid., Ludovic.

*jugicola* (Dz.) Par. — *Tortella* ?

*juniperoidea* C. M. in *Linn.* 1880-82, p. 429.

*Fl.* et fr. ign. — *Am. merid.* : Alp. Argentin. subtropic.

*khasiana* (Mitt.) Jaeg. — *Trichostomum*.

*Knighitii* (Mitt.) Jaeg. — *Tortella*.

*laevigata* (Mitt.) Jaeg. *Ad. I.* p. 274 (1871-72).

*Tortula laevigata* Mitt. *M. austr. am.* p. 160 (1869).

2. — *Am. merid.* : And. Quitens.

- laevipila* { (Brid.) Bryol. eur. — *Tortula*.  
 { Garov. — *Tortula papillosa*.  
 { Hpe. — *Tortula Preissiana*.  
 { Mac. — *Tortula pulvinata* var. *lato-excisa*.  
 { Schwaegr. — *Tortula laevipila* var. *laevipilaeformis*.  
 var. *laevipilaeformis* Husn. { — *Tortula laevipila* var. *laevipi*,  
 var. *marginata* Lindb. { laeformis.  
 var. *meridionalis* W. P. Sch. \
- lagunicola* C. M. in *Bull. hb. Boissier*, 1897, p. 104.  
 2. — *Am. sept.* : Guatemala.
- lamprocalyx* C. M. *Syn. I*, p. 599 (1849).  
 2. — *Am. merid.* : Chile.
- lanceolata* { Hedw. — *B. unguiculata* var. *obtusifolia*.  
 { Schum. — *B. unguiculata*.  
*lateritia* Kindb. — *B. brachyphylla* var. *angustifolia*.  
*latifolia* { Brid. — *Tortula agraria*.  
 { Br. (Bryol. eur.) — *Tortula*.  
 { Lindb. — *Desmatodon*.  
*lato-excisa* C. M. et Kindb. — *Tortula pulvinata* var. *lato-excisa*.  
*Laureri* Kindb. — *Desmatodon*.
- Laureriana* Ltz. *Moosst. p.* 161 (1864).  
 2. — *Afr.* : Prom. B. Sp.
- leptocarpa* Besch. *Prodr. bryol. mexic. p.* 35 (1871).  
 2. — *Am. sept.* : Mexic.
- leptotheca* W. P. Sch. — *Aloina*.  
*lepto-Tortella* C. M. — *Trichostomum*.  
*lepto-tortuosa* C. M. — *Trichostomum*.  
*leptotricha* C. M. et Kindb. — *B. pulvinata* subsp. *leptotricha*.
- leucocalyx* Mont. in *Ann. sc. nat.* 1838, p. 53; C. M. *Syn. I*,  
 p. 615 (1849).  
*Tortula leucocalyx* Mont. in *Gay, Hist. Chil., Crypt. VII*, p. 152 (1850).  
 et *Syll. p.* 38 (1856); *Mitt. M. austr. am. p.* 153 (1869).  
 2. Terr. — *Am. merid.* : Chile.
- leucodontoides* C. M. — *Geheebia*.  
*leucostoma* R. Br. — *Desmatodon suberectus*.
- ligularis* (Mitt.) Jaeg. *Ad. I*, p. 273 (1871-72).  
*Tortula ligularis* Mitt. *op. cit.* p. 156.  
 2. — *Am. merid.* : M. Pinchincha.
- limbata* { Lindb. — *Tortula sublimbata*.  
 { Mitt. — *Tortula*.  
*Lindigiana* Hpe. — *Gyroweisia*.  
*linearis* { (Sw.) Brid. — *Trichostomum*.  
 { Warnst. — *Trichostomum Warnstorffii*.  
*linoides* Brid. — *B. unguiculata*.
- lonchodonta* C. M. in *Linn.* 1878-79, p. 335.  
 2. Limic. — *Am. merid.* : Argentin. Cordob.
- lonchostega* C. M. in *Bull. hb. Boissier*, 1897, p. 195.  
 2. — *Am. sept.* : Guatemala.
- longifolia* Griff. — *Trichostomum*.

*Louisiadum* Broth. in *Oefv. af Finska Vet.-Akad. Foerh.* 1899, p. 113.  
2. fr. ign. Terr. — Pac. : Ins. Louisiad.

Luhmanni Broth. et Geh. in *op. cit.* 1895, p. 56.  
2. — Pac. : Austral. or. (Victoria).

lurida Hsch. *Fl. brasil.* I, p. 18 (1840); *C. M. Syn.* I, p. 608 (1849).

B. Glaziovii Sb. in *Jaeg. Ad. II*, p. 672 (1877-78).  
B. gracilentia Hpe. *Symb.* 1874, p. 484 et *Enum. M. Brasil.* p. 10 (1869).  
Hyophila lurida Jaeg. *op. cit.* I, p. 205 (1871-72).  
Weisia lurida Mitt. *M. austr. am.* p. 137 (1869).

2. — Am. merid. : Brasil. austr. or.

*lurida* Lindb. — Didymodon.

*lusitanica* Kindb. — Trimmiella Barbula.

*luteola* Mitt. — Tortula pungens.

*lutescens* Laur. — Barbula flavipes.

Mac Leana Rehm. *M. Afr. austr. n.* 482.  
— Afr. : Prom. B. Sp.

*macrocarpa* Brid. — Tortula subulata?

*macrogonia* Besch. — Tortula?

*macrorhyncha* Kindb. — Aloina.

*macrostegia* Kindb. — Rhamphidium dicranoides.

*madagassa* R. C. — Tortula?

*magellanica* } (Mont.) C. M. — Tortula.  
C. M. (1882). — Tortula fuegiana.

*magnifolia* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 255.  
Ster. — As. : China centr. (prov. Schen-Si).

*majuscula* C. M. in *op. cit.* 1898, p. 182.  
Ster. — As. : China centr. (prov. Schen-Si).

*mallacophylla* C. M. — Tortella.

Mandoni Par. *Ind. p.* 82 (1894).  
B. fusca W. P. Sch. in *Mandon Pl. Boliv. n.* 1624.  
— Am. merid. : And. Boliv.

*Manniae* C. M. — Crossidium.

*Mauchii* Rehm. — B. torquescens.

*melanocarpa* C. M. et Kindb. in *Mac. Cat. p.* 154 (1892).  
2. Rupic. — Am. sept. : Columb. brit.

*membranifolia* } Garov. — Crossidium griseum.  
} (Hook.) Sch. — Crossidium squamigerum.  
var. *brevisetia* Besch. — Crossidium squamigerum var. *brevisetum*.  
var. *grisea* Vent. — Crossidium griseum.

mendozensis (Mitt.) Jaeg. *op. cit.* p. 275.  
Tortula mendozensis Mitt. *M. austr. am.* p. 154 (1869).  
2. — Am. merid. : And. Chilens.

*microcarpa* Schultz. — B. unguiculata var. *microcarpa*.

*microglottis* C. M. in *Hedwig.* 1898, p. 232.  
2. Terr. ? — Am. sept. : S. Doming.

*nnioides* Schwaegr. — Calyptopogon.  
var. *prostrata* C. M. — Calyptopogon mnioides var. *prostratum*.

- mobilis* C. M. in *Linn.* 1878-79, p. 482.  
2. — *Am. merid.* : Venezuela.
- montana* (Mitt.) *Jaeg. op. cit.* p. 272.  
*Tortula montana* Mitt. *op. cit.* p. 156.  
2. — *Am. merid.* : And. Quitens.
- Moritziana* C. M. — *Didymodon*.  
*mucronata* Brid. — *Dialytrichia Brebissoni*.
- mucronulata* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1894, 11, p. 114.  
Ster. Terr. — *Afr.* : W. Madagascar.
- multiflora* C. M. — *Timmiella*.  
*muralis* var. *capensis* Brid. — *T. muralis* var. *incana* ?  
*mutabilis* Baumg. — *Dicranella Schreberi*.  
*mutica* Brid. — *Tortella caespitosa*.  
*nana* C. M. — *Tortula* ?
- nanocaulis* C. M. in *Hedwig.* 1898, p. 125.  
2. — *Pac.* : Tasman.
- nano-tortuosa* C. M. — *Tortella Knightii*.  
*natalensi-caespitosa* C. M. — *Tortella*.
- natalensis* C. M. in *Rehm. M. Afr. austr. n.* 104 et in *Hedwig.* 1899, p. 106.  
Ster. — *Afr.* : Natal.
- nervosa* { Brid. — *Tortella inclinata*.  
Milde. — *Tortula atro-virens*.  
Schultz. — *Barbula fallax*.  
var. *inclinata* Brid. — *Tortella inclinata*.
- nigrescens* Mitt. *M. Ind. or.* p. 36 (1859).  
*Tortula fallax* Wils. in *Kew Journ. bot.* IX, p. 322 (1857).  
2. — *As.* : W. Himalaya, Sikkim.
- nitida* (Lindb.) Jur. — *Trichostomum*.  
*nivalis* Spr. — *Geheebia gigantea*.  
*Northiana* C. M. — *Tortella caespitosa*.  
*norvegica* Lindb. — *Tortula aciphylla*.
- Novae-Caledoniae* C. M. in *Hedwig.* 1898, p. 128.  
Ster. — *Pac.* : Nov. Caledon.
- novo-granatensis* Hpe. in *Linn.* 1861-62, p. 518.  
*Tortula novo-granatensis* Mitt. *M. austr. am.* p. 158 (1869).  
2. — *Am. merid.* : And. Nov. Granat.  
« var. *gracilior* Hpe. in *op. et l. cc.* (1).  
Ad rad. arb. — *Am. merid.* : Cum typ.
- novo-guineensis* Broth. in *Oefv. af Finska Vet.-Akad. Foerh.* 1895, p. 56.  
2. — *Pac.* : Nov. Guinea.
- obliquifolia* C. M. — *Aloina*.  
*oblongifolia* W. P. Sch. — *Tortula Vahlana*.
- obtusatula* C. M. in *hb. Saharampour, n.* 3745.  
Ster. — *As.* : N. W. Himalaya.

---

(1) « *B. teretiusculas* W. P. Sch. aemula : an spec. propr. ? » Hpe. l. c.



- obtusifolia* { (Schleich.) Schwaegr. — Tortula.  
                   { Schultz. — *B. unguiculata* var. *obtusifolia*.
- obtusula* Lindb. *M. scand.* p. 22 (1879).  
     2. Ad rup. calcar. — *Eur.* : Suecia.
- oenea* (Didymodon?) C. M. et Kindb. in *Mac. Cat.* p. 55 (1892).  
     2. fr. ign. Ad. terr. humid. — *Am. sept.* : Columb. brit.
- olivacea* Besch. *Prodr. bryol. mexic.* p. 37.  
     *Tortula olivacea* Mitt. *M. austr. am.* p. 631 (1869).  
     2. Terr. — *Am. sept.* : Mexic.
- orientalis* (Wild.) Broth. *Natür. Planzenfam. l. c.*  
     *B. indica* Brid. *Bryol. univ. I.* pp. 544 et 828 excl. syn. Willd. (1826);  
     *Bryol. jav. I.* p. 99, t. 81 (1855-61).  
     *Tortula indica* Hook. *M. ex. II.* p. 7, t. 135 excl. syn. Willd. (1820);  
     Mitt. *M. Ind. or.* p. 30 (1859).  
     *Trichostomum indicum* Schwaegr. *Suppl. I, P. I.* p. 142, t. 36 (1811).  
     *Tr. orientale* Willd. *hb.*; C. M. *Syn. I.* p. 568 (1849).  
     2. Muric. et terr. — *As.* : Quang Tcheou Van, Hong Keng, Nepal.,  
     Nepal., Bengal., Ceylan, Java, Banca, ins. Ceram, Borneo. *Pac.* :  
     Nov. Guin. occid. (ins. Kei).
- orizabensis* C. M. in Linn. 1876, p. 63.  
     *B. atlantica* W. P. Sch. in *hb. Mus. Paris* (fd. Besch.).  
     *B. graminicolor* Besch. *Prodr. bryol. mexic.* p. 37 (1871).  
     2. Muric. — *Am. sept.* : Mexic.
- orthodonta* C. M. — *Trichostomum*.
- ovata* Mitt. — *Didymodon*.
- pachyloma* Broth. in Oefv. af Finska Vet.-Akad. Foerh. 1893.  
     *Calymperes dissifolium* ej. in sched.  
     2. — *Pac.* : Nov. Guinea.
- pallens* Brid. — *Tortula agraria*.
- pallido-viridis* C. M. — *Tortella*.
- paludosa* Schleich. *Cent. III*, n. 23; W. et M. *Bot. Taschb.* p. 482 (1807);  
     Schwaegr. *Suppl. I, P. I.* p. 124, t. 30 (1811); Brid. *Bryol. univ. I.*  
     p. 562 (1826); *Bryol. eur. fasc.* 13-15 (*Vol. II*), *Mon.* p. 21,  
     t. 144 (1842); C. M. *Syn. I.* p. 607 (1849); W. P. Sch. *op. cit.*  
     p. 212.  
     *B. crocea* W. et M. *op. cit.* p. 484.  
     *B. crocea* β *paludosa* Roehl. *Deutschl. Fl. III*, p. 79 (1813).  
     *Tortula crocea* Brid. *Sp. M. I.* p. 257 (1806).  
     *T. fallax* β *paludosa* D<sup>na</sup> Fior. *Mass. Bryol. rom. II ed.* p. 8.  
     *T. paludosa* H. et Grev. in *Edinb. Journ. of. sc. I.* p. 301.  
     2. In udis praecip. calcar. — *Eur.* : Norveg. merid., Transilv., Hun-  
     gar., Salisb., Bavar., Francon., Helvet., Wurtemb., Hassia; Ital.  
     super.; deest in Britann. *Afr.* : Alger.
- « var. *Funckiana* (Schultz) *Bryol. eur.*, C. M. et W. P. Sch. *ll. cc.*  
     *B. Funckiana* Schultz *Rec. de Barbula*, p. 26, t. 33, f. 26, (1823);  
     Brid. *Bryol. univ. I.* pp. 512 et 828 (1826).  
     *Tortula crocea* β *humilis* Lindb. de *Tortul.* p. 247 (1864).  
     *T. Funckiana* H. et Gr. in *op. cit.* p. 29.  
     Ut typ. — *Eur.* : liad. loc.

- papillinervis* (Ltz.) Broth. *Natür. Pflanzenfam. l. c.*  
*Hyophila papillinervis* Jaeg. *Ad. I*, p. 205 (1871-72).  
*Pottia papillinervis* Ltz. *Moosst.* p. 160 (1861).  
 Ster. — *Afr.* : Zanzibar.
- papillinervis* C. M. et Kindb. — *Tortula pulvinata subsp. papillinervis.*
- paramattana* (Mitt.) Par. *Suppl. Ind.* p. 29 (1900).  
*Tortula paramattana* Mitt. in *Trans. and Proceed. of the roy. Soc. of Victoria*, 1882, p. 60.  
 2. — *Pac.* : Austral. or. (Nov. Vales).
- parvula* Spreng. — *Tortula recurvata.*
- patagonica* (Mitt.) Jaeg. *Ad. I*, p. 273 (1871-72).  
*Tortula patagonica* Mitt. *M. austr. am.* p. 156 (1869).  
 2. — *Am. merid.* : Patagon.
- patagonica* C. M. — *Tortula fuegiana.*
- pellata* W. P. Sch. — *Pottia.*
- perezilis* C. M. — *Didymodon.*
- perlinealis* C. M. in *Bull. hb. Boissier*, 1897, p. 195.  
 2. — *Am. sept.* : Guatemala.
- perlinearis* C. M. in *Hedwig*, 1899, p. 107.  
 Ster. — *Afr.* : Orange.
- pernana* C. M. in *Linn.* 1878-79, p. 338.  
 2. — *Am. merid.* : Argentin. subtropic.
- perrevoluta* C. M. in *op. cit.* p. 334 (nom. emend. p. 486).  
 2. — *Am. merid.* : Argentin. Cordob.
- « *var. acutifolia* C. M. in *op. cit.* p. 355.  
 Ut et ubi typ.
- « *var. linearifolia* C. M. *op. et l. cc.*  
 Ut et ubi typ.
- pertorquata* C. M. — *Tortula* ?
- perrufula* C. M. — *Tortella.*
- pilifera* (Hook.) Brid. *Bryol. univ. I*, p. 572 (1826); *C. M. Syn. I.* p. 641 (1849); *Bryol. jav. I*, p. 103, t. 83 (1855-61).  
*B. crinita* Schlutz in *Nov. Act. Acad. Leop. XII, I*, p. 226, t. 34 (1823).  
*B. flavipila* W. P. Sch. in *Breutel M. capens.*  
*Tortula pilifera* Hook. *M. ex. t.* 12 (1818).  
 2. — *As.* : Manille, Java. *Afr.* : Prom. B. Sp. *Am. merid.* : Chile.
- « *var. aquatilis* C. M. in *Rehm. M. Afr. austr. n.* 111.  
 — *Afr.* : Prom. B. Sp.
- « *var. gracilis* C. M. *Syn. I*, p. 642.  
 — *As.* : Manille. *Afr.* Prom. B. Sp.
- « *var. sessilis* C. M. in *Rehm. Coll. cit.* p. 113.  
 — *Afr.* : Prom. B. Sp.
- pilifera* Brid. — *Tortula marginata.*
- pilosa* Br., Schultz. — *Tortula laevipila.*
- platyneura* C. M. et Kindb. in *Mac. Cat.* p. 52 (1892).  
 Ster. Ad rup. humid. — *Am. merid.* : Columb. brit.

- plebeja* C. M. in *Linn.* 1880-82, p. 426.  
 Fl. et fr. ign. — *Am. merid.* : Uruguay.
- Podocarp* C. M. — *Tortula* ?
- Poeppigiana* C. M. in *Linn.* 1843, p. 585.  
*Tortula Australasiae* Hook. (*fid. Mitt.*).  
 2. — *Am. merid.* : Chile.
- polyseta* C. M. — *Tortula* ?
- porphyroneura* C. M. — *Tortula* ?
- Potanini Broth.* in *sched.*; C. M. in *Nuov. Giorn. bot. ital.* 1896, p. 99.  
 2. — *As.* : Mongolia.
- Preissiana* } C. M. — *Tortula*.  
                   } Hpe. — *Tortula latrobeana*.  
*princeps* C. M. — *Tortula Müller*.
- pruinosa* (*Mitt.*) *Jaeg. op. cit.* p. 288.  
*Tortula pruinosa* *Mitt. op. cit.* p. 152.  
 2. — *Am. merid.* : And. Quitens.
- pseudo-caespitosa* C. M. — *Tortella*.  
 var. *brachybasis* C. M. — *Tortella ps. caespitosa var. brachybasis*.  
 var. *pungens* C. M. — *Tortella ps. caespitosa var. pungens*.
- pseudo-gracilis* C. M. in *Nuov. Giorn. bot. ital.* 1872, p. 13.  
 2. — *Afr.* : Abyss. sept.
- pseudo-pilifera* Hpe. et C. M. in *Linn.* 1853, p. 493.  
*B. luteola* *Mitt.*  
*B. pungens* H. f. et W. { *fid. Mitt.*  
 2. — *Pac.* : Tasman.
- pseudo-rigidula* *Kindb.* in *Mac. Cat.* p. 264 (1892).  
*B. rigidula* *Mac. Canad. M. n.* 42 p.p.  
 2. — *Am. sept.* : Columb. brit.
- punctulata* *Jaeg.* — *Tortula robusta*.  
*pungens* (H. f. et W.) *Jaeg.* — *Tortula* ?
- purpurea* C. M. in *Fl.* 1875, p. 78.  
*B. brachyphylla* *Sull. et Lesq. M. bor. am. II ed. n.* 135 p.p.  
 2. *Rupic.* — *Am. sept.* : Calif.
- purpurea* *Agh.* — *B. convoluta*.
- purpuripes* C. M. in *Bull. hb. Boissier*, 1897, p. 558.  
 2. *Rupic.* — *Am. sept.* : Jamaic.
- pygmaea* C. M. in *Fl.* 1888, p. 412.  
 Ster. In gramin. — *Afr.* : Kilima N'Djaro.
- pygmaeola* C. M. — *Rhamphidium*.
- rectifolia* *Tayl.* in *Journ. of bot.* 1846, p. 49.  
*Tortula rectifolia* *Mitt. M. austr. am. p.* 158 (1869).  
 2. — *Am. sept.* : Guatemala. *Am. merid.* : And.  
 Nov. Granat., Quitens. et Peruv.

*recurvicuspis* C. M. in *Bull. hb. Boissier*, 1887, p. 557.

B. *Harrisii* Par. *Suppl. Ind.* p. 25 (1900).

2. — *Ann. sept.* : Jamaic.

*recurvifolia* { Mitt. — *Geheebia gigantea*.  
(Wils.) W. P. Sch. — B. *reflexa*.

var. *robusta* Braithw. — B. *reflexa* var. *robusta*.

*recurvipatula* C. M. — Aloina.

*reflexa* (Brid.) Brid. *Mont. M.* p. 93 (1819).

B. *fallax* var. *recurvifolia* *Husn. Musc. gall.* p. 105 (1886).

B. *fallax* γ *reflexa* Brid. *Bryol. univ.* I, p. 556 (1826).

B. *recurvifolia* W. P. Sch. *Coroll. Suppl.* (1856) et *Syn. II ed.* p. 206 (1876).

*Schistidium*? *recurvifolium* Wils. *ms.*; *Spr.* in *Ann. a. Mag. of nat.*

*hist. II ser., III*, p. 491 (1819).

*Tortula fallax* var. *recurvifolia* Wils. *Bryol. brit.* p. 124 (1855).

T. *recurvifolia* *Herk. Handb. brit. M.* p. 258 (1863).

T. *reflexa* Brid. *Sp. M. I.* p. 255 (1806).

2. Ad mur. et rup. calcar. — *Eur.* : Britann., Suecia, Siles., Tatra, M. Beskid., Salisb., Stir., Bavar., Harz, Thuring., vall. Rhén., Jurass., Helvet., Tirol. *As.* : Caucas. *Am. sept.* : New Jersey. Niagara, Colorado.

« var. *robusta* Braithw. *Brit. Moosfl.* p. 263 (1887).

— *Eur.* : Hibern.

*Rehmanni* C. M. in *Rehm. M. Afr. austr.* p. 106 (1890).

2. In albis.

*replicata* Th. in *II. Bryol. europ.*

*Syn. Bryol. europ.*

T. *replicata* Th. in *II. Bryol. europ.*

2. Muric. — *Am. merid.* : And. Quindus.

*revoluta* (Schr.) Brid. in *Schr. Journ. III, P. II*, p. 299 (1801) et *Bryol. univ.* I, p. 571 (1826); *Bryol. eur. fusc.* 13-15 (Vol. II), *Mon.* p. 27, t. 153 (1842); C. M. *Syn.* I, p. 621 (1849); W. P. Sch. *op. cit.* p. 213.

*Bryum stellare nitidum pallidum*, capsulis tenuissimis Dill. *Hist. M.* p. 381, t. 48, f. 44 (1741) et *hb.*

*Tortula nervosa* Engl. *Bot. n.* 2383.

T. *revoluta* Schrad. *Syst. Samml. krypt. Gew.* I, n. 54 (1796).

2. Muric. — *Eur.* : Ccpiosior in merid. quam in med., var. in sept. *As.* Persia. *Afr.* : Alger., Tunet., Canar.

*revoluta* W. et M. { — B. *Hornschuchiana*.  
var. *Hornschuchiana*

*revolvens* W. P. Sch. *Syn. II ed.* p. 195 (1876).

B. *atro-virens* var. *robusta* ej. in *Husn. M. Gall. n.* 513 a.

*Desmatodon revolvens* *Husn. Musc. gall.* p. 96 (1885).

1 (W. P. Sch.!). Ad mur. muscos. — *Eur.* : Monspel., Galloprov.

*rhaetica* Amann in *Rev. bryol.* 1889, p.p. 51 et 90.

Ster. Ad terr. lapidos. — *Eur.* : Alp. Rhaet.

*rigida* { Brid. — Aloina *ambigua*.  
Schultz. — Aloina.

- var. *mucronulata* Bryol. eur. }  
 var. *obtusa* (Jaeg.) Par. } — *Aloina rigida* varr. ejusd. nom.  
 var. *pilifera* Bryol. eur. }
- rigidicaulis* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 255.  
 Ster. — *As.* : China centr. (prov. Schen-Si).
- rigidula* { Besch. — B. Bescherellei.  
 { Mac. p.p. — B. pseudo-rigidula.  
 { (Hedw.) W. P. Sch. — Didymodon.  
 var. *densa* W. P. Sch. — Didymodon rigidulus var. *densus*.
- riparia* C. M. in *Linn.* 1880-82, p. 428.  
 2. Terr. — *Am. merid.* : Alp. Argentin. subtropic.
- robustifolia* C. M. et Kindb. in *Mac. Cat.* p. 56 (1892).  
 B. *rubiginosa* Mac. *Canad. M.* n. 75.  
 Ster. Rupic. — *Am. sept.* : Columb. brit., Washington, ins. Vancouver.
- romana* C. M. — *Dialytrichia Brebissoni*.  
*rosulata* C. M. — *Timmiella*.  
*rotundata* Lindb. et Arn. — Didymodon.  
*rotundifolia* Jens. — *Tortula papillosa*.  
*rubella* { (H. f. et W.) Jaeg. — *Tortula*.  
 { Mitt. — Didymodon.  
 var. *brevifolia* L. et A. — Didymodon *rubellus* var. *brevifolia*.
- rubiginosa* Mitt. in *Journ. of the Linn. Soc.* 1864, p. 27, t. 5; *Lesq. et Jam. Man.* p. 126 (1884).  
*Bottia rubiginosa* Wats. *Bot. Calif.* II, p. 362.  
 2. — *Am. sept.* : Ins. Vancouver, Oregon.
- rubiginosa* { C. M. — Didymodon *argentinicus*.  
 { Mac. — B. *robustifolia*.
- rubra* { Lindb. — Didymodon.  
 { Mitt. — *Tortula*.  
*rufa* { Jur. — Didymodon.  
 { W. P. Sch. — *Tortula*.  
*rufescens* Mitt. — *Hyophila*.  
*rufidula* C. M. — Didymodon.
- rufipes* W. P. Sch. in *Besch. Prodr. bryol. mexic.* p. 36 (1871).  
 2. — *Am. sept.* : Mexic.
- rufiseta* Tayl. — B. *Australasiae*.  
*ruraliformis* Besch. { — *Tortula ruralis* var. *ruraliformis*.  
 f. *gigantea* Rob. Farn. {
- ruralis* (L.) Hedw. var. *calva* Bryol. eur. — *Tortula montana* var. *calva*.  
 var. *pulvinata* Limpr. — *Tortula pulvinata*.  
 var. *rupestris* Bryol. eur. — *Tortula montana*.
- ruicola* C. M. in *Bot. Zeit.* 1875, p. 381.  
 1. Muric. — *Am. merid.* : Brasil.
- ruvenzorensis* (Broth.) Par. — *Trichostomum*.  
*Sartorii* Jaeg. — *Desmatodon*.  
*Saussuriana* Brid. — *Ceratodon purpureus*.
- Schenkii* Broth. in *Hedwig.* 1894, p. 128.  
 Ps. 2. Ad rup. humid. — *Am. merid.* : Brasil. or.

- schensiana* C. M. in *Nuov. Giorn. bot. ital.* 1896, p. 101 et 1898, p. 182.  
2. Rupic. — As. : China centr. (prov. Schen-Si).
- « *var. longifolia* C. M. in *op. et ll. cc.*  
Ster. — As. : Ubi typ.
- « *var. tenuissima* C. M. in *op. et ll. cc.*  
Ut et ubi typ.
- Schreberi* Brid. — *Dicranella*.
- scleromitra* Besch. in *Bull. Soc. bot. Fr.* 1894, p. 81.  
2. In udis coctis. — As. : Tonkin. super.
- sedifolia* C. M. — *Aloina*.
- semi-torta* Sull. in *Pac. R. R. IV*, p. 186, t. 3; *Lesq. et Jam. Man.* p. 126 (1884).  
2. — Am. sept. : Calif.
- serpenticaulis* C. M. in *Nuov. Giorn. bot. ital.* 1898, p. 183.  
2. — As. : China centr. (prov. Schen-Si).
- serripungens* Ltz. et C. M. — *Tortula*.
- serrulata* H. et Gr. (Brid.) — *Tortula*.
- setacea* Hedw. — B. convoluta.
- setifolia* (C. M.) Broth. *op. et l. cc.*  
*Trichostomum setifolium* C. M. in *Hedwig.* 1898, p. 234; *Par. Suppl. Ind.* p. 325 (1900).  
Ster. Ad flum. — Am. sept. : Porto-Rico.
- sinensi-fallax* C. M. in *Hedwig.* 1897, p. 100.  
Ster. — As. : China centr. (prov. Schen-Si)
- sinensis* C. M. — *Tortula alpina*.
- sinuosa* (Wils.) Jaeg. *Ad. I*, p. 286 (1871-72); *Braithw. Brit. Moossfl* p. 270 (1887).  
B. cylindrica γ *sinuosa* Lindb. *M. scand.* p. 22 (1879).  
B. vinealis « *luxurians* » Jur. mss. ; *Geh. in Fl.* 1864, p. 80.  
*Dicranella sinuosa* Wils. mss. ; *Seem. Journ. bot.* 1871, p. 294.  
*Didymodon sinuosus* W. P. Sch. *Syn. II ed.* p. 166 (1876); *Par. Ind.* p. 380 (1895).  
*Tortula sinuosa* Mitt. in *Lond. Journ. of bot.* 1867, p. 327.  
*Trichostomum sinuosum* Lindb. in *Seem. Journ. bot. l. c.*  
2. Muric. et terr. — Eur. : Anglia, Belgia, Gall. sept. or. As.  
Caucas. Am. sept. : Washington.
- Solmsiana* Hpe. — B. convoluta *var. commutata*.
- Somaliae* C. M. in *Linn.* 1876, p. 294.  
Ster. — Afr. : Somali.
- sordida* Besch. in *Bull. Soc. bot. Fr.* 1894, p. 80.  
2. — As. : Tonkin.
- spadicea* Mitt. — *Didymodon*.
- sparsidens* C. M. et Kindb. in *Mac. Cat.* p. 51 [1892].  
2. Terr. — Am. sept. : Columb. brit., Wisconsin.
- sparsifolia* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1894, II, p. 115.  
Ster. — Afr. : Ins. Bourbon.
- spathulata* Dz. et Mk. — *Hyophila commutata*.

- speirostega* C. M. in *Hedwig*. 1898, p. 125.  
 Ster. — *Pac.* : Austral. or. (Nov. Vales).
- spiralis* W. P. Sch. in C. M. *Syn.* I, p. 622 (1849).  
*Tortula spiralis* Mitt. *M. austr. am.* p. 151 (1869).  
 2. — *Am. sept.* : Mexic.
- Sprengelii* Schwaegr. — *Trichostomum Berteroanum*.  
*squamigera* Viv. — *Crossidium*.  
*squarrosa* Brid. — *Tortella*.  
 var. *nitida* R. Farn. — *Tortella squarrosa* var. *nitida*.  
*stellata* Brid. — *Tortula agraria*.
- stenocarpa* Hpe. in *Ann. sc. nat. V ser., III*, 1866, p. 346.  
*Tortula stenocarpa* Mitt. *op. cit.* p. 162.  
 2. Terr. — *Am. merid.* : And. Nov. Granat.
- stenophylla* (Mitt.) Jaeg. — *Trichostomum*.  
*stricta* Hedw. — *B. unguiculata*.
- strictidens* C. M. in *Bull. hb. Boissier*, 1897, p. 192.  
 2. — *Am. sept.* : Guatemala.
- Stuhlmanni* Broth. *op. et l. cc.*  
*Anoctangium Stuhlmanni* ej. in *Engler's Bot. Jahrb.* 1897, p. 177.  
 2. fr. ign. Terr. — *Afr.* : Zanzibar.
- subcaespitosa* (Hpe.) Broth. *op. et l. cc.*  
*Anacalypta subcaespitosa* Hpe. in *Linn.* 1863, p. 127.  
*Pottia subcaespitosa* Par. *Ind.* p. 1029 (1897).  
*Weisia subcaespitosa* Mitt. *M. austr. am.* p. 140.  
 1 (?). Terr. — *Am. merid.* : And. Nov. Granat.
- subcalycina* C. M. in *Linn.* 1867-68, p. 617.  
 2. — *Pac.* : Austral. or.
- subcomosa* Broth. in *Hedwig*. 1899, p. 211.  
 2. — *As.* : Japon.
- subconvoluta* C. M. in *Nuov. Giorn. bot. ital.* 1898, p. 183.  
 2. — *As.* : China centr. (prov. Schen-Si).
- subcylindrica* Broth. in *Bot. Centralbl.* 1890, n. 51 et in *Hedwig*.  
 1893, p. 205.  
 2. fr. ign. — *Am. sept.* : Washington.
- subduriuscula* C. M. — *Trichostomum*.  
*suberecta* Kindb. — *Desmatodon*.
- suberythropoda* C. M. in *Bull. hb. Boissier*, p. 194.  
 Ad rup. hument. — *Am. sept.* : Guatemala.
- subfallax* C. M. in *Bot. Zeit.* 1862, p. 338; *Lesq. et Jam. Man.*  
 p. 221 (1884).  
*B. fallax* Sull. in *Bot. R. R. Survey* (Whipple), p. 186.  
 2. — *Am. sept.* : S. Nevada Calif.
- subglaucescens* C. M. — *B. boliviana*.
- subgracilis* C. M. et Kindb. in *Mac. Cat.* p. 53 (1892).  
 2. Rupic. — *Am. sept.* : Columb. brit., Montana.  
 « var. *viridior* Kindb. in *op. cit.* p. 263.  
 Ut typ. — *Am. sept.* : Ontario.

- subicmadophila *C. M. et Kindb. in op. cit. p. 53.*  
*B. gracilis* var. *squarrulosa* Kindb. in *Bull. Torr. bot. Club, XVII*,  
 p. 89 (1890).  
 2. Ad rup. humid. — *Am. sept.* : Columb. brit.  
*subnigra* (Mitt.) Jaeg. — Didymodon?  
*subpellucida* Mitt. *M. Ind. or. p. 35* (1859).  
 2. — *As.* : Tibet., N. W. Himalaya.  
*subpilosa* Brid. — *Tortula marginata*.  
*subrevoluta* } *C. M.* — *B. apoclada*.  
 } Hpe. — Didymodon.  
*subtorquata* Hpe. et *C. M. in Linn. 1853, p. 492; F. v. Müll. Analyt.*  
*draw. Austr. M. t. 3* (1864).  
 Ad terr. arenos. — *Pac.* : Austral. or., Tasman.  
*subtortuosa* *C. M.* — *Tortella tortuosa*.  
*subulata* (L.) P. B. var. *inermis* Bryol. eur. — *Tortula inermis*.  
 var. *integrifolia* Boul. — *T. subulata* var. *angustata*.  
*subulifolia* Sull. in *Proced. of the Amer. Acad. 1861, p. 277.*  
*B. crispula* Hpe. in *Jaeg. Ad. I, p. 286* (1871-72).  
*B. gracilis* Sull. in *op. et l. cc.*  
*Tortula subulifolia* Mitt. *M. austr. am. p. 161* (1869) (1.).  
 2. Ad rup. irror. — *Am. sept.* : Cuba. *Am. merid.* : Guian., And.  
 Quitens.  
*subunguiculata* W. P. Sch. in *Savatier M. Japon. n. 203; Besch. in*  
*Ann. sc. nat. 1893, p. 337.*  
 2. — *As.* : Japon.  
*Sullivaniana* *C. M.* — *Aloina*  
*Swarziana* *C. M. in Hedwig. 1898, p. 232.*  
 — *Am. sept.* : Jamaic.  
*syrrhopodontoides* Rehm. *M. Afr. austr. n. 175.*  
 — *Afr.* : Transvaal.  
*tasmanica* Hpe. — *Tortula*?  
*tectorum* *C. M. in Nuov. Giorn. bot. ital 1896, p. 101.*  
 2. Ad tecta. — *As.* : China centr. (prov. Schen-Si).  
*tenuirostris* Brid. — *B. angustifolia* (2).  
*tenuis* Kindb. — *Leptobarbula berica*.  
*teretinscula* W. P. Sch. in *C. M. Syn, I. p. 614* (1849).  
*Tortula teretiuscula* Mitt. *M. austr. am. p. 160* (1869).  
 2. — *Am. sept.* : Mexic.

(1) Sec. Hpe. in Jaeg. op. cit. pp. 269 et 286, pl. Wright. cubens. n. 32 est *B. subulifolia*, pl. n. 27 *B. crispula*. Cl. Mitt. n. 27 et 32 sub nom. *B. subulifoliae* conjungit.

(2) *tenera* Brid. *Bryol. univ. I. pp. 577 et 835.*  
*Bryum tenerum* Vill. Pl. Dauph. III,  
 p. 366 (excl. syn. Dill.) } Quid? Deleatur.  
*Tortula tenera* P. B. Prodr. p. 93.



- thrausta* C. M. in *Dusen M. Camer. n.* 352.  
— Afr. : Cameroon.
- ticinensis* Kindb. — *Tortula muralis* var. *ticinensis*.
- tokyensis* Besch. in *Journ. de bot.* 1898.  
2. — As. : Japon.
- tophacea* Mitt. — *Trichostomum*.
- torquata* Tayl. in *Hook. Journ. of bot.* 1846, p. 50; C. M. *Syn. I.*, p. 611 (1849); *Hook. f. et Wils. Fl. of the N. Zeal. II*, p. 70 (1855) et *Handb.* p. 419 (1867).  
*Tortula torquata* Mitt. in *Journ. of the Linn. Soc.* 1859, p. 70.  
— Pac. : Austral., Nov. Seeland. sept., Tasman.
- torquatifolia* Geheeb in *Bull. hb. Boissier*, 1896, p. 409.  
2. — Afr. : Ondonga merid. or.
- torquescens* W. P. Sch. in *Bot. Zeit.* 1853, p. 163; C. M. in *Hedwig.* 1899, p. 105 (*descr. locupletior*).  
B. Mauchii Rehm. *M. Afr. austr. n.* 109.  
2. — Afr. : Prom. B. Sp.
- tortellifolia* C. M. et Kindb. in *Mac. Cat.* p. 156 (1892).  
B. cylindrica Mac. *Canad. M. n.* 81 p. p.  
B. flexifolia ej. *Coll. cit.* p. 399.  
2. Ad rup. marit. humid. — Am. sept. : Columb. brit., ins. Vancouver.
- tortelloides* C. M. in *Linn.* 1878-79, p. 337.  
2. — Am. merid. : Argentin. subtropic.
- tortuosa* (L.) W. et M. — *Tortella*.  
var. *angustifolia* Jur. { — *T. tortuosa* varr. ejusd. nom.  
var. *fragilifolia* Jur. {  
var. *incrassata* Brid. — *T. tortuosa*.  
var. ? *longifolia* Ren. — *T. tortuosa* var. *longifolia*.  
var. *nitida* Ren. — *Trichostomum nitidum*.  
var. *pseudo-fragilis* Ther. { — *T. tortuosa* varr. ejusd. nom.  
var. *viridis* Delogne }
- trachyphylla* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 259.  
B. eroso-denticulata C. M. in *op. cit.* 1896, p. 102.  
2. Ad saxa rivul. — As. : China centr. (prov. Schen-Si).
- trichostomacea* C. M. in *Rehm. M. Afr. austr. n.* 97 et 483, et in *Hedwig.* 1899, p. 108.  
2. — Afr. : Prom. B. Sp.  
var. *chlorophyllosa* C. M. — B. *trivialis*.
- trichostomifolia* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 256.  
2. — As. : China centr. (prov. Schen-Si).
- trichostomoides* Besch. *Prodr. bryol. mexic.* p. 38 (1871).  
— Am. sept. : Mexic.
- trifaria* Mitt. — *Didymodon luridus*.
- trivialis* C. M. in *Hedwig.* 1899, p. 107.  
B. *trichostomacea* var. *chlorophyllosa* ej. in *Rehm. M. Afr. austr. n.* 98; *Par. Ind. p.* 99 (1894).  
— Afr. : Prom. B. Sp., Orange.

*Turneri* Brid. — *Didymodon rubellus*.

*Uleana* C. M. in *Ule Bryoth. brasil. n.* 14.

— *Am. merid.* : Brasil. austr. or.

*umbrosa* C. M. in *Linn.* 1878-79, p. 348.

2. Muric. et terr. — *Am. merid.* : Argentin. Cordob. et subtropic.

*uncinicomma* C. M. in *op. cit.* p. 345.

2. — *Am. merid.* : Argentin. Cordob.

*unguiculata* (Huds.) Hedw. *Fund. M. II*, p. 92 (1782); *Brid. Bryol. univ. I*, p. 563 (1826); *Bryol. eur. fasc.* 13-15 (*Vol. II*), *Mon.* p. 19, t. 142 (1842); C. M. *Syn. I*, p. 612 (1849); W. P. Sch. *op. cit.* p. 203.

B. *amœna* Schum. *Fl. Sacl. II*, p. 48 (1801); *Brid. op. cit.* p. 538.

B. *aristata* Brid. *Mant. M.* p. 92 (1819).

B. *dubia* ej. *op. et l. cc.*

B. *incerta* Schum. *op. cit.* p. 44.

B. *lanceolata* Schum. *op. cit.* p.

B. *linoides* Brid. *Mant. M.* p. 90 et *Bryol. univ. I*, p. 543.

R. *stricta* Hedw. *Sp. M.* p. 119, t. 26 (1801); *Brid. op. cit.* p. 540.

B. *unguiculata* var. *aristata* Brid. *op. cit.* p. 547.

B. *unguiculata* var. *dubia* ej. *op. et l. cc.*

Bryum angustis { pallidis foliis, capitulis erectis, longioribus pediculis insidentibus, calyptra rectiore Dill. *Cat. Giss.* p. 225 (1718).  
viridibus foliis, capitulis erectis, brevibus pedunculis insidentibus, calyptra falcata vel avium unguiculos referente Dill. *op. et l. cc.*

Br. *aristatum* Dicks. *Pl. crypt. fasc. III*, p. 12, t. II, f. 7 (1793).

Br. *barbatum* Curt. *Fl. Lond. fasc. IV*, t. 65 (1777-92).

Br. *linoides* Dicks. *op. cit.* p. 8, f. 3.

Br. *mucronulatum* ej. *op. cit.* p. 3.

Huds. *Fl. angl.* p. 410 (1762).

Br. *unguiculatum* { surculis in summitate crassioribus Dill. *Hist. M.* p. 383, t. 48, f. 47 (1741).  
et *barbatum* { tenuius et stellatum ej. *op. cit.* p. 384, t. 48, f. 48.

*Dicranum laxum* Brid. *Musc. rec. II, P. I*, p. 175 excl. *syn.* 1789).

*Mollia unguiculata* Schrank Bayer. *Fl. II*, p. 457 (1798).

*Orthopyxis aristata* P. B. *Prodr.* p. 79 (1805).

*Tortula amœna* Brid. *Sp. M. I*, p. 249 (1806).

T. *aristata* Sm. *Fl. brit. III*, p. 1261 (1804-5).

T. *barbata* ej. *op. cit.* p. 260.

T. *decepiens* Sibth. *Fl. Oxon.* p. 285 (1794).

T. *dubia* P. B. *Prodr.* p. 32 (1805).

T. *fallax* H. et Grev. in *Edinb. Journ. of sc. I*, p. 300.

T. *humilis* Turn. *Musc. hib.* p. 45 (1804).

T. *linoides* Brid. *Mant. M.* p. 90 et *Bryol. univ. I*, p. 543.

T. *mucronulata* Sm. *op. cit.* p. 1250.

T. *stricta* Brid. *Sp. M. I*, p. 249.

T. *unguiculata* Roth. *Tent. fl. germ. III, P. I*, p. 206 (1800).

*Trichostomum linoides* Brid. *op. cit.* p. 234.

2. Ad mur. et rup., agger. et vias. — *Eur.* : Tota, vulgatiss. *As.* : Japon., Hong-Kong, Jenissei med., *Caucas.* *Afr.* : Alger. *Am. sept.* : Canada or., Columb. brit., fere tota.

• var. *apiculata* (Hedw.) *Bryol. eur. et W. P. Sch. II. cc.*

B. *apiculata* Hedw. *Sp. M.* p. 117, t. 26, f. 1-1 (1801); *Brid. Bryol. univ. I*, pp. 560 et 831.

*Tortula apiculata* Brid. *Sp. M. I*, p. 259.

T. *unguiculata* var. *apiculata* H. et T. *Musc. brit.* p. 30 (1818).

*Trichostomum pensilvanicum* Spreng. *Syst. veg. IV, P. I*, p. 170.

Loc. argilloso-calcar., cum typ.

var. *aristata* B. — *B. unguiculata*.

« var. *brachypus* C. M. *op. cit.* p. 613.

*B. brachypus* Brid. *Bryol. univ. I*, p. 551.

*Tortula acuminata* ej. *Sp. M. I*, p. 252.

Ut typ. — Eur. : Thuring., Catalaun.

« var. *cuspidata* (Schultz) Brid. *Bryol. univ. I*, p. 565 ; *Bryol. eur. l. c.* t. 143 ; C. M. *op. cit.* p. 613 ; W. P. Sch. *op. cit.* p. 204.

*B. cuspidata* Schultz *Rec. de Barbula*, p. 14, t. 32, f. 14 a (1823).

*Bryum nervosum* Hoffm. *Deutschl. Fl. II*, p. 44 (1795).

*Tortula mucronulata* Sw. *M. Succ.* p. 40 (1799).

*T. mucronulata* β Turn. *Musc. hib.* p. 47 (1804).

Ut typ. — Eur. : lisd. loc. ; Sardin. Afr. : Alger.

var. *dubia* Brid. — *B. unguiculata*.

« var. *fastigiata* (Schultz) *Bryol. eur. l. c.*

*B. fastigiata* Schultz *op. cit.* t. 33 ; Brid. *Bryol. univ. I*, p. 554.

Ut typ. — Eur. : Reg. alp.

var. *latifolia* de Breb. — *Dialytrichia Brebissoni*.

« var. *microcarpa* (Schultz) *Bryol. eur. et W. P. Sch. II. cc.*

*B. microcarpa* Schultz *op. et l. cc.* ; Brid. *op. cit.* p. 561.

Ut et cum var. *apiculata*.

« var. *nitido-costata* R. Farn. in *M. Prov. Pavia*, 1891, p. 20.

Ad rup. umbros. — Eur. : Ital. super.

« var. *obtusifolia* (Schultz) *Bryol. eur. et W. P. Sch. II. cc.*

*B. lanceolata* Hedw. *Sp. M.* p. 119, t. 26 (1801) ; Brid. *op. cit.* p. 568.

*B. obtusifolia* Schultz *op. et l. cc.*

*Tortula lanceolata* P. B. *Prodr.* p. 92 (1805) ; Brid. *Sp. M. I*, p. 260 (1806).

In agris. — Eur. : Rarior.

« var. *patagonica* C. M. in *Linn.* 1843, p. 583 et *Syn. I*, p. 613 (1849).

*B. ericetorum* Brid. *Musc. rec. II, P. I*, p. 196 (1798).

*Bryum ericetorum* Dickx. *Pl. Crypt. fasc. II*, p. 5 ? (1790).

*Br. ruralis* facie, foliis minoribus et non pilosis Dill. *Hist. M.* p. 354, t. 45, f. 13 (1741).

*Syntrichia ericetorum* Brid. *Bryol. univ. I*, p. 588 (1826).

*Tortula ericetorum* Sm. *Fl. brit. III*, p. 1258 (1804).

Ut typ. — Eur. : Scotia ? Am. merid. : Patagon.

*unguiculata* Dz. et Mk. — *Hyophila Sieboldi*.

*unguiculatula* C. M. in *Linn.* 1878-79, p. 333.

2.

— Am. merid. : Alp. Argentin. subtropic.

*uruguayensis* Broth. in *Bihang till K. Sv. Vet.-Akad. Foerh.* 1900.

2.

— Am. merid. : Uruguay.

*uruquensis* Par. — *B. decidua*.

*vaginans* Lindb. — *Didymodon spadiceus*.

*vancouveriensis* Kindb. — *Timmiella*.

*Ventanae* C. M. in *Helwig.* 1897, p. 102.

2.

— Am. merid. : Argentin. patagon.

*vinealis* Brid. *Bryol. univ.* I, p. 830 (1826); *Bryol. eur. fasc.* 13-15 (Vol. II). *Mon.* p. 24, t. 148 (1842); *C. M. Syn.* I, p. 617 (1849); *W. P. Sch. op. cit.* p. 209.

*B. cylindrica* var. *vinealis* Lindb. *M. scand.* p. 22 (1879).

*B. fallax* var. *vinealis* Hüb. *Musc. germ.* p. 327 (1833).

*Tortula fallax* var. *vinealis* de Not. in *Mém. Acad. Torin.* XL, p. 319 (1838).

*T. vinealis* Spr. in *Hook. Journ. bot.* IV, p. 194 (1845).

*Weisia monocladus* Brid. *op. cit.* p. 360.

2. Ad mur. et terr. lapid. calcar., rarius arenac., sæpius steril. — *Eur.* : Ins. Faroër, Britann., Scand. et Fenn. merid., Dania, Siles., Boruss. or., Saxon., Harz, etc., vall. Rhen., Galloprov., Ital., ins. Melite. *As.* : Yunnan, Nepal., Kurdistan, Persia, Minor. *Afr.* : Alger., Tunet. *Am. sept.* : Terra nov., Ontario, Washington, Calif., Mexic.

• var. *cylindrica* (Tayl.) Boul. *M. Fr.* p. 430 (1884).

*B. Beecheyi* Lesq. *ms.*; *Watson Bot. Calif.* II, p. 372.

*B. cylindrica* W. P. Sch. in *Hedwig.* 1873, p. 47 et *Syn.* II ed. p. 208 (1876).

*B. insulana* Husn. *M. N. O.* p. 81 (1873).

*B. vinealis* var. *flaccida* *Bryol. eur. l. c.*

*Tortula cylindrica* Lindb. in *Bot. Not.* 1865, p. 76.

*T. insulana* de Not. in *Mém. Acad. Torin.* XL, p. 320 (1838).

*T. vinealis* Wils. *Bryol. brit.* p. 124, t. 42 (1855).

*Zygotrichia cylindrica* Tayl. in *Mackay Fl. hib.* II, p. 26 (1836).

Lignic. et ad terr. arenos. humid. — *Eur.* : Britann., Scand., Boruss. or., Voges., Sardin., etc., ubi typ. *As.* : Caucas. *Afr.* : Alger., *Am. sept.* : Fret. Behring, ins. Vancouver, Columb. brit., Washington, Oregon, Montana, Calif.

var. *flaccida* *Bryol. eur.* — *B. vinealis* var. *cylindrica*.

*virescens* Lesq. in *Trans. of the Am. Phil. Soc.* XIII, p. 4; *Lesq. et Jam. Man.* p. 124 (1884).

2. Rupic. — *Am. sept.* : Calif.

*viridi-flava* Kindb. — *Trichostomum*.

*viridis* W. P. Sch. — *B. gracilis* var. *viridis*.

*vulcanica* Ltz. *Moosst.* p. 161 (1864).

2. Rupic. — *Am. merid.* : And. Quitens.

*Waghornei* Kindb. *Eur. and N. Amer. Bryin.* p. 264 (1897).

— *Am. sept.* : Terra nov.

*Wallichii* (Mitt.) Jaeg. — *Didymodon*.

*Willdii* Broth. — *Didymodon*.

*Wilhelmii* C. M. — *Calyptopogon*.

*Woodii* W. P. Sch. — *Amphoridium Mougeotii*.

*Wrightii* Sb. in *Jaeg. Ad.* II, p. 673 (1877-78).

*B. obscura* Sull. in *Proced. of the Amer. Acad.* 1861, p. 277; *Jaeg. Ad.* I, p. 280 (1871-72).

*Tortula obscura* Mitt. *M. austr. am.* p. 150 (1869).

2. — *Am. sept.* : Cuba.

*xanthocarpa* C. M. — *Tortella*.

*Zetterstedtii* Kindb. — *Didymodon spadiceus*.

*zygodontifolia* C. M. — *Didymodon rubellus*.

**BARTRAMIA** Hedw. Descr. II, p. III, t. 40 (1789). — 104 spec.

Species omnes non indicatae sunt PHILONOTIDES ejusd. nom.

*abyssinica* C. M. — Anacolia.

*acerosa* C. M. et Hpe. in Linn. 1856, p. 208.

B. papillata H. f. et W. (fd. Mitt.).

2. — Pac. : Austral. or.

*aciphylla* Wils. — Breutelia.

*affinis* } Hsch. — Philonotis erythrocaulis.  
 } Schwaegr. — Breutelia commutata.

*afro-ithyphylla* Broth. in Engler's Bot. Jahrb. p. 1897, p. 249.

Syn. Ad arbor. — Afr. : Kilima N'Djaro.

*afro-scoparia* C. M. — Breutelia.

*afro-stricta* C. M. in Rehm. M. Afr. austr. n. 203-204 et in Hedwig. 1899, p. 94.

Ster. — Afr. : Pr. B. Sp.

*alpina* Schleich. — B. Oederi.

*ambigua* Mont. in Ann. sc. nat. 1845, IV, p. 103 et Syll. p. 28 (1856);  
 C. M. Syn. I, p. 502 (1849).

2. Rupic. et terr. — Am. merid. : Peruv., Chile.

*angulata* Tayl. — Philonotis rufflora.

*angustata* Wils. } — Philonotis angusta.  
 } p.p. — Philonotis fontana var. falcata.

*angustifolia* Mitt. M. austr. am. p. 273 (1869).

B. longifolia ej. in Jameson M. exs.

— Am. merid. : And. Quitenses.

*aprica* C. M. in Linn. 1875, p. 397.

Syn., fr. ign. Rupic. — Afr. : Littor occ. mar. Erythraei.

*aristaria* C. M. — Glyphocarpus.

*aristata* W. P. Sch. in Lechl. Pl. Chil. n. 834; C. M. in Bot. Zeit. 1862, p. 338.

Ad rup. madid. — Am. merid. : Chilè.

*aristata* Mitt. — B. aristifolia.

*aristifolia* Jaeg. Ad. I, p. 533 (1873-74).

B. aristata Mitt. M. austr. am. p. 269 (1869).

2? Rupic. — Am. merid. : And. Quitens.

*aristifolia* Broth. — B. Jungneri.

*arcuata* { Brid. — Breutelia.  
 } Schwaegr. p.p. } Breutelia.  
 } Breutelia tomentosa.

*auricola* C. M. in Nuov. Giorn. bot. ital. 1897, p. 42.

2. — Am. merid. : And. Boliv.

*australis* Mitt. { (M. austr. am.) — Conostomum.  
 } (1876). — Philonotis anisothecioides?  
 } (Handb.) — Philonotis.

*austro-arcuata* C. M. — Breutelia.

*austro-georgica* Par.

*B. subpatens* C. M. *Bryol. Austr. Georg.* p. 29 (1890).

Syn. — *Am. merid.* : Austr. Georg.

*azorica* Mitt. — *Breutelia*.

*Baldwinii* C. M. in *Fl.* 1896, p. 448.

— *Pac.* : Ins. Hawai.

*bogotensis* Hpe. — *Q. flavicans*.

*boschbergiana* C. M. — *Philonotis afro. fontana* var. *breviseta*.

*Boulayi* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1890, I, p. 177; *Ren.*

*M. Masc. Mad.* n. 174 et *Prodr.* p. 171 (1897).

Fr. ign. — *Afr.* : Ins. Borboniae.

*brachyphylla* C. M. — *Cryptopodium*.

var. *viridissima* C. M. — *Cryptopodium* var. *viridissima*.

*brachypus* Br. et W. P. Sch. in W. Sch. *M. Abyss.* n. 441; C. M.

*Syn.* I, p. 497 (1849).

Syn. — *Afr.* : Alp. Abyss.

*Breutelii* W. P. Sch. in *Breut. M. capens*; C. M. in *Bot. Zeit.* 1858, p. 162.

2. — *Afr.* : Pr. B. Spei.

*brevifolia* Brid. *Bryol. univ.* II, p. 737 (1827); C. M. *Syn.* II, p. 619 (1851).

B. *Mülleri* Mont. in C. M. *Syn.* I, p. 498 (1849) et *Syll.* p. 29 (1856); Mitt. *M. austr. am.* p. 273 (1869).

B. *patens* Hook. in *Kunth Pl. Aequinoct.* p. 55 (1822-28).

B. *potosica* C. M. in *Linn.* 1847, p. 204; *Hpe. ibid.* p. 76.

2 ? — *Am. merid.* : Columb.

*brevifolia* W. P. Sch. — B. *Schimperi*.

*breviseta* } Boul. — B. *ithyphylla* var. *strigosa*.

} Lindb. — B. *ithyphylla* var. *breviseta*.

*Brownii* Brid. — *Philonotis affinis*.

*caespitosa* Wils. — *Philonotis fontana* var. *caespitosa*.

*calabrica* Lindb. — *Philonotis rigida* var. *gracilis*.

*Campylopus* W. P. Sch. in C. M. *Syn.* II, p. 619 (1851).

— *Am. sept.* : M. Mexic.

*carinata* Mitt. — *Breutelia*.

*chilensis* Ltz. — *Breutelia*.

*chrysea* C. M. — *Breutelia* ?

*chrysura* C. M. — *Glyphocarpus*.

*circinulata* C. M. et Kindb. in *Mac. Cat.* p. 105 (1892).

2 ? *Rupic.* — *Am. sept.* : Alaska, Columb. brit.

*commutata* Mitt. in *Journ. of the Linn. Soc.* 1863, p. 153.

B. *patens* *Schwaegr. Suppl.* I, II, p. 55, t. 62 (fid. Mitt.).

Syn. — *Afr.* : M. Cameroon.

*commutata* Hpe. — *Breutelia*.

*comosa* } Hpe. et C. M. — *Glyphocarpus*.

} Hook. f. — *Breutelia*.

*compacta* Hsch. — *Glyphocarpus*.

*condensata* *Schwaegr.* — B. *Æderi*.

*conostoma* *Bryol. eur.* — *Conostomum boreale*.

*consimilis* Hook. f. — Breutelia.

*costa-ricensis* C. M. *Syn. II*, p. 618 (1851).

Syn. — *Am. sept.* : Costa-Rica.

*crassa* H. f. et W. — Breutelia.

*crassicaulis* C. M. — Breutelia.

*crassinervia* Mitt. in *Handb. of the N. Zeal. Fl.* p. 417 (1867).

Syn. *Silvic.* — *Pac.* : Nova Seelandia merid.

*crassinervis* Wils. p.p. } — *Philonotis calcarea*.  
                                       } — *Philonotis fontana*.  
                                       } — *Philonotis lutea*.

*crispa* Sw. — *B. pomiformis* var. *crispa*.

var. *major* W. et M. — *B. pomiformis* var. *crispa*.

var. *minor* W. et M. — *B. pomiformis*.

var. *nana* Hartm. — *B. pomiformis*.

var. *nigrescens* Kindb. — *B. pomiformis*.

var. *Normanni* Holmgr. — *B. pomiformis* var. *heteromalla*.

var. *pomiformis* Lindb. — *B. pomiformis*.

*crispata* W. P. Sch. in *Savatier M. Japon n.* 511; *Besch. in Ann. sc. nat.* 1893, p. 348.

l. Ad arbor. — *As.* : Japon.

*crispo-ithyphylla* C. M. in *Nuov. Giorn. bot. ital.* 1896, p. 105.

Syn. — *As.* : China centr. (prov. Schen-Si).

*Cryptopodium* C. M. — *Cryptopodium bartramioides*.

*curvirostra* Mitt. — *Conostomum*.

*cuspidatissima* C. M. — Breutelia.

*cygnea* Mont. — *Bartramidula*.

*declivium* C. M. — Breutelia.

*defoliata* C. M. in *Linn.* 1864, p. 597.

Ster. — *Am. merid.* : And. Nov. Granat.

*dicranacea* C. M. — Breutelia.

*dicranoides* Wils. — *Philonotis longicolla*.

*didymocarpa* W. P. Sch. — *Anacolia*.

*diminutiva* C. M. in *Engler's Bot. Jahrb.* 1883, p. 79 et in *Forschungsreise...* p. 20 (1889).

Syn. — *Afr.* : Ins. Kerguelen.

*divaricata* } C. M. — Breutelia Mülleri.

                  } Hook. — Breutelia.

*dumosa* Mitt. — Breutelia.

*Elliotii* Broth. in *Engler's Bot. Jahrb.* 1897, p. 26.

Ster. — *Afr.* Reg. lacustr.

*elongata* Mitt. — *Prionodon*.

*erecta* Mitt. — *Bartramidula*.

*exigua* Sull. — *Bartramidula*.

*falcata* Hook. — *Philonotis fontana* var. *falcata*.

*faucium* C. M. in *Bull. hb. Boissier*, 1898, p. 47 et in *Ule Coll. M. Brasil.* n. 1821.

2? In fauc. rup. — *Am. merid.* : Brasil. or. (S. do Itatiaia).

*Fendleri* C. M. — *Bartramidula*.

*filiformis* Hsch. — *Philonotis tenella* var. *gracillima*.

*flavicans* Mitt. in *Lond. Journ. of bot.* 1851, p. 55 et *M. austr. am* p. 270 (1869); *C. M. Syn.* II, p. 618 (1851).

*B. bogotensis* Hpe. in *Linn.* 1863, p. 142 et in *Ann. sc. nat. ser. V, III*, p. 374 (1866).

Syn. — *Afr.* : Ins. Kerguelen (??). *Am. merid.* : And. Nov. Granat.

*fontana* Schwaegr. — { *Philonotis*.  
                                  *Philonotis rigida*.  
                                  *Philonotis tenella* var. *humilis*.  
var. *falcata* Mont. — *Philonotis macrocarpa* ?  
var. *marchica* H. et T. — *Philonotis marchica* ?

*fragilifolia* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 40.  
— *Am. merid.* : And. Boliv.

*fragilis* Mitt. — *B. papillata*.

*fuscescens* Aongstr. in *Oefv. af K. Vet.-Akad. Foerh. Stockholm*, n. 4, p. 18 (1872).  
2. — *Am. merid.* : Brasil. austr. or.

*gigantea* } Bory. — *Breutelia vulcanica*.  
              } H. f. et W. — *Breutelia divaricata*.

*glauca* Ltz. *Moosst.* p. 160 (1864).  
Fl. et fr. ign. — *Am. sept.* : Mexico.

*glauco-viridis* C. M. et Kindb. in *Mac. Cat.* p. 105 (1892).  
1. Ad rup. humid. — *Am. sept.* : Columb. brit.

*globosa* C. M. — *Bartramidula Breuteliana*.

*gnaphalea* C. M. — *Breutelia*.

*gonioclada* Wils. — *Philonotis fontana* var. *falcata*.

*gracilis* Floörke. — B. Oederi.

var. *tomentosa* C. M. — B. Oederi var. *condensata*.

*graminicola* C. M. — { *Philonotis*.  
                                  *Philonotis kerguelensis*.

*granatensis* W. P. Sch. — *Anacolia Webbii*.

*grandiflora* Schwaegr. — B. Oederi.

*grandis* Geh. et Hpe. — *Breutelia*.

*Griffithsiana* Wils. } *Philonotis*.  
                                  *Philonotis imbricatula*.

*hakonensis* Besch. in *Journ. de bot.* 1898, p.  
— *As.* : Japon.

*Halleriana* (Hedw.) Hedw. *Descript.* II, p. III, t. 40 (1789); *Brid.* *Bryol. univ.* II, p. 33 (1827); *Bryol. eur. fasc.* 12, *Monog.* p. 14, t. 302 (1842); *C. M. Syn.* I, p. 495 (1849); W. P. Sch. *Syn.* II ed. p. 511 (1876).

*B. lateralis* Clairville (1811).

*B. norvegica* Lindb. in *Oefv. af Vet.-Akad. Foerh.* XX, n. 7, p. 389 (1863).

*Bryum alpinum* Wulf. in *Berlin. Schrift.* Vol. III, fasc. I, p. 156 (sec. Ehrh. *Beitr.* VI, p. 42).

Br. *Bartramia* Gmel. *Syst. nat.* II, p. 1335 (1791).

Br. foliis mollibus subulatis, setis brevissimis alaribus, capsulis ovatis *Haller Hist. St. Helvet.* n. 1802, t. 45, f. 8 (1768).

Br. foliis subulatis capsulis sphaericis striatis in pedunculis brevibus Oeder in *Fl. Dan.* Vol. III, fasc. IX, p. 4, t. 533, f. 3 (1770).

Br. laterale Lightf. *Fl. Scot.* II, p. 727 (1777).

Br. norvegicum Gunn. *Fl. norveg.* II, p. 138, n. 1064 (1772).

Br. pendulum Oeder in *Fl. Dan.* V, t. 823, n. 1 (1789).



- Br. pomiforme var. 1 Scop. *Fl. Carn. No.* 1307 (1760).  
 Br. pomiforme  $\beta$  Halleri Vill. *Pl. Dauph. III*, p. 878 (1786).  
 Br. recurvum Wulf. in Jacq. *Collect. II*, p. 224 (1788).  
 Mnium laterale Hoffm. *Deutschl. Fl. II*, p. 54 (1796).  
 Sphagnum subulatum viridissimum, capsulis ovatis Dill. *Hist. M.* p. 245, t. XXXII, f. 3 (1741).  
 Webera clandestinia Hedw. *Fund. II*, p. 104, t. 6, f. 30 (1782).  
 W. Halleriana ej. in op. cit. p. 95 (1782).  
 1. Ad rup. humid. umbr. — Eur. : Zon. sept. et med., rar. merid., reg. mont. usq. in alp. As. : China centr. (prov. Schen-Si), Yunnan, Sikkim, Himalaya, Kashmir, Tibet, Caucas. Afr. : M. Cameroen (?). Am. sept. : M. Rupestr. Pac. : Nov. Seeland., Tasmano. ?

*Halleriana* { H. f. et W. — B. Mossmanniana.  
 { Mitt. — B. magellanica.

Hampeana C. M. in *Bot. Zeit.* 1858, p. 162.

Glyphocarpus asperrimus Hpe. in Jaeg. *Ad. I*, p. 512 (1872-73).

2. — Afr. : Pr. B. Sp.

*Hasskarliana* Hpe. — Breutelia.

*hawaica* C. M. — Bartramidula.

? *heleniana* Bursh. — Philonotis fontana.

*Henoni* Duby. — B. pomiformis var. crispa.

*hercynica* Floerke. — B. pomiformis var. crispa.

*heteromalla* Brid. — B. pomiformis var. heteromalla.

*hispida* Mitt. — Breutelia.

*Hookeri* Wils. — Philonotis longicollis.

*humilis* Mitt. — Glyphocarpus.

*incana* Tayl. — Breutelia.

*inclinata* Hpe. et Ltz. — Breutelia.

inconspicua Mitt. in *Journ. of the Linn. Soc.* 1877, p. 62.

Ster. — Afr. : Ins. Tristan da Cunha.

*incurva* Hoppe. — B. pomiformis var. crispa.

*inserta* Sull. et Lesq. — Glyphocarpus.

*integrifolia* { Sull. — Breutelia aureola ?

{ Tayl. — Breutelia.

*intermedia* Hpe. — Breutelia.

*intertexta* C. M. p. p. — { Anacolia.  
 { Anacolia setifolia.

ithyphylla (Haller) Brid. *Muscol. rec. II*, P. III, p. 132, t. I, f. 6 (1803)  
 et *Bryol. univ. II*, p. 43 (1826); *Bryol. eur. fasc. 12, Mon. p. II*,  
 t. 317 (1842); C. M. *Syn. I*, p. 493 (1849); W. P. Sch. *Syn. II ed.*  
 p. 510 (1876).

B. pomiformis (L. Fl. succ.) Wahlenb. *Fl. lapp.* p. 362 (1812).

Bryum alpinum viridissimum et majus capitulis crassissimis Haller *Enum.*  
*St. Helv.* p. 109, t. 3, f. 4 (1740) et *Hist. Stirp. Helv. No* 1803, p. 44  
 var.  $\gamma$ , t. 46, f. 7 (1768).

Br. pomiforme  $\beta$  majus Web. *Fl. Gott.* p. 120 (1778).

Syn. Ad terram et in fiss. rup. — Eur. : E reg. campestr. usq. in  
 supraalp. praepr. zonae sept. : Spitzberg, S. d'Estrella et Nevada.  
 As. : Kamschatka, Jenissei inf., Kashmir, Caucas. Am. sept. : Tota  
 arctica et temperata.

\* var. *arenae* Besch. in *Miss. sc. Cap. Horn. Bot.* V, p. 288 (1889).

— Am. merid. : Fret. Magellan.

- *var. breviseta* (Lindb.) Husn. *Muscol. gall.* p. 266 (1890).  
*B. breviseta* Lindb. in *Not. ur Saellsk p. Fauna et Flora fenn. Foerh. IX.*  
p. 255 (1868).  
*Ut typ.* — *Eur.* : Finmark, Alp. centr. *Am. sept.* : Groenland, Behring.  
*var. breviseta* Kindb. — *B. ithyphylla var. strigosa.*  
*var. rigidula* W. P. Sch. — *B. ithyphylla var. strigosa.*
- *var. strigosa* Wahlenb. *Fl. lapp.* p. 362 (1812).  
*B. ithyphylla var. breviseta* Kindb. *Arten d. Laubm. Schwed. et Norv*  
p. 153 (1883).  
*B. ithyphylla var. rigidula* W. P. Sch. *Syn. II ed.* p. 510 (1876).  
*B. rigidula* Kindb. *Eur. a. N. Amer. Bryin.* p. 323 (1897).  
*Ut typ.* — *Eur.* : Finmark, Alp. Norveg. *Am. sept.* : Groenland, fret.  
Scoresby.  
*var. subulata* Husn. — *B. subulata.*
- ithyphylloides* W. P. Sch. in *Lechler M. Chil. No. 875; C. M. in Bot.*  
*Zeit.* 1859, p. 229.  
— *Am. sept.* : Mexic. (*fide Besch. Prodr.* p. 57).  
*Am. merid.* : Chile.
- Jacquenotiana* W. P. Sch. in *Lechler Pl. Magell. n.* 1025.  
— *Am. merid.* : Fret. Magell.
- jamaicensis* Mitt. — *Breutelia.*
- Jamesoni* Tayl. — *Cryptopodium Hookeri.*
- javanica* Dz. et Mk. in *Zoll. Syst. Verz.* pp. 24 et 50 ( ) et *Bryol. jav.*  
*I.* p. 154, t. 123 (1855-61).  
2. — *As.* : Simla, Java.
- Jungneri* Par. *Suppl. Ind.* p. 35 (1900).  
*B. aristifolia* Broth. in *Engler's Bot. Jahrb.* 1897, p. 249.  
*Ster.* Ad rupes vulcan. — *Afr.* : Cameroon.
- Karsteniana* C. M. — *Breutelia* ?  
*kilima N'Djarica* C. M. — *Breutelia.*
- Krauseana* C. M. in *Bot. Centralbl.* 1883, No. 2-4.  
*Syn.* — *As.* : Penins. Tschutschica.
- Krausei* } *Bryol. eur.* — *Glyphocarpus compactus.*  
} C. M. — *Bartramidula.*
- lacte-virens* Rehm. — *Breutelia afro-scoparia.*
- laevisphaera* Tayl. — *Glyphocarpus.*
- lamprocarpa* W. P. Sch. — *Anacolia intertexta.*
- lanceolata* Schrad. — *B. Ederi.*
- lateralis* Clairv. — *B. Halleriana.*
- Lechleri* W. P. Sch. in *Lechl. Pl. Magell. n.* 1170.  
— *Am. merid.* : Fret. Magellan.
- Leikiipiae* C. M. in *Fl.* 1890, p. 479.  
*Ster.* — *Afr.* : M. Aberdare.
- leptodonta* Wils. in *Kew Journ. bot.* IX, p. 369 (1857); *Mitt. M. Ind.*  
*or.* p. 58 (1859).  
1 vel syn. — *As.* : N. W. Himalaya (Dehra Doun.)
- leucolomacea* C. M. *Bryol. Austr. Georg.* p. 28 (1890).  
2. In argill. — *Am. merid.* : Austr. Georg.

*Lindigii* Mitt. — Bartramidula.

*lineata* C. M. in Linn. 1878-79, p. 464.

2? — *Am. merid.* : Venezuela.

*longifolia* Hook. M. ex. t. 68 (1818-20); *Brid. Bryol. univ.* II, p. 39 (1827); C. M. Syn. I, p. 496 (1849); *Mitt. M. austr. am.* p. 273 (1869).

Ad rup. granit. — *Am. merid.* : And Nov. Granat.

*longifolia* } Hsch. — H. Martiana.  
                  } Mitt. — B. angustifolia.

*longiseta* } Brid. — B. Oederi.  
                  } Mich. — Philonotis radicalis.

*Lorentzi* C. M. — Breutelia.

*lutescens* Mitt. — Cryptopodium.

*Lycopodium* C. M. in Helms M. Nov. Zealand. n. 67.

— *Pac.* : Nov. Seeland.

*Mac Owaniana* C. M. in Hedwig. 1899, p. 95.

Ster. Ad scaturig. — *Afr.* : Pr. B. Sp.

*macrocarpa* Hpe. — Breutelia tomentosa.

*macrosubulata* C. M. in Bot. Centralbl. 1883, No. 2-4.

Syn. — *As.* : Penins. Tschutschica.

*macrotheca* Hpe. — Breutelia tomentosa.

*magellanica* Aongstr. in Oefv. af K. Vet.-Akad. Foerh. Stockholm, 1872. No. 4, p. 8.

B. Halleriana Mitt. M. austr. am. p. 272 (1869).

l. Ad rup. madid. — *Am. merid.* : Argentin., fret. Magellan., Fuegia.

*magellanica* Mitt. — Conostomum.

*marchica* Sull. — Philonotis Muhlbergii.

*marginalis* Rehm. M. Afr. austr. n. 202.

Ster. — *Afr.* : Orange.

*Martiana* Hpe. Enum. M. Brasil. p. 22, nom. (1879).

B. longifolia Hsch. Fl. brasil. I, p. 40 (1840).

— *Am. merid.* : Brasil. austr. or.

*Matthewsii* Mitt. M. austr. am. p. 273 (1869).

— *Am. merid.* : And. Quitens., Peruv.

*mauiensis* C. M. in Fl. 1896, p. 449.

Ster. — *Pac.* : Ins. Hawai.

*Menziesii* } Lesq. et Jam. — Anacolia Baueri p.p.

                  } Turn. — Anacolia.

*microbasis* C. M. in Linn. 1880-82, p. 409.

Fl. et fr. ign. — *Am. merid.* : Argentin. Cordob.

*microdonta* Mitt. — Glyphocarpus.

*microstoma* Mitt. M. austr. am. p. 272 (1869).

— *Am. sept.* : Guatemala.

*mollis* var. Wils. — Philonotis leptocarpa.

*nniocarpa* C. M. } — Breutelia macrocarpa.  
                          } — Breutelia.

*Mossmanniana* C. M. in *Bot. Zeit.* 1851, p. 552.

B. *Halleriana* H. f. et W. *Fl. of the N. Zeal.* II, p. 88 (1855) et *Handb.* p. 446 (1867) (*jur. fide Mitt. !*).

1. Arb. et rupic. — *Pac.*: Austral. or., N. Seeland., Tasman.

*Mülleri* Mont. — B. *brevifolia*.

*nana* C. M. — *Bartramidula*.

*nevadensis* C. M. — B. *thrausta*.

*nitida* Wils. } — *Philonotis*.  
                   } — *Philonotis Turneriana*.

*Normanni* Holmgr. — B. *pomiformis* var. *heteromalla*.

*norvegica* Lindb. — B. *Halleriana*.

Æderi (Gunn.) *Schwaegr. Suppl. I, P. II, p. 49, t. 59* (1816); *Brid. Bryol. univ. II, p. 46* (1827); *Bryol. eur. fasc. 12, Mon. p. 12, t. 318* (1842); *W. P. Sch. Syn. II ed. p. 512* (1876).

B. *alpina* Schleich. *Cent. IV, No. 17*.

B. *gracilis* Floerke in *Schrad. Journ. f. Bot. II, p. 171* (1799); C. M. *Syn. I, p. 508* (1849); *Drumm. M. bor. am. I ed. No. 21*.

B. *grandiflora* *Schwaegr. Suppl. I, P. II, p. 48, t. 58* (1816).

B. *lanceolata* *Schrad.*

B. *longiseta* *Brid. Musc. rec. II, P. III, p. 136, t. 2, f. 10* (1803).

B. Æderi γ *longiseta* *Brid. Bryol. univ. II, p. 47* (1827).

B. Æderiana Sw. in *Schrad. Bot. Journ. II, p. 181, t. 3 B, f. 5* (1800).

B. *pomiformis* *Sull. M. Allegh. n. 121* (1846).

B. *subintegrifolia* P. Beauv. *Prodr. p. 44* (1805).

*Bryum* foliis striatis acuminatis reflexis, capsulis reflexis subnutantibus, maturitate striatis Æder *Fl. Dan. III, fasc. 8, p. 9, t. 478* (1769).

Br. *lacerum* *Vill. Pl. Dauph. III, p. 879* (1786).

Br. Æderi *Gunn. Fl. norv. No. 1005* (1772).

Br. *pomiforme* β Æderi *Vill. op. et l. cc. p. 879*.

*Mnium gracile* *Funch. Fasc. crypt. IV, p. 93*.

*Plagiopus* Æderi *Limpr. Laubm. Deutschl. etc. p. 548* (1895).

*Syn.* Ad rup. humid. praepr. calcar. — *Eur.*: Reg. mont. et subalp. As.: Penins. Tschutschica, Jenissei med.; Amour, Japon, China centr., Nepal, Himalaya, Kashmir, Tchitral, Gilgit, Pandjab. Am. sept.: Canada, de Nov. Angl. ad M. Rupestr., Columb. brit.

var. *alpina* *Schwaegr.* — B. Æderi var. *condensata*.

« var. *condensata* *Brid. Sp. M. III, p. 87* (1817) et *Bryol. univ. II, p. 48* (1827).

B. *gracilis* β *tomentosa* C. M. *Syn. I, p. 509*.

B. Æderi β *alpina* *Schwaegr. Suppl. I, P. II, p. 48* (1816).

B. Æderi var. *microcarpa* *Kindb. Enum. Bryin. Dove. p. 28* (1888).

B. Æderi var. *subnivalis* *Mol. Alg. Moosst. p. 82* (1865).

*Plagiopus* Æderi var. *condensata* *Limpr. Laubm. Deutschl. etc. p. 550* (1895).

Pl. *serratus* *Brid. Bryol. univ. I, p. 596* (1826); C. M. *Syn. I, p. 511* (1849) (1).

In uligin. (*Brid.*); ad rup. apricas (*Limpr.*) — *Eur.*: Alp. Algov., Stir., Carinth., Tirol., Rhaet., Helvet. As.: Japon, Kashmir, Tchitral, Gilgit.

« var. *glacialis* *Mol. in Un. itin. 1863, No. 20*.

Rupic. — *Eur.*: Alp. Tirol.

(1) Cf. C. M. et *Limpr.* II. cc.

α var. *groenlandica* C. Jens. in *Saërtryk af Medd. om Groenland*, 1898, p. 395.

— *Am. sept.* : Fret. Scoresby.

var. *longiseta* Brid. — B. Oederi.

var. *microcarpa* Kindb. — B. Oederi var. *condensata*.

α var. *minor* Kindb. in *Mac. Cat.* p. 105 (1892).

Ad rup. madid. : *Am. sept.* : M. Rupestr., Columb. brit.

var. *subnivalis* Mol. — B. Oederi.

*Oederiana* Sw. — B. Oederi.

*Oreadella* C. M. *Bryol. austr. georg.* p. 29.

Syn. Rupic. — *Am. merid.* : Austr. Georg.

*papillata* H. f. et W. *Fl. of the N. Zeal.* p. 89, t. 86, f. 4 (1855) et *Handb.* p. 447 (1867); *Fl. Tasm. II*, p. 195 (1860).

B. *acerosa* Hpe. (fid. Mitt.).

B. *fragilis* Mitt. in *Journ. of the Linn. Soc. IV*, p. 81 ( ) et *Fl. Tasm.* p. 196, t. 174, f. 8 (1860).

2.

— *Pac.* : N. Seeland., Austral., Tasman.

*patens* Brid. *Sp. M. III*, p. 82 (1817) et *Bryol. univ. II*, p. 38 (1827) *excl. syn.* *Schwaegr. et Turn.*; C. M. *Syn. I*, p. 494 (1849); Mitt. *M. austr. am.* p. 275 (1869).

B. *reticulata* P. B. *Prodr.* p. 44 (1805).

Syn.

— *Afr.* : Ins. Kerguelen. *Am. merid.* : Terr.

Magell., ins. Maclov., Eremitae.

*patens* { Hook. — B. *brevifolia*.  
Schwaegr. { (fid. C. M.) — B. *vulcanica*.  
(fid. Mitt.) — B. *commutata*.

*patula* Mitt. — *Bartramidula*.

*pendula* { Hook. — B. *eutelia*.  
Mitt. (1876). — *Glyphocarpus chrysurus*.  
Spreng. — *Breutelia Sieberi*.  
Sull. — *Breutelia dumosa* ?

*penicillata* C. M. in *Hedvig*, 1899, p. 84.

B. *Somerseti* Par. *Suppl. Ind.* p. 36 (1900).

Ster. Ad scaturig. — *Afr.* : Pr. B. Sp.

*penicillatula* C. M. in *Linn.* 1880-82, p. 411.

Fl. et fr. ign. — *Am. merid.* : Alp. Argent. sub trop. Tucuman.

*pentasticha* Brid. — *Conostomum australe*.

*perpumila* C. M. in *Linn.* 1880-82, p. 411.

2. — *Am. merid.* : Alp. Argent. sub trop. Tucuman.

*perpusilla* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 41.

2. — *Am. merid.* : And. Boliv.

*pinnata* Hpe. — *Breutelia*.

*polygastrica* C. M. in *Linn.* 1874, p. 600.

Ster. — *Am. merid.* : Summ. Alp. Peruv.

*polytrichoides* C. M. in *Linn.* 1874, p. 597.

Ster. — *Am. merid.* : And. Nov. Granat

*pomiformis* (L. ex. p.) *Hedw. Sp. M.* p. 164 (1801); *Roth Tent. III*, p. 258 (1800); *Brid. Bryol. univ. II*, p. 39 (1827); *Bryol. eur. fasc. 12* (Vol. IV), *Mon.* p. 13 (1842); *C. M. Syn. I*, p. 499 (1849); *W. P. Sch. Syn. II ed.* p. 511 (1876).

*B. crispa*  $\beta$  *minor* *Sw. mss.*, *W. et M. Bot. Taschb.* p. 272 (1807).

*B. crispa* var. *nigrescens* *Kindb. Art. Laubm. Schwed. und Norw.* p. 153 (1883).

*B. crispa* var. *pomiformis* *Lindb. M. scand.* p. 15 (1879).

*B. vulgaris* *Lam. et DC. Fl. fr. I*, p. 509 (1805).

*Bryum capillaceum*, capsulis sphaericis *Dill. Hist. M.* p. 339, t. 44, f. I (1741).

*Br. pomiforme* *L. Sp. pl.* p. 1580, n. 3 sed non *Herbar.*

*Br. trichodes virescens*, erectis majusculis capitulis maliformibus *Dill. Cat. Giss.* p. 224 (1718).

*Mnium pomiforme* *L. jun. Meth. M.* p. 364 (1787).

*Webera pomiformis* *Hedw. Fund. M. II*, p. 95 (1802).

1. Ad terr. lapid. et ad rup. praepr. aren. et granit. — *Eur.*: Reg. camp., et mont. *As.*: Kamschatka, Amour, Japon, China or., Himalaya. *Caucas. Am. sept.*: Reg. arct. et bor. totae. *Am. merid.*: Fret Macell., *Fuegia. Pac.*: N. Seeland.

« var. *crispa* *Bryol. eur.*, *C. M. et W. P. Sch. II. cc.*

*B. crispa* *Sw. Disp. M. Succ.* p. 73 (1799); *Brid. Bryol. univ. II*, p. 41 (1827); *Lindb. M. scand.* p. 15 excl. var.  $\beta$  et  $\gamma$  (1879).

*B. crispa*  $\alpha$  *major* *W. et M. Bot. Taschb.* p. 173 (1807).

*B. Henoni* *Duby in Mem. Soc. Geneve* 1876, II, 2, f. 2 (*B. japonica* in tab.).

*B. hercynica* *Floerke in Schrad. Journ. Bot. II*, p. 171 (1799).

*B. incurva* *Ipe. in Sturm Deutsch. Fl. II, Heft. 6* (1803).

*B. pomiformis*  $\beta$  *elongata* *Turn. Ann. bot. I*, p. 1526 ( ).

*B. pomiformis*  $\alpha$  *major* *H. et T. Muscol. brit.* p. 85 (1818).

*Bryum montanum* *Smith. in Hb. Linn.*

*Locis humidior.* — *Eur.*: Reg. mont. super. vel in zona frig. *As.*: Unalaschka, *Jenissei* super., Amour, Japon., alp. *Yunnan.*, *Kashmir*, *Caucas. Am. sept.*: *Groenland*: Washington (1).

var. *elongata* *Turn.* — *B. pomiformis* var. *crispa*.

« var. *heteromalla* (*Brid.*) *C. M. I. c.*

*B. crispa*  $\beta$  *Normanni* *Holmgr. mss.*; *Hartm. Skand. Fl. X ed. P. II*, p. 47 (1871).

*B. heteromalla* *Brid. Bryol. univ. II*, p. 42 (1827).

*B. Normanni* *Holmgr. mss.*; *Hartm. op. et l. cc.*

Ut typ. — *Eur.*: *Scandin.* *Am. sept.*: Sine loco spec.

var. *major* *H. et T.* — *B. pomiformis* var. *crispa*.

*pomiformis* } *Sull.* — *B. Oederi*.  
                  } *Sw.* — *B. ithyphylla*.

*potosica* *Mont. in Ann. sc. nat.* 1838, p. 56, in *d'Orbigny Voy.* p. 95 (1839) et *Syll.* p. 29 (1856); *C. M. Syn. I*, p. 494 (1849).

*Arb. et rupic.* — *Am. merid.*: *And. Nov. Granat.*, *Quitens.* et *Boliv.*, *Chile*.

*potosica* *C. M. et Hpe.* — *B. brevifolia*.

*Prabaktiana* *Dz. et M'k.* — *Philonotis secunda* var. *Prabaktiana*.

(1) In *Mac. Cat. Canad. Pl.* non. *indicata*.

- praetenuis* Wils. — *Philonotis angusta*.  
*pseudo-Cryptopodium* C. M. in *Linn.* 1880-82, p. 412.  
 Fl. et fr. ign. — *Am. merid.* : Alp. Argent. sub trop. Tucuman.
- pusilla* } C. M. — *Bartramidula*.  
               } Hook. — *Bartramidula exigua*.
- pyncocoleos* C. M. *Bryol. austr. georg.* p. 28 (1890).  
 Ster. Rupic. — *Am. merid.* : Austr. Georg.
- quadrata* Mitt. — *Glyphocarpus marionensis*.  
*radicalis* var. H. f. et W. — *Philonotis pallida*.
- radiculosa* Mitt. in *Journ. of the Linn. Soc.* 1877, p. 62.  
 Ster. In arenos. — *Afr.* : Ins. Tristan da Cunha.
- ramentosa* C. M. in *Hedwig.* 1899, p. 94.  
 Ster. — *Afr.* : Transvaal, pr. B. Sp.
- Readeriana* Col. — *Breutelia*.  
*recurvifolia* Duby. — *Campylopus penicillatus*.  
*remotifolia* H. f. et W. — *Philonotis appressa*.  
*reticulata* P. B. — *B. patens*.
- Rietmanniana* Hpe. in *Jaeg. Ad. I*, p. 534 (1873-74),  
 — *Pac.* : Nov. Hebrid.
- rigida* Sull. — *Philonotis Sullivantii*.
- rigidissima* W. P. Sch. in *Mandon Pl. Boliv.* n. 1674.  
 — *Am. merid.* : And. Boliv.
- rigidula* Kindb. — *B. ithyphylla* var. *strigosa*.  
*rivalis* C. M. — *Breutelia*.
- robusta* H. f. et W. *Fl. antarct.* p. 153, t. 59, f. 4 (1844-47) et *Handb.*  
*of the N. Zeal. Fl.* p. 447 (1867); C. M. *Syn. I*, p. 495 (1849) et *II*,  
 p. 619 (1851).  
 2. Coll. humid. — *Afr.* : Ins. Heard, Kerguelen. *Pac.* : Ins. Auckland,  
 Campbell.
- robusta* Sull. et Lesq. — *B. Wilkesiana*.  
*Roylei* C. M. — *Bartramidula*.
- rufescens* Hpe. *Symb.* 1874, p. 491.  
 2. — *Am. merid.* : Brasil. austr. or.
- rupestris* Mitt. — *Breutelia*.
- rupta* C. M. in *Linn.* 1880-82, p. 413.  
 Fl. et fr. ign. — *Am. merid.* : Argent. Cordob.
- Rusbyana* C. M. — *Breutelia thrausta*.
- ruvenzorensis* Broth. in *Engler's Bot. Jahrb.* 1877, p. 249.  
 Syn. — *Afr.* : Reg. lacustr.
- scabrida* Schwaegr. — *Philonotis uncinata*.
- Schimperi* C. M. *Syn. II*, p. 617 (1851).  
*B. brevifolia* W. P. Sch. in *Hb. Mus. Paris*.  
 Syn. — *Am. sept.* : Mexic.
- Schmidiana* C. M. in *Bot. Zeit.* 1858, p. 162.  
 Syn. — *As.* : M. Neilgherr.
- scoparia* Schwaegr. p.p. } *Breutelia*.  
                                       } *Breutelia Sieberi*.

- scorpioides* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 47.  
Ster. — *Am. merid.* : And. Boliv.
- secunda* W. P. Sch. in *Mandon Pl. Boliv. No.* 1671; C. M. in *op. et l. cc. (descript.)*  
— *Am. merid.* : And. Boliv.
- secundifolia* C. M. — Breutelia.
- sericea* Hsch. — Glyphocarpus.
- setifolia* Mitt. pp. } Anacolia.  
                                  } Anacolia intertexta.
- Sieberi* Hsch. — Breutelia.
- Somerseti* Par. — B. penicillatula.
- Spielhausi* C. M. — Breutelia.
- squarrosa* } Mitt. — Breutelia.  
                  } Turn. — B. vulcanica?
- stricta* Brid. M. rec. II, P. III, p. 132, t. I, f. 5 (1803) et *Bryol. univ.* II, p. 45 (1827); *Bryol. eur. fasc.* 12, Mon. p. 10 (Vol. IV), t. 316 (1842); C. M. Syn. I, p. 500 (1849); W. P. Sch. Syn. II ed. p. 509 (1876).  
B. strictifolia Tayl. in *Lond. Journ. of Bot.* 1846, p. 51.  
Syn. Rupic. et terr. — *Eur.* : Zon. merid. et medit.; *Britann.*? *As.* : Minor; Unalascika? *Afr.* : Alger., Canar., Madeir., Cameroon.  
*Am. sept.* : Colorado, Calif. *Am. merid.* : Fret. Magell.? *Pac.* : Austral. occ. (Swan River), Tasman.
- stricta* Schwaegr. — Glyphocarpus compactus.
- strictae-affinis* C. M. in *Dusen M. Camer. N.* 337.  
— *Afr.* : Cameroon.
- strictifolia* Tayl. — B. stricta.
- strictula* C. M. in *Fl.* 1888, p. 412.  
Ster. — *Afr.* : Kilima N'Djaro.
- strumosa* Mitt. — Glyphocarpus.
- subarcuata* C. M. — Breutelia.
- subasperrima* C. M. in *Rehm. M. Afr. austr. n.* 199, 213 et in *Hedwig.* 1899, p. 95.  
Ster. — *Afr.* : Pr. B. Sp.
- subbrevifolia* C. M. in *Linn* 1880-82, p. 464.  
2? — *Am. merid.* : Venezuela.
- subdisticha* Hpe. — Breutelia.
- subexigua* C. M. — Bartramidula.
- subgnaphalea* C. M. — Breutelia.
- subintegrifolia* P. B. — B. Oederi.
- subithyphylla* Besch. *Prodr. bryol. mexic.* p. 58 (1871).  
Ster. — *Am. sept.* : Mexic.
- submicrocarpa* C. M. — Breutelia.
- subpatens* Hpe. in *Ann. sc. nat. V ser. III*, p. 375 (1866).  
Syn. — *Am. merid.* : And. Nov. Granat.
- subpatens* C. M. — B. austro-georgica.
- subpellucida* Mitt. M. Ind. or. p. 59 (1859).  
Syn. — *As.* : Kashmir, Himalayn.
- subsessilis* C. M. — Glyphocarpus levisphaerus.



- substricta* W. P. Sch. mss.; C. M. in Bot. Zeit. 1858, p. 162.  
2. — Afr. : Pr. B. Sp.
- subtomentosa* Hpe. — Breutelia.
- subulata* Bryol. eur. fasc. 31, Suppl. p. 1 (1846); C. M. Syn. I, p. 493 (1849); W. P. Sch. Syn. II ed. p. 508 (1876).  
B. ithyphylla var. subulata Husn. Musc. Gall. p. 266 (1890).  
B. viridissima Kindb. Eur. and N. Am. Bryin. p. 323 (1897).  
Weisia viridissima Brid. Bryol. univ. I, p. 364 (1826).  
Syn. In fiss. rup. — Eur. : Summ. Alp. Salisb., Carinth., Rhaet., Engadin. As. : Kashmir, Baltistan, Sikkim. Am. sept. : Groenland, Alaska, Sitka, Colorado.
- tenella* Wils. — Philonotis imbricatula.  
var. Wils. — Philonotis heterophylla.  
var. erecta C. M. — Philonotis glaucescens.
- tenuifolia* Mitt. — Breutelia.
- thelioides* C. M. in Fl. 1875, N. 34-35.  
Ster. — Am. merid. : And. Nov. Granat.
- thrausta* W. P. Sch. in Mandon Pl. Boliv. N. 1673; C. M. in Nuov. Giorn. bot. ital. 1897, p. 41 (descr.)  
B. nevadensis C. M. in Linn. 1880-82, p. 410.  
B. Rusbyana ej. in Herb. Nov. Eborac.  
2. — Am. merid. : And. Boliv., Alp. Argent  
subtrop.
- ticinensis* Kindb. in Rev. bryol. 1892, p. 102.  
Ad rupes schistos. — Eur. : Ital. super. (Lugano).
- tomentosa* { C. M. Breutelia.  
Schwaegr. — Breutelia gnaphalea.
- Trianae* Hpe. — Breutelia.
- Türkheimii* C. M. — Bartramidula.
- Uleana* C. M. in Ule Bryoth. bras. n. 14.  
— Am. merid. : Brasil. austr. or.
- Ulei* C. M. — Breutelia.
- uncinata* { C. M. et W. — Philonotis Gardneri.  
Wils. — Philonotis lutea.
- vaginans* Rehm. M. Afr. austr. N. 201 et 534.  
Ster. — Afr. : Orange.
- viridissima* { C. M. — Cryptopodium Hookeri.  
Kindb. — B. subulata.
- vulcanica* Brid. Bryol. univ. II, p. 37 (1827); C. M. Syn. I, p. 497 (1849).  
B. gigantea Bory Voy. en. Afr. III, p. 96 ( ).  
B. patens Schwaegr. Suppl. I, P. II, p. 55, t. 62 (1816).  
B. squarrosa Turn. in König et Sims. Ann. bot. I, p. 329, t. 2, f. 2? (1805-6).  
Syn. Rup. basalt. — Afr. : Ins. Borboniae.
- vulgaris* Lam. et DC. — B. pomiformis.
- Webbii* C. M. — Anacolia.
- Wilkesiana* Par. Index p. 115 (1893).  
B. robusta Sull. in Wilkes Expl. Exped. p. 12 ( ); C. M. in Fl. 1885, p. 413.  
— Am. merid. : Fuegia.

*Willii* C. M. *Bryol. austr. georg.* p. 30 (1890).

Ster. Rupic. — *Am. merid.* : Austr. Georg.

*Wilsoni* C. M. — *Bartramidula*.

*xanthochacta* W. P. Sch. in *Mandon Pl. Boliv. N.* 1672.

— *Am. merid.* : And. Boliv.

**BARTRAMIDULA** *Bryol. eur. fasc. 29-30* (1846). — 18 spec.

*Breuteliana* W. P. Sch. in *Breutel M. Capens.*

*Bartramia globosa* C. M. in *Hedwig*. 1899, p. 90.

*Glyphocarpus Breutelii* *Jaeg. Ad. I*, p. 521 (1873-74).

Syn. — *Afr.* : Pr. B. Sp.

« *var. tenuicaulis* (C. M.) *Par. Suppl. Ind.* p. 36 (1900).

*Bartramia globosa var. tenuicaulis* C. M. in *op. et l. cc.*

*Bartramidula capensis* W. P. Sch. *hb. (fid. C. M.)*.

*Glyphocarpus Schimperii* *Hpe. hb. (fid. C. M.)*.

Ut et cum typ.

*capensis* W. P. Sch. — *B. Breuteliana var. tenuicaulis*.

*cernua* Lindb. — *B. Wilsoni*.

*curta* (*Hpe*) *Par. Ind.* p. 115 (1893).

*Glyphocarpus curtus* *Hpe. Enum. M. Brasil.* p. 21 (1879).

*Philonotula curta* *ej. Symb.* 1874, p. 722.

Syn. — *Am. merid.* : Brasil. austr. or.

*cygnaea* (*Mont.*) *Jaeg. Ad. II*, p. 698 (1877-78).

*Bartramia cygnaea* *Mont. in Ann. sc. nat.* 1845, IV, p. 102, *Hist. Chil. Bot. crypt.* t. 2, f. 2 (1850) *et Syll.* p. 28 (1856); C. M. *Syn. I*, p. 479 (1849).

*Glyphocarpus cygnaeus* *Jaeg. Ad. I*, p. 523 (1873-74).

2. *Paludic.* — *Am. merid.* : Chile.

*erecta* (*Mitt.*) *Jaeg. Ad. II*, p. 698 (1877-78).

*Bartramia erecta* *Mitt. M. austr. am.* p. 255 (1869).

*Glyphocarpus erectus* *Jaeg. Ad. I*, p. 523 (1873-74).

1. *Terr.* — *Am. merid.* : And. Quitens.

*exigua* (*Sull.*) *Jaeg. Ad. II*, p. 698 (1877-78).

*Bartramia exigua* *Sull. in U. St. Expl. Exped.* p. II, t. 8 ( ).

*B. pusilla* *Hook. Journ. bot.* 1856, p. 316.

*Glyphocarpus exiguus* *Jaeg. Ad. I*, p. 522 (1873-74).

— *Am. merid.* : Pr. Horn.

*Fendleri* (C. M.) *Par. Ind.* p. 115 (1893).

*Bartramia Fendleri* C. M. in *Linn.* 1878-79, p. 403.

Syn. — *Am. merid.* : Venezuela.

*Glaziovii* (*Hpe.*) *Jaeg. Ad. II*, p. 698 (1877-78).

*Glyphocarpus Glaziovii* *Hpe. Symb.* 1872, p. 18.

2. — *Am. merid.* : Brasil. austr. or.

*hawaica* (C. M.) *Par. Suppl. Ind.* p. 36 (1900).

*Bartramia hawaica* C. M. in *Fl.* 1896, p. 447.

Syn. ? — *Pac.* : Ins. Hawaï.

*Krausei* (C. M.) *Jaeg. Ad. II*, p. 698 (1877-78).

*Bartramia Krausei* C. M. in *Linn.* 1874, p. 595.

*Glyphocarpus Krausei* *Jaeg. Ad. I*, p. 523 (1873-74).

2. — *Am. merid.* : Chile.

*Lindigii* (Hpe.) *Jaeg. Ad. II*, p. 698 (1877-78).

*Bartramia Lindigii* Mitt. *M. austr. am.* p. 255 (1869).

*Glyphocarpus Lindigii* Hpe. in *Linn.* 1861-62, p. 522.

Syn. Terr. — *Am. merid.* : And. Nov. Granat.

*mexicana* W. P. Sch. in *Besch. Prodr. bryol. mexic.* p. 58 (1871).

*Glyphocarpus mexicanus* *Jaeg. Ad. I*, p. 522 (1873-74).

Syn. — *Am. sept.* : Mexic. (Orizaba).

*nana* (C. M.) *Par. Suppl. Ind.* p. 36 (1900).

*Bartramia nana* C. M. in *Fl.* 1896, p. 447.

Syn. ? — *Am. merid.* : And. Venezuel.

*patula* (Mitt.) *Jaeg. Ad. II*, p. 698 (1877-78).

*Bartramia patula* Mitt. *op. cit.* p. 255.

*Glyphocarpus patulus* *Jaeg. Ad. I*, p. 522 (1873-74).

Syn. Terr. — *Am. sept.* : Guatemala.

*pusilla* (Hook. f. et Wils.) *Jaeg. Ad. II*, p. 698 (1877-78).

*Bartramia pusilla* C. M. *Syn. I*, p. 480 (1849) et *II*, p. 616 (1851).

*Glyphocarpus pusillus* H. f. et W. in *Lond. Journ. of bot.* 1844, p. 515 et *Fl. Tasm. II*, t. 174, f. 2 (1860).

1. — *Pac.* : Tasman.

*Roylei* *Bryol. eur. fasc.* 12, *Mon.* p. 2 (1842).

*Bartramia Roylei* C. M. *Syn. I*, p. 478 (1849).

*Glyphocarpa Roylei* Hook. f. in *Hook. Ic. pl. rar. II*, t. 194 (1841) et in *Lond. Journ. of bot.* 1841, *II*, p. 8.

Syn. — *As.* : Himalaya, M. Neilgherr.

*subexigua* (C. M.) *Par. Ind.* p. 116 (1893).

*Bartramia subexigua* C. M. in *Engler's Bot. Jahrb.* 1883, p. 79 et in *Forschungsreise.....* p. 21 (1889).

*Philonotis appressa* Mitt. in *Journ. of the Linn. Soc.* 1876, p.

2. — *Afr.* : Ins. Kerguelen.

*Türckheimii* (C. M.) *Par. Suppl. Ind.* p. 36 (1900).

*Bartramia Türckheimii* C. M. in *Bull. hb. Boissier*, 1897, p. 187.

Syn. Rupic. — *Am. sept.* : Guatemala.

*Wilsoni* *Bryol. eur. fasc.* 29-30, t. 327 (1846); W. P. Sch. *Syn. II ed.* p. 516 (1876).

*Bartramia* (Philonotula) *Wilsoni* C. M. *Syn. I*, p. 479 (1849).

*Bartramidula cernua* Lindb. in *Oefv. Vet.-Akad. Föerh.* 1863, n. 7, p. 7.

*Glyphocarpus* ? *cernuus* Wils. in *Hook. Journ. of bot.* 1841, p. 8, *memorata*.

Gl. *Wilsoni* *Jaeg. Ad. I*, p. 521 (1873-74).

Syn. Ad terr. humos. — *Eur.* : Hibern., Scot., Cambrovall. *As.* : Alp. Yunnan. *Afr.* : Ins. Fernando Po. *Am. sept.* : New Jersey ?

**BARTRAMIOPSIS** Kindb. in Rev. bryol. 1894, p. 35. — 1 spec.

*Lescurii* (Jam.) Kindb. in op. et l. cc.

*Atrichum Lescurii* Jam. in Bull. Torr. bot. Club. VI, p. 33 (1879);  
Lesq. et Jam. Man. p. 257 (1884); Par. Ind. p. 53 (1894).

*Oligotrichum Lescurii* Mitt. in Trans. of the Linn. Soc. 1891, p. 191.

*Philocrya aspera* Hag. u. Jens. in Meddel. om Groenland, XV, 1898,  
p. 388 c. ic.

2. — As. : Japon. Am. sept. : Alaska, Sitka, fret.  
Scoreshy.

» *subsp. sitkana* Kindb. in op. et l. cc.

— Am. sept. : Sitka.

**BECCARIA** C. M. in Nuov. Giorn. bot. ital. 1872, p. 11. — 2 spec.

*elatio* C. M. in op. et l. cc.

1 vel ps. 2. Terr. — Afr. : Abyss.

*minor* C. M. in op. et l. cc.

1 vel ps. 2. Ad rup. madid. — Afr. : Abyss.

**BECKETTIA** C. M. in Hedwig. 1898, p. 77.

*bruchioides* C. M. — Henediella.

**BERGIA** Fühnr. in Bot. Zeit. 1829, Ergänz. bl. p. 25.

(*Absque nom. spec.*) — Entosthodon Bergianus.

**BESCHERELLIA** Duby in Bull. Soc. bot. Fr. 1873, p. 30, c. ic. — 2 spec.

*brevifolia* Hpe. in Linn. 1876, p. 317.

2. — Pac. : Austral. or. (Queensland).

*elegantissima* Duby l. c. c. ic.

1. Cortic. — Pac. : Nov. Caledon.

**BLINDIA** Bryol. eur. fasc. 33-36 (1846). — 25 spec.

Spec. omn. non indic. sunt DICRANOWEISIA ejusd. nom.

*acuta* (Huds.) Bryol. eur. l. c. Mon. p. 3, t. 114; C. M. Syn. I, p. 342  
(1849) et II, p. 584 (1851); W. P. Sch. Syn. II ed., p. 131 (1876).

*Bryum acutum* Huds. Fl. angl. II ed., p. 481 (1778).

Br. filiforme Vill. Pl. Dauph. III, p. 875 (1786).

Br. pilosum sphagni subulati facie Dill. Hist. M. p. 374, t. 47, f. 34 (1741).

Br. splachnoides Dicks. Fusc. II Pl. crypt. p. 5 (1790).

Br. verticillatum Lightf. Pl. Scot. II, p. 733 (1777).

Br. verticillatum  $\beta$  Huds. Fl. angl. p. 411.

Dicranum splachnoides Brid. Musc. rec. II, P. I, p. 181 (1812).

Grimmia acuta Turn. Muscol. hibern. p. 29 (1804).

Gr. rupincola W. et M. Reite in Schwed. t. 2, f. 3 a-d (1804).

Gr. schisti Engl. Bot. t. 1958.

Gymnostomum subulatum Bryol. germ. I, p. 147, t. 10, f. 12 (1823).

Seligeria acuta de Not. Epil. p. 653 (1869).

Weisia acuta Hedw. Descr. III, p. 85, t. 35 (1792); Brid. Bryol. univ. I,

p. 362 (1826).

Weisia acuta  $\beta$  rupestris Brid. op. cit. p. 363.

Weisia acuta  $\gamma$  rupincola ej. l. c.

W. fastigiata Bryol. germ. II, t. 35, f. 31 (1831).

W. rupestris Hedw. Sp. M. t. 14 (1801).

2. Ad rup. granit. vel schist., nunquam calc. — *Eur.* : Island., Spitzberg ; e subalp. usque in supraalp. per tot. zon. interm. et septentr. *As.* : Caucas. *Am. sept.* : Groenland, Alaska, M. Rupestr., Columb. brit., ins. Vancouver, Oregon, Terra nova, M. Alb., Catskill, New York. *Am. merid.* : Ins. Eremitæ ? *Pac.* : Tasman.
- var. *arenacea* Mol. — *Bl. acuta* var. Seligeri.  
var. *curviseta* Mitt. — *Bl. curviseta*.
- \* var. *flexipes* R. C. in *Rev. bryol.* 1892, p. 79.  
Ut typ. — *Am. sept.* : Alaska, Oregon.
- \* var. *irrorata* Pfeff. *Bryogeogr. Stud.* p. 29 (1869); *Jaeg. Ad. I.* p. 179 (1871-72.)  
Ut. typ. — *Eur.* : Engadin.
- \* var. Seligeri (*Brid.*) *Limpr. Laubm. Deutschl. etc., I.* p. 475 (189C).  
*Bl. acuta* var. *arenacea* Mol. *Moosst. Algaü* p. 65 (1865).  
*Bl. acuta* var. *trichodes* Braithw. in *Journ. bot.* 1870, p. 228.  
*Bl. trichodes* Lindb. *mss.* ; *Philib. in Rev. bryol.* 1881, p. 90 ; *Par. Ind.* p. 119 (1893).  
*Dicranum rupestre* W. et M. *Bot. Taschb.* pp. 185 et 469 (1807).  
D. Seligeri *Brid. Mant. M.* p. 59 (1819) et *Bryol. univ. I.* n. 129 (1826).  
D. *trichodes* Wils. *M. brit. n.* 77.  
Ut typ. — *Eur.* : Britann., Riesengeb., Siles.
- var. *trichodes* Braithw. — *Bl. acuta* var. Seligeri.  
*acuta* H. f. et W. — *Bl. curviseta*.
- arcuata* Mitt. *M. austr. am.* p. 55 (1869).  
Ad rup. madid. — *Am. merid.* : Fuegia, ins. Eremitæ. *Pac.* : Tasman.
- aschistodontoides* C. M. in *Engler's Bot. Jahrb.* 1883, p. 78 et in *Forschungreise.... Bot.* p. 18 (1889).  
Ster. — *Afr.* : Ins. Kerguelen.
- \* var. *nigrescens* C. M. in *Forschungreise.... I. c.*  
*Cynodontium Hookeri* Mitt. in *Journ. of the Linn. Soc.* 1876 ?  
Ut et ubi typ.
- auriculata* C. M. in *Fl.* 1885, p. 405.  
l. *Rupic.* — *Am. merid.* : M. Fuegiae.
- brevipes* C. M. *Bryol. austr. georg.* p. 24 (1890).  
l. *Rupic.* — *Am. merid.* : Austro-Georg.
- caespiticia* Lindb. -- *Stylostegium*.
- chrysea* C. M. in *Trans. and Proceed. of the N. Zeal. Inst.* 1892, p. 290.  
l. *Rupic.* — *Pac.* : Nov. Seeland.
- churuccana* Besch. in *Bull. Soc. bot. Fr.* 1885, p. LVIII et in *Miss. sc. Cap Horn, V, Bot.* p. 266 (1889).  
l. — *Am. merid.* : Terr. Magell.
- contacta* (*Hook. f. et Wils.*) C. M. *Syn. I.* p. 344 (1849) et *II.* p. 584 (1851).  
*Dicranoweisia contacta* *Par. Ind.* p. 340 (1895).  
*Weisia contacta* H. f. et W. in *Journ. of bot.* 1844, p. 540, *Fl. antarct. I.* p. 127, t. 58 (1860) et *Handb. of the N. Zeal. Fl.* p. 405 (1867).  
2 ? — *Afr.* : Ins. Kerguelen. *Am. merid.* : Fret. Magell. *Pac.* : Ins. Campbell.

*curviseta* Mitt. *M. austr. am.* p. 56 (1869).

Bl. *acuta* H. f. et W. *Fl. Antarct. l. c.* p. 8.

Bl. *acuta* var. *curviseta* Mitt. in *Journ. of the Linn. Soc.* 1859, p. 68.

2. Ad rup. madid. — *Am. merid.* : And. Quitens., Chile, ins. Eremit., Maclov., Fuegia. *Pac.* : Tasman.

*curviseta* Mitt. p.p. — Bl. *gracillima*.

*dicranellacea* C. M. *Bryol. austr. georg.* p. 26.

Ster. *Rupic.* — *Am. merid.* : Austr. Georg.

*dryptodontoides* C. M. — *Dicranoweisia*.

*ferruginea* (Hook. f. et Wils.) Broth. *Natür. Pflanzenfam.* p. 306.

*Dichodontium ferrugineum* Jaeg. *Ad. II*, p. 635 (1877-78); *Par. Ind.* p. 322 (1894).

*Dicranum ferrugineum* H. f. et W. *Fl. Tasm.* p. 171, t. 172, f. I (1860).

2. Terr. — *Pac.* : Tasman.

*flexipes* Kindb. — Bl. *acuta* var. *flexipes*.

*gracillima* Mitt. in *Journ. of the Linn. Soc.* 1879, (*extra vol.*), p. 26.

Bl. *curviseta* ej. in op. cit. XV, p. 198 ( ).

In lacunis. — *Afr.* : Ins. kerguelen.

*leptotrichocarpa* C. M. in op. cit. p. 405.

2. Paludic. — *Am. merid.* : M. Fuegiae.

\* var. *strictiuscula* C. M. in op. et l. cc.

Ut et ubi typ.

*lygopodina* C. M. in op. cit. p. 406.

2. Ad rup. marit. — *Am. merid.* : Fuegia.

*magellanica* W. P. Sch. mss.; C. M. in *Bot. Zeit.* 1862, p. 328.

— *Am. merid.* : Ins. Eremitae.

*microcarpa* Mitt. — *Dicranoweisia Kerguelensis*.

*pulvinata* C. M. in Engler's *Bot. Jahrb.* 1883, p. 79 et in *Forschungsreise.... Bot.* p. 19 ( ).

*Cynodontium conicum* Mitt. in *Journ. of the Linn. Soc.* 1876?

Ster.

— *Afr.* : Ins. Kerguelen.

*robusta* Hpe. in Linn. 1859-60, p. 627; F. v. Müll. *Anal. draw. Austr. M. t. VII* (1864).

1. In saxos. glacial. — *Pac.* : Alp. Austral. or., Nov. Seeland., Tasman.

*Savatieriana* C. M. in Miss. sc. *Cap Horn*, V, *Bot.* p. 267 (1889).

1.

— *Am. merid.* : Terr. Magellan.

*seligerioides* Lindb. in Broth. *Enum. M. Caucas.* p. 64 (1892).

2. Ad rup. trachyt. — *As.* : Caucas. centr.

*Sonsoniae* C. M. in Linn. 1874, p. 593.

Ster.

— *Am. merid.* : And. Nov. Granat.

*sordida* (Wils.) C. M. in *Bot. Zeit.* 1864, p. 349.

*Campylopus sordidus* Jaeg. *Ad. I*, p. 138 (1869-70).

*Dicranum sordidum* Wils. in *Kew Journ. bot.* IX, p. 297 (1857); Mitt. *M. Ind. or.* p. 18 (1859).

— *As.* : Nepal. or.

*stricta* (Hook. f. et Wils.) C. M. Syn. I, p. 343 (1849), II, p. 585 (1851) et in *Forschungsreise...* Bot. p. 17 (descript. locupletiss.) (1889).

*Leptotrichum strictum* H. f. et W. Fl. Tasman. II, p. 177 (1860).

*Weisia stricta* H. f. et W. in Lond. Journ. of bot. 1844, p. 510 et Fl. antarct. II, p. 401, t. 152, f. 4 (1847).

2. — Afr. : Ins. Kerguelen.

*stylostegium* C. M. — *Stylostegium caespiticium*.

*Taylori* Hpe. — *Syrrhæpodon*.

*tenuifolia* (Hook. f. et Wils.) Mitt. M. austr. am. p. 59 (1869).

*Dicranum tenuifolium* H. f. et W. in Lond. Journ. of bot. 1844, p. 512 et Fl. antarct. II, p. 407, t. 152, f. 7 (1847); C. M. Syn. I, p. 365 (1849) et II, p. 592 (1851).

— Am. merid. : Fuegia, ins. Eremitæ. Pac. : Austral. or. (Victoria), Nov. Seeland.

*tortelloides* C. M. in Engler's Bot. Jahrb. 1883, p. 78 et in *Forschungsreise...* Bot. p. 18 (1889).

*Dicranoweisia tortelloides* Par. Ind. p. 312 (1895).

Ster. — Afr. : Ins. Kerguelen.

*tortifolia* (Hook. f. et Wils.) C. M. Syn. I, p. 343 (1849) et II, p. 584 (1851).

*Dicranoweisia tortifolia* Par. op. et l. cc.

*Weisia tortifolia* H. f. et W. in Journ. of bot. 1844, p. 510 et Fl. antarct. II, p. 404, t. 152, f. 5 (1847).

2. Afr. : Ins. Kerguelen.

*trichodes* Lindb. — *Bl. acuta* var. Seligeri.

**Blindiadelphus** Lindb. M. Scand. p. 25 (1879) (sect. SELIGERIÆ).

**BRACHELYMA** W. P. Sch. Syn. II ed. p. 557 (1876). — 1 spec.

*subulatum* (P. B.) W. P. Sch. op. et l. cc. ; Cardot Mon. p. 132 (1892).

*Cryphaea inundata* Nees in Pflanz. Maxim. v. Wied., p. 27 ( ).

*Dichelyma subulatum* Myr. in Act. reg. Acad. sc. Holm. 1832, t. 7 b, f. 18-11 ; Bryol. eur. fasc. , Mon. p. 9, t. 434 ( ).

*Fontinalis subulata* P. B. Prodr. p. 158 (1805) ; Brid. Bryol. univ. II, p. 661 (1827).

*Neckera subulata* C. M. Syn. II, p. 145 (1851).

2. Ad rup. et rad. arb. submers. — Am. sept. : Illinois, Arkansas., Georg., Ludovic.

**BRACHYDONTIUM** (Bruch. hb.) Furrn. in Fl. 1827, II, Reil. I, p. 37. — 2 spec.

*flexisetum* Hpe. in Linn. 1863, p. 132.

2. Ad saxa ripar. — Am. merid. : And. Nov. Granat.

*trichodes* (Web. fl.) Bruch hb. ; Furrn. in op. et l. cc.

*Anoetangium trichodes* Schwægr. Suppl. I, P. I, p. 33, t. 10 (1811).

*Brachyodon trichodes* Furrn. in Fl. 1827, II, Beil. p. 112, Fl. 1829, II, p. 594 et Ergänzbl. II, p. 59.

*Brachyodus trichodes* Bryol. germ. II, P. II, p. 5 (1831) ; Bryol. eur. fasc. 33-36, t. 115 (1816) ; C. M. Syn. I, p. 416 (1849) ; W. P. Sch. Syn. II ed. p. 132 (1876).

*Grimmia trichodes* Web. fl. in W. et M. Bot. Ind. M. pl. cr. 3 (1803) et Archiv. Naturg. I, P. I, p. 124, 4, f. 1 a-d (1804).

*Weisia trichodes* H. et T. Muscol. brit. p. 45, t. 15 (1818).

1. Ad rup. aren., granit. et vulcan., numquam calcar. — *Eur.* : Reg. mont. zonae sept. et centr., copiose in Voges. et Silv. nigra, Siles., Saxon, etc. ; Gall. centr., Pyren., Etruria. *As.* : Caucas. *Am. sept.* : Rarius. : New Hampshire.

**BRACHYMENIUM** Hook. M. exot. III (1818-20). — 129 spec.

Species omn. non indicat. sunt ACIDODONTIA ejusd. nom.

abyssinicum *Br. et W. P. Sch. in W. Sch. M. Abyss. II coll. N. 434.*

*Bryum abyssinicum C. M. Syn. I, p. 265 (1849).*

*Syn.* — *Afr.* : Alp. Abyss.

acuminatum *Harv. in Hook. Ic. pl. rar. t. 19, f. 3 (1841) et in Lond. Journ. of bot. 1840, II, p. 10.*

*Br. cuspidatum Griff. Not. p. 442 et Ic. pl. asiat. t. 102, f. I (1849).*

*Bryum Harveyanum C. M. Syn. I, p. 313 (1849); Mitt. M. Ind. or. p. 68 (1859).*

angolense (*Welw. et Duby*) *Jaeg. Ad. I, p. 579 (1873-74).*

*Bryum angolense W. et Dy. in Mem. .... Genève, 1870, p. 5, t. I, f. 4.*

*Ad arb.* — *Afr.* : Angola.

angustatum *W. P. Sch. in Besch. Prodr. bryol. mexic. p. 51 (1871).*

— *Am. sept.* : Mexic.

appressifolium *Ren. et Card. in Bull. Soc. roy. bot. Belg. 1895, p. 61.*

*Ster.* — *As.* : Sikkim.

atro-purpureum *W. P. Sch. in Mandon Fl. Boliv. N. 1661.*

— *Am. merid.* : And. Boliv.

Barbae-Montis *C. M. in Bull. Soc. roy. bot. Belg. 1892, I, p. 165.*

— *Am. sept.* : Costa Rica.

barbuloides (*C. M.*) *Par. Suppl. Ind. p. 37 (1900).*

*Bryum barbuloides C. M. in Nuov. Giorn. bot. ital. 1897, p. 21.*

2. — *Am. merid.* : And. Boliv.

Baroni *Mitt. in Wright Journ. (1888).*

— *Afr.* : Madagascar.

Bordazii *Ren. et Card. in Bull. Soc. roy. bot. Belg. 1890, I, p. 178.*

*Ster. Muric.* — *Am. sept.* : Martinic.

Borgenianum *Hpe. in Linn. 1874, p. 211.*

2. — *Afr.* : Usambara, Madagascar, ins. Franciae.

brachypelma *C. M. in Bull. Soc. roy. bot. Belg. 1892, I, p. 162.*

2. — *Am. sept.* : Costa-Rica.

Breutelii (*C. M.*) *Par.* — Haplodontium.

brevicaule (*Hpe.*) *Jaeg. Ad. I, p. 582 (1873-74).*

*Bryum brevicaule Hpe. in C. M. Syn. I, p. 323 (1849).*

*Br. nepalense Dz. Mk. M. frond. ined. Archip. ind. p. 12 ? (1851).*

1. — *As.* : Java.

brevipes *Broth. in Hedwig. 1894, p. 131.*

1. *Cortic.* — *Am. merid.* : Brasil. austr. or. (ins. S. Catharina).

bryoides *Hook.* — *Br. Weisia.*



capillare W. P. Sch. *ms. et in Besch. Prodr. bryol. mexic. p. 50* (1871).

Br. capillifolium C. M. *Syn. II, p. 578* (1851).

2. — *Am sept. : Mexic.*

\* var. longicollis Besch. *op. et l. cc.*

Ut et ubi typ.

capitulatum (Mitt.) Par. *Ind. p. 120* (1893).

Bryum capitulatum Mitt. in *Journ. of the Linn. Soc.* 1886, p. 306

1. Ad *Erythrinam*. — *Afr. : Cameroon, Usagara, Kilima N'Djaro, Madagascar.*

cellulare (Hook.) Jaeg. — Bryum.

cernum W. P. Sch. — Bryum japonense.

ciliare Mont. — Br. leptostomoides.

clavariaeforme (C. M.) Jaeg. *Ad. I, p. 581* (1873-74).

Bryum clavariaeforme C. M. in *Bot. Zeit.* 1853, p. 24.

2. — *As. : M. Neilgherries.*

clavigerum Mitt. — Br. clavulum.

clavulum Mitt. in *Trans. of the Linn. Soc.* 1891, p. 165.

Br. clavigerum ej. in *Bisset's M. Japon. n. 207* (in *Herb. British Mus.* .

1. — *As. : Japon.*

coarctatum (C. M.) *Bryol. jav. I, p. 140, t. 115* (1855-61).

Bryum coarctatum C. M. *Syn. I, p. 312* (1849).

2. Calcic. — *As. : Java, Pac. : Ins. Samoa, Tonga, Nov. Seeland., Nov. Caledon.*

columbicum (de Not.) *Broth. Natur. Pflanzenfam. p. 557.*

Br. globosum Jaeg. *Ad. I, p. 574* (1874-75); *Par. Ind. p. 132* (1894).

Bryum columbicum E. S. Salmon. *in litt.*

Br. globosum Mitt. *M. austr. am. p. 289* (1869).

Osculatia columbica de Not. in *Mem. Acad. sc. Torino*, 1859, p. 445, t. VI; *Jaeg. op. cit. p. 516; Par. op. cit. p. 990* (1896).

2. — *Am. merid. : Columb. super., And. Quitens.*

commutatum (C. M.) Jaeg. *Ad. I, p. 575* (1873-74).

Bryum commutatum C. M. in *Bot. Zeit.* 1858, p. 156.

Br. dicranoides W. P. Sch. in *W. Sch. M. Abyss. N. A. 10* (*fid. C. M.*).

2. — *Afr. : Abyss.*

consimile (Mitt.) Jaeg. *Ad. I, p. 582* (1873-74).

Bryum consimile Mitt. *M. austr. am. p. 281* (1869).

2. — *Am. merid. : And. Quitens.*

crinitum Jaeg. — Br. systylium.

cristatum (C. M.) Jaeg. *op. cit. p. 583.*

Bryum cristatum C. M. *Syn. II, p. 580* (1851).

2. — *As. : Bombay.*

cuspidatum Griff. — Br. acuminatum.

daguense (Besch.) Par. *Suppl. Ind. p. 38* (1900).

Peromnium daguense Besch. in *Bull. hb. Boissier*, 1897, p. 332.

2. Cortic. — *Am. merid. : Columb.*

*densifolium* (Aongstr.) Broth. op. cit. p. 558.

*Mielichhoferia densifolia* Aongstr. in Oefv. af K. Vet.-Akad. Foerh. 1873, n. 5, p. 139; Par. Ind. p. 815 (1896).

2. — Afr. : Ins. Franciae.

*dicranoides* (Hsch.) Jaeg. op. cit. p. 575.

*Bryum dicranoides* Hsch. in Linn. 1841, p. 131; C. M. Syn. I, p. 309 (1849).

2. Muric. et rupic. — Afr. : Pr. B. Sp.

*didymodontium* (Mitt.) Jaeg. op. cit. p. 574 (1).

*Bryum didymodontium* Mitt. M. austr. am. p. 289 (1869).

2. — Am. sept. : Mexic.

*Duseni* (C. M.) Par. op. et l. cc.

*Bryum Duseni* C. M. in Dusen M. Camer. n. 772; Par. Ind. p. 178 (1893).

— Afr. : M. Cameroon.

*ellipticum* (Mitt.) Jaeg. op. cit. p. 575.

*Bryum ellipticum* Mitt. M. austr. am. p. 289 (1869).

2. — Am. merid. : Chile.

*eriophorum* Rehm. M. Afr. austr. n. 545.

Ster. — Afr. : Transvaal.

*erubescens* (C. M.) Par. Ind. p. 121 (1893).

*Bryum erubescens* C. M. in Linn. 1880-82, p. 379.

Fl. et fr. ign. — Am. merid. : Argent. tropic.

*eurychaelium* Besch. Fl. bryol. Réunion, etc. p. 90 (1880-81).

1 ? Terr. — Afr. : Ins. Bourbon.

*exile* (Dz. Mh.) Bryol. jav. I, p. 139 (1855-61).

*Bryum exile* Dz. Mh. in Ann. sc. nat. 1840, p. 300 et M. frond. ined Archip. ind., p. 13, t. 5 (1854); C. M. Syn. I, p. 341 (1849).

2. As. : Java, Sumatra.

*extenuatum* (Mitt.) Jaeg. op. cit. p. 578.

*Bryum extenuatum* Mitt. M. Ind. or. p. 68. (1859).

Syn. — As. : Carnatic.

*fabronioides* (C. M.) Par. op. cit. p. 122.

*Bryum fabronioides* C. M. in Linn. 1878-79, p. 289.

2. — Am. merid. : Argentin. subtropic.

*filiforme* } Griff. — *Anomobryum cymbifolium*.  
} Wils. p.p. — *Anomobryum auratum*.

*flaccidisetum* (C. M.) Jaeg. op. cit. p. 576.

*Bryum flaccidisetum* C. M. in Bot. Zeit. 1853, p. 23.

2. — As. : M. Neilgherr.

*flexifolium* Br. et Sch. in W. Sch. M. Abyss. n. 452.

*Bryum flexifolium* C. M. Syn. I, p. 266 (1849).

2. — fr. : Abyss., Cameroon.

---

(1) *Pohlia* *fid.* Broth. Natur. Pflanzenfam. p. 552).

- fragile* (C. M.) Jaeg. *Ad. II*, p. 703 (1877-78).  
*Bryum fragile* C. M. in *Linn.* 1865, p. 378.  
 Ster. *Ad Filices*. — *Afr.* : Niam-Niam.
- fusiferum* (Mitt.) Jaeg. *Ad. I*, p. 576 (1873-74) (1).  
*Bryum fusiferum* Mitt. *M. austr. am.* p. 290 (1869).  
 2. Terr. — *Am. merid.* : And. Quitens.
- Gilliesii* (Hook.) Jaeg. — *Bryum*.
- glaucum* (C. M.) Jaeg. *op. cit.* p. 582.  
*Bryum glaucum* C. M. in *Linn.* 1869-70, p. 36.  
 1. — *As.* : Ceylan.
- globirameum* (C. M.) Par. *op. et l. cc.*  
*Bryum globirameum* C. M. in *op. cit.* 1878-79, p. 478.  
 2. — *Am. merid.* : Venezuela.
- globosum* (Mitt.) Jaeg. — *Br. columbicum*.
- Heribaudi* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1891, *II*, p. 187;  
*Ren. M. Masc. Mad. n.* 26 (1).  
 2 ? — *Afr.* : Ins. Bourbon, Madagascar.
- Holstii* Broth. in *Engler's Bot. Jahrb.* 1894, p. 189.  
*Ad arb.* — *Afr.* : Usambara.
- Hookeri* Bryol. jav. — *Br. nepalense*.
- Hornschuchianum* Mart. *lc. crypt. Brasil.* XXXVI, t. 20, f. A ( ) et  
*in Fl. Brasil.* I, p. 36 (1840).  
*Bryum Hornschuchianum* C. M. *Syn. I*, p. 322 (1849); *Mitt. op. cit.* p. 281.  
 2. Terr. in silv. — *Am. merid.* : Brasil. austr. or.
- huillense* (Welw. et Duby) Jaeg. *op. cit.* p. 572.  
*Bryum huillense* W. et Dy. in *Mém. . . Genève*, 1870, p. 6, t. I, f. 5.  
 2. *Ad arb.* — *Afr.* : Angola.
- imbricatum* W. P. Sch. *mss. et hb.*  
*Bryum imbricatifolium* C. M. *Syn. II*, p. 578 (1851); *Mitt. op. cit.* p. 282.  
 2. — *Am. sept.* : Mexic.
- indicum* (Dz. Mk.) Bryol. jav. I, p. 141 (1855-61).  
*Bryum indicum* Dz. Mk. *M. frond. ined. Archip. ind.* p. 22, t. II (1854);  
*C. M. Syn. I*, p. 313 (1849).  
*Syn. Calcic.* — *As.* : Amboine, Java.  
 « *var. corrugatum* Besch. *Fl. bryol. Nouv.-Caled.* p. 213 (1873).  
*Saxic.* — *Pac.* : Nov. Caledon.
- Jamesoni* Tayl. in *Lond. Journ. of bot.* 1848, p. 282.  
*Bryum Jamesoni* C. M. *Syn. II*, p. 580 (1851); *Mitt. op. cit.* p. 283.  
*Peromnium Jamesoni* Jaeg. *op. cit.* p. 584.  
 2. *Ad arb.* — *Am. sept.* : Jamaic. *Am. merid.* : Columb., And. Quitens. et Peruv.

---

(1) Pohlia (fd. Broth. *Natur. Pflanzenfam.* p. 552).

*japonense* Besch. — Bryum.

*julaceum* Hsch. in Linn. 1841, p. 133.

Bryum koratranum C. M. Syn. I, p. 324 (1849).

2. Ad arb. — Afr. : Transvaal, Pr. B. Sp.

*Klotzschii* (Schevaegr.) Par. Ind. p. 123 (1893).

Br. Sellowianum Jaeg. op. cit. p. 578.

Bryum Klotzschii Mitt. op. cit. p. 283.

Br. Sellowianum C. M. op. cit. p. 319.

Didymodon Klotzschii Schwaegr. Suppl. IV, t. 340 (1832).

2. — Am. merid. : Brasil.

*Krausei* Hpe. et Ltz. in Bot. Zeit. 1869.

Peromnium Krausei Jaeg. op. cit. p. 585.

2. Silvic. — Am. merid. : Aequator.

*lanceolatum* Hook. f. et Wils. Fl. Tasman. II, p. 188, t. 173, f. 8 (1860).

Bryum lanceolatum Mitt. in Trans. and Proceed. of the roy. Soc. of Victoria, 1883, p. 71.

1. — Pac. : Tasman.

*Lefebvrei* Kindb. — Bryum.

*leptophyllum* W. P. Sch. in hb. Hpe ; Jaeg. op. cit. p. 573.

Bryum leptophyllum Br. et W. P. Sch. in W. Sch. M. Abyss. Coll. II, n. 451 et 460 ; C. M. op. cit. p. 273 ; Par. Ind. p. 38.

1. Ad arb. — Afr. : Abyss.

*leucostomum* Bryol. jav. — Webera.

*leucotrichum* C. M. in Linn. 1875, p. 377.

Ster. — Afr. : Niam-Niam.

*longipedicellatum* (C. M.) Par. Ind. p. 123 (1893) (1).

Bryum longipedicellatum C. M. in Linn. 1878-79, p. 477.

2. — Am. merid. : Venezuela.

*longissimum* (C. M.) Par. Suppl. Ind. p. 39 (1900).

Bryum longissimum C. M. in Micholitz M. Philipp. n. 150.

— As. : Ins. Philipp. (Luzon).

*luteolum* (C. M.) Jaeg. Ad. I, p. 579 (1873-74).

Bryum luteolum C. M. in Linn. 1874, p. 625.

2. — Am. sept. : Mexic.

*madagassum* Hpe. in Linn. 1874, p. 242.

2. — Afr. : Madagascar.

*magellanicum* (Sull.) Par. op. et l. cc.

Bryum magellanicum C. M. Syn. II, p. 579 (1851).

Peromnium magellanicum Sull. in Lond. Journ. of bot. 1850, p. 316.

Syn. — Am. merid. : Fuegia.

*malachiticum* (C. M.) Par. op. et l. cc.

Bryum malachiticum C. M. in Linn. 1880-83, p. 378.

2. Cortic. — Am. merid. : Argent. subtropic. Tucuman.

---

(1) Pohlia (fid. Broth. Natur. Pflanzenfam. p. 552.

- mammillosum (C. M.) *Par. Suppl. Ind. p. 39 (1900) (1).*  
*Bryum mammillosum C. M. in Bull. hb. Boissier, 1897, p. 550.*  
 2. — *Am. sept.* : Jamaic.
- megalacrimon (Schwaegr.) *Jaeg. Ad. I, p. 578 (1873-74).*  
*Bryum megalacrimon Schwaegr. Suppl. II, P. I, p. 104 (1823) ; C. M. Syn. I, p. 312 (1449).*  
 2. Ad arb. — *Afr.* : Ins. Franciae, Rodriguez, Tristan da Cunha.
- melanopyxis (C. M.) *Par. Suppl. Ind. p. 39 (1900).*  
*Bryum melanopyxis C. M. in Fl. 1896, p. 530.*  
 — *Am. merid.* : And. Venezuel.
- melanothecium (C. M.) *Jaeg. op. cit. p. 576.*  
*Bryum coarctatum Mitt. Sam. M. p. 177 (1867).*  
*Br. melanothecium C. M. M. Polyn. p. 62 (1874).*  
*Syn. Muric. — Pac.* : Ins. Samoa, Tonga, Taïti, archip. Gambier.
- mexicanum *Mont. in Ann. sc. nat. 1838, IX, p. 54 et Syll. p. 34 (1856).*  
*Bryum erectum Hook. in Kunth Syn. Pl. æquinoct. ? (1822).*  
*Br. mexicanum C. M. Syn. I, p. 252 (1849) ; Mitt. M. austr. am. p. 282 (1869).*  
*Br. Webera erecta Brid. Bryol. univ. I, p. 626 (1826).*  
 2. — *Am. sept.* : Mexic.
- Meyenianum (Hpe.) *Jaeg. Ad. I, p. 175 (1871-72).*  
*Bryum Meyenianum C. M. op. cit. p. 296.*  
*Miehlichhoferia chilensis Hpe. in Hb. roy. Berol.*  
*Webera Meyeniana ej. in Linn. 1837, p. 268.*  
 2. — *Am. sept.* : Chile.
- micans (Mitt.) *Jaeg. op. cit. p. 583 (1873-74).*  
*Bryum micans Mitt. M. Ind. or. p. 68 (1859).*  
 2. — *As.* : Ind. or.
- microstomum *Harv. in Lond. Journ. of bot. 1840, II, p. 10 et in Hook. Ic. Pl. rar. t. 19, f. 4 (1841).*  
*Bryum microstomum C. M. op. cit. p. 325.*  
 — *As.* : Nepal.
- miehlichhoferioides C. M. in *Nuov. Giorn. bot. ital. 1872, p. 16.*  
 2 ? — *Afr.* : Abyss.
- minutulum *Hpe. in Bot. Zeit. 1870, p. 51.*  
 2 ? *Rupic.* — *Am. sept.* : Mexic.
- Molleri (C. M.) *Par. Ind. p. 124 (1893).*  
*Bryum Molleri C. M. in Fl. 1886, p. 281.*  
 2. — *Afr.* : Ins. São Thoma.
- morasicum *Besch. in Bull. hb. Boissier, 1894, p. 392.*  
 2. Ad arb. — *Am. merid.* : Columb.
- murale W. P. Sch. in *Besch. Prodr. bryol. mexic. p. 51 (1871).*  
 2. — *Am. sept.* : Mexic.

---

(1) *Pohlia (fd. Broth. Natur. Pflanzenfam. p. 552).*

*Myurella* (C. M.) Broth. *Natür. Pflanzensfam.* p. 558.

*Bryum Myurella* C. M. in Linn. 1875, p. 380; *Par. Ind.* p. 197 (1894).

Ster. Terr. — Afr. : Dar Fertit.

*myurelloides* Par.

Br. Dusenii Broth. *op. et l. co.*

*Bryum Myurella* C. M. in Dusen M. Camer. n. 314.

Br. myurelloides Par. *Ind.* p. 197 (1894).

*Pohlia Myurella* Broth. in Engler's Bot. Jahrb. 1897, p. 244.

2. Rupic. — Afr. : M. Cameroon.

*Neesii* (C. M.) Par. *Suppl. Ind.* p. 39 (1900).

*Bryum Neesii* C. M. in Hedwig. 1899, p. 66.

2? — Afr. : Pr. B. Sp.

*nepalense* Hook. in Schwaegr. *Suppl. II, P. II*, p. 131, t. 135 (1826);

*Brid. Bryol. univ. I*, p. 602 (1826).

Br. Hookeri Bryol. jav. I, p. 138, t. 114 (1856-61).

*Bryum contortum* Griff. *Not.* p. 440 et  *Ic. pl. asiat. II*, t. 100, f. 2 (1849).

Br. Hookeri Spreng. *Syst. Veg. IV*, p. 212 (1827); C. M. *Syn. I*, p. 323 (1849).

Br. nepalense Mitt. *M. Ind. or.* p. 73 (1859).

1. Cortic. et terr. — As. : N. W. Himalaya, Nepal., Mysore; Célebes, Borneo, Java, Sumatra.

*Nietneri* (C. M.) Jaeg. *Ad. I*, p. 576 (1873-74).

*Bryum Nietneri* C. M. in Linn. 1869-70, p. 37.

2. — As. : Ceylan.

*nigrescens* Besch. *Fl. bryol. Réunion, etc.* p. 91.

1. — As. : Madagascar.

*Nordenskioldii* Besch. in Oefv. af Vet.-Akad. Foerh. 1900, p. 291.

1. — As. : Japon.

*orthodontioides* Jaeg. — *Bryum Gilliesii*.

*orthopelma* (C. M.) Jaeg. *Ad. II*, p. 703 (1877-78).

*Bryum orthopelma* C. M. in Linn. 1874, p. 550.

1. — As. : Ins. Philippin.

? *ovatum* H. f. et W. — *Leptotheca Gaudichaudii*.

*pallido-julaceum* (C. M.) Par. *Suppl. Ind.* p. 39 (1900).

*Bryum pallido-julaceum* C. M. in Hedwig. 1899, p. 67.

Ster. — Afr. : Transvaal.

*papillosum* (C. M.) Jaeg. *Ad. II*, p. 122 (1874-75) (1).

*Bryum papillosum* C. M. *Syn. I*, p. 326 (1849); Mitt. *M. austr. am* p. 290 (1869).

2. Saxic. et terr. — Am. sept. : Jamaic. Am. merid. : Venezuela, And. Nov. Granat., Quitens. et Peruv.

*patulum* W. P. Sch. *ms. et hb.*

*Bryum patulum* C. M. *Syn. II*, p. 579 (1851).

1. Ad arb. — Am. sept. : Mexic.

---

(1) *Pohlia* (fid. Broth. *Natür. Pflanzensfam.* p. 552).

- pendulum *Mont. in Ann. sc. nat. XVI*, p. 254 ( ) et *Syll.* p. 34 (1856)  
*Bryum Montagneanum C. M. Syn. I*, p. 225 (1849); *Mitt. M. Ind. or.*  
 p. 69 (1859).
2. Terr. in silv. — *As.* : M. Neilgherr.
- peraristatum (*C. M.*) *Par. Ind.* p. 125 (1893).  
*Bryum peraristatum C. M. in Linn.* 1878-1879, p. 478.
2. — *Am. merid.* : Venezuela.
- Philonotula (*Hpe.*) *Broth. in Engler's Bot. Jahrb.* 1894, p. 191.  
*Bryum Philonotula Hpe. in Hildebr. M. Madag. n.* 2136.  
 — *Afr.* : Usambara, Madagascar.
- piriforme *W. P. Sch. in Jaeg. Ad. I*, p. 577 (1873-74).  
 — *Afr.* : Pr. B. Sp.
- Pittieri *Ren. et Card. in Bull. Soc. roy. bot. Belg.* 1892, I, p. 163.  
 1. — *Am. sept.* : And. Costa-Ric.
- Plumella (*C. M.*) *Par. Suppl. Ind.* p. 39 (1900) (1).  
*Bryum Plumella C. M. in Hedwig.* 1899, p. 67.  
 — *Afr.* : Transvaal.
- polychaete (*C. M.*) *Par. op. et l. cc.*  
*Bryum polychaete C. M. in Dusen M. Camer. n.* 243; *Par. Ind.* p. 26  
 (1893).  
 — *Afr.* : Cameroon.
- Preissianum (*Hpe.*) *Jaeg. Ad. I*, p. 575 (1873-74).  
*Bryum Preissianum Hpe. Ic. M. t.* 25 (1841); *C. M. Syn. I*, p. 310 (1849).  
 2 ? Rupic. — *Pac.* : Austral. or. (Victoria) et occid. ; Tasman.
- procerrimum *Broth. in Engler's Bot. Jahrb.* 1897, p. 243.  
 1. — *Afr.* : Reg. lacustr.
- pulchellum *Hsch. Fl. brasil. fasc. I*, p. 36 (1840).  
*Bryum Martianum C. M. op. cit.* p. 320; *Mitt. M. austr. am.* p. 283 (1869).  
 2 ? Ad arb. — *Am. merid.* : Brasil.
- pulchrum *Hook. Bot. Misc.* 1830, I, p. 136, t. 38.  
*Bryum pulchrum C. M. op. cit.* p. 324.  
 2 ? Ad arb ? — *Afr.* : Reg. lacustr., Transvaal, Pr. B. Sp., Natal, ins.  
 Rodriguez.
- radiale (*C. M.*) *Kindb. Enum.*  
*Bryum radiale C. M. in Hildebrandt M. Mad. n.* 2115.  
 — *Afr.* : Madagascar.
- « var. leptoradiale (*C. M.*) *Par. Suppl. Ind.* p. 39 (1900).  
*Bryum radiale var. lepto-radiale C. M. in coll. cit.*  
 — *Afr.* : Madagascar.
- radiculosum (*Schwaegr.*) *Par. Ind.* p. 125 (1894).  
*Bryum Peromnium C. M. Syn. I*, p. 321 (1849).

---

(1) *Pohlia (fid. Broth. Natur. Pflanzenfam.* p. 552).

- Br. radiculosum *Mitt. op. cit.* p. 283.  
 Peromnium radiculosum *Schwaeagr. Suppl. III, P. I, II, t. 250* (1828).  
 2. Ad arb. — *Am. merid.* : Brasil. austr. or.
- Regnellii *Hpe. in Linn.* 1849, p. 582.  
 Bryum Regnellii *C. M. Syn. II, p. 581* (1851).  
 Streblophilum Regnellii *Aongstr. in Oefv. af K. Vet.-Akad. Föerh.* 1876, n. 4, p. 18.  
 2. Rupic. — *Am. merid.* : Brasil. austr. or.
- revolutum *Broth. in Engler's Bot. Jahrb.* 1894, p. 190.  
 1. Cortic. — *Afr.* : Cameroon.
- rostratum (*C. M.*) *Jaeg. Ad. I, p. 577* (1873-74) (1).  
 Bryum rostratum *C. M. in Bot. Zeit.* 1858, p. 156.  
 Syn. Muric. — *As.* : Malabar.
- rosulatum *Besch.* — Bryum.
- rugosum (*C. M.*) *Jaeg. op. cit.* p. 580.  
 Bryum rugosum *C. M. in Bot. Zeit.* 1853, p. 23; *Mitt. M. Ind. or.* p. 69 (1859).  
 2. — *As.* : M. Neilgherr.
- Salaminae (*C. M.*) *Jaeg. op. cit.* p. 576 (1).  
 Bryum Salaminae *C. M. in Linn.* 1874, p. 580.  
 2. — *Am. merid.* : And. Novo Granat.
- scabridens (*Mitt.*) *Broth. in Hedwig.* 1899, p. 218.  
 Bryum scabridens *Mitt. in Journ. of the Linn. Soc.* 1861, p. 151.  
 Webera brevicolla *Sande Lac. in Miq. Ann. mus. bot. Lugdun. Batav. II, p. 294* ( ).  
 W. scabridens *Jaeg. op. cit.* p. 592; *Par. Ind. p.* 1359 (1893).  
 2. — *As.* : China or., Japon.
- Schenkii *Broth. in Hedwig.* 1894, p. 132.  
 1. Cortic. — *Am. merid.* : Brasil. or.
- Sellowianum *Jaeg.* — Br. Klotzschyi.
- semperlaxum (*C. M.*) *Jaeg. op. cit.* p. 576.  
 Bryum semperlaxum *C. M. in Linn.* 1874, p. 582.  
 Syn. — *Pac.* : Austral. or.
- sikkimense *Ren. et Card. in Bull. Soc. roy. bot. Belg.* 1899, II, p. 220.  
 Ster. — *As.* : Sikkim.
- smaragdinum (*C. M.*) *Jaeg. op. cit.* p. 577.  
 Bryum smaragdinum *C. M. Syn. II, p. 576* (1851); *Mitt. M. austr. am.* p. 290 p.p. (1869).  
 2. — *Am. merid.* : And. Venezuel.
- sordidissimum (*C. M.*) *Par. Ind. p.* 126 (1893).  
 Bryum sordidissimum *C. M. in Linn.* 1878-79, p. 474.  
 2. — *Am. merid.* : Venezuela.
- spathidophyllum *Besch. Fl. bryol. Réunion, etc., p. 90* (1880-81).  
 1. — *Afr.* : Ins. Bourbon.

---

(1) Pohlia (*fid. Broth. Natur. Pflanzenfam.* p. 552).



- spatulifolium* Ren. et Card. in Bull. Soc. roy. bot. Belg. 1892, I, p. 164.  
2. Ad terr. in silv. — Am. sept. : Costa-Rica.
- speirocladum* C. M. in Besch. op. cit. p. 91.  
2. — Afr. : Usambara, Comor. magn.
- spirale* Besch. in Journ. de bot. 1891.  
2. — Am. merid. : Paraguay.
- spirifolium* (C. M.) Jaeg. op. cit. p. 581.  
Bryum spirifolium C. M. in Linn. 1864, p. 623.  
2. Ad arb. rad. — Am. sept. : Mexic.
- splachnoides* Harv. — Bryum.
- stenopyxis* C. M. in Rehm. M. Afr. austr. n. 220.  
Bryum liliputanum C. M. in Hedwig. 1899, p. 66.  
— Afr. : Pr. B. Sp.
- stereoneuron* C. M. in Nuov. Giorn. bot. ital. 1872, p. 17.  
2. — Afr. : Abyss.
- subcapillifolium* (C. M.) Jaeg. op. cit. p. 579.  
Bryum subcapillifolium C. M. in Linn. 1874, p. 624.  
2. Ad arb. — Am. sept. : Mexic. (Orizaba).
- subcinitum* Par. Suppl. Ind. p. 40 (1900).  
Br. rugosum C. M. in Bull. hb. Boissier, 1898, p. 30 et in Ule Coll. M. brasil. n. 1776, 1777.  
2. Ad arb. — Am. merid. : Brasil. or. (S. do Itatiaia).
- suberectum* Mitt. in Journ. of the Linn. Soc. 1863, p. 155.  
1. — Afr. : Ins. Fernando Po.
- subflexifolium* Ren. et Card. in Bull. Soc. roy. bot. Belg. 1891, II, p. 148;  
Ren. M. Masc. Mad. n. 222.  
1. — Afr. : Madagascar.
- subglobosum* W. P. Sch. in Besch. Fl. bryol. Antill. fr. p. 30 (1876) (1).  
Acidodontium subglobosum Jaeg. Ad. II, p. 703 (1877-78).  
2. Ad terr. lapid. — Am. sept. : Guadalup.
- subsmaragdinum* (C. M.) Jaeg. Ad. I, p. 577 (1873-74).  
Bryum smaragdinum Mitt. in Spr. M. And. n. 355 et M. austr. am. p. 290  
(quoad pl. quitensem) (1869).  
Br. subsmaragdinum C. M. in Linn. 1874, p. 581.  
2. — Am. merid. : And. Quitens.
- subuliferum* Mitt. in Journ. of the Linn. Soc. 1863, p. 155.  
1. — Afr. : Ins. Sao Thoma.
- systylium* (C. M.) Jaeg. Ad. II, p. 117 (1874-75).  
Acidodontium Kunthii Hsch. in Deppe et Schiede M. Mexic.  
Br. crinitum Jaeg. Ad. I, p. 581 (1873-74).  
Bryum crinitum Mitt. in Lond. Journ. of bot. III, p. 56 ( ); C. M.  
Syn. II, p. 580 (1851).  
Br. systylium C. M. Syn. I, p. 320 (1849); Mitt. op. cit. p. 280.  
2. — Am. sept. : Mexic. Am. merid. : And. Quitens.

---

(1) Pohlia (fid. Broth. Natur. Pflanzenfam. p. 552).

tenellum W. P. Sch. mss. et hb.

Bryum tenellum C. M. Syn II, p. 572.

2. — Am. sept. : Mexic. (M. Orizaba).

torquescens W. P. Sch. in Q. Dillon et Petit M. Abyss.

— Afr. : Abyss.

velutinum (C. M.) Jaeg. op. cit. p. 583.

Bryum velutinum C. M. in Bot. Zeit. 1853, p. 33; Mitt. M. Ind. or. p. 73. (1859).

2. — As. : M. Neilgherr.

vernicosum (C. M.) Par. Suppl. Ind. p. 40 (1900) (1).

Bryum vernicosum C. M. in Nuov. Giorn. bot. ital. 1897, p. 22.

— Am. merid. : And. Holiv.

Walkeri Broth. in Records of the bot. Survey of India, 1899, p. 320.

2. Ad rup. gneiss. et terr. argill. — As. : M. Ghates (Coorg).

Weisia Harv. in Lond. Journ. of bot. 1840, II, p. 10 et in Hook. Ic. pl. rar. t. 19, f. 1 (1841).

Bryum bryoides Hook. in Schwaeagr. Suppl. III, II, p. 134, t. 135 (1828); Brid. Bryol. univ. I, p. 603 (1826).

Br. exile Dz. Mkh. (fid. Mitt. M. Ind. or. p. 685).

Br. nepalense Spreng. Syst. Veg. IV, p. 211 (1827); C. M. Syn. I, p. 311 (1849).

Br. Weisia C. M. op. cit. p. 325; Mitt. op. et l. cc.

1. Terr. — As. : Himalaya, Nepal., Khasia, M. Neilgherr.

Welwitschii (Duby) Jaeg. Ad. II, p. 119 (1874-75).

Bryum Welwitschii Dy. in Mém.... Genève, 1870, t. 2, f. 3.

2. In spongiosis. — Afr. : Angola.

Wilsoni (Mitt.) Jaeg. — Leptobryum.

Wolkensii Broth. in Engler's Bot. Jahrb. 1894, p. 189.

2. Cortic. ad Erythrinam. — Afr. : Kilima N'Djaro.

Wrightii (Sull.) Broth. op. cit. p. 559.

Leptotheca Wrightii Sull. in Proceed. of the Amer. Acad. ... V, p. 281;

Par. Ind. p. 736 (1896).

Peromnium Wrightii Jaeg. Ad. I, p. 385 (1873-74).

Ps. 1. Lignic. — Am. sept. : Florid., Cuba, ins. S. Vincent.

BRACHYMITRIUM Tayl. in Lond. Journ. of bot. 1846, p. 44.

Jamesoni ) Tayl. — Tayloria.

) W. P. Sch. — Tayloria Mandoni.

BRACHYDODON Fürnr. in Fl. 1827, II, Beil. I, p. 112 et 1829, II, p. 594 et Ergänzbl. p. 59.

trichodes Fürnr. — Brachydontium.

BRACHYDODUS Bryol. germ. II, P. II, p. 3 (1831).

trichodes Bryol. germ. — Brachydontium.

BRACHYPODIUM Brid. Bryol. univ. I, p. 147 (1826).

crispatum Brid. — Ptychomitrium.

parasiticum Brid. — Calymperes.

(1) Pohlia (fid. Broth. Natur. Pflanzenfam. p. 552.

**BRACHYSTELEUM** *Rehb. Comp. p. 34, N. 619* (1828).

Spec. omn. non indicatae sunt **PTYCHOMITRIA** ejusd. nom.

- crispatum* C. M. p.p. } — **Ptychomitrium**.  
                                   } — **Ptychomitrium nigricans**.  
     β *brachycarpum* Hsch. — **Ptychomitrium cucullatifolium**.  
*Drummondii* C. M. — **Glyphomitrium**.  
*Hampeanum* C. M. — **Ptychomitrium incurvum**.  
*polyphyllum* C. M. p.p. } — **Ptychomitrium**.  
                                   } — **Ptychomitrium nigricans**.

**BRACHYTHECIUM** *Bryol. eur. fasc. 52-54* (1853). — 240 spec.

*abyssinicum* *Par. Suppl. Ind. p. 40* (1900).

*Br. trachypodioides* *W. P. Sch. in W. Sch. M. Abyss.*

*acuminatum* (*Hedw.*) *Kindb. in Mac. Cat. p. 191* (1892).

*Camptothecium acuminatum* *Kindb. Eur. and N. Am. Bryin. p. 88* (1897).

*Homalothecium acuminatum* *Jaeg. Ad. II, p. 375* (1876-77).

*Hypnum acuminatum* *P. B. Prodr. p. 60* (1805); *C. M. Syn. II, p. 334* (1851); *Sull. Ic. M. p. 187, t. 116* (1864); *Lesq. et Jam. Man. p. 336* (1884).

*H. bryiforme* *Brid. Musc. rec. II, P. II, p. 100* (1801).

*H. erectum* *Drumm. M. bor. am. I ed., n. 224*.

*H. julaceum, erectum, Bryi argentei habitu* *Dill. Hist. M. p. 322, t. 41 A.B.D.* (1745).

*H. pulchrum* *Drumm. Coll. cit. n. 180 p.p.*

*Isothecium bryiforme* *Brid. Bryol. univ. II, p. 355* (1827).

*Leskea acuminata* *Hedw. Sp. M. p. 227, t. 56, f. 1-7* (1801); *Brid. op. cit. p. 296*.

*L. Beyrichii* *Hpe. in Linn. 1839, p. 47 et Ic. M. t. 7* (1844).

*L. imbricatula* *Hedw. Sp. M. p. 224, t. 2* (*fid. Card. ex autops.*).

2. *Ad rad. arb. et terr. — Am. sept. : Reg. sept., orient., centr. et merid.; Aniniboine.*

« *var. rupincolum* (*Hedw.*) *Ren. et Card. in Rev. bryol. 1893, p. 17.*

*Hypnum acuminatum var. rupincolum* *Lesq. et Jam. op. et l. cc.*

*H. rupincolum* *P. B. op. cit. p. 69.*

*Leskea rupicola* *Hedw. op. cit. p. 54; Brid. Bryol. univ. II, p. 302* (1827).

*Rupic. et terr. — Am. sept. : Cum typ.*

« *var. setosum* (*Hedw.*) *Ren. et Card. in op. et l. cc.*

*Hypnum acuminatum var. setosum* *Lesq. et Jam. op. et l. cc.*

*H. setosum* *P. B. op. et l. cc.*

*Leskea setosa* *Hedw. op. cit. p. 236, t. 57; Brid. op. et l. cc.*

*Lignic. — Am. sept. : Cum typ.*

« *subsp. stenocarpum* *Kindb. in Hedwig. 1896, p. 66.*

*Camptothecium stenocarpum* *ej. Eur. and N. Am. Bryin. p. 89* (1897).

— *Am. sept. : Wisconsin.*

« *var. subalbicans* *Ren. et Card. in Bot. Gaz. 1890, p. 60.*

*Ad arb. — Am. sept. : Florid., Ludovic.*

*acutum* *Sull. — Br. Mildeanum.*

*afro-glareosum* (*Broth.*) *Par. Suppl. Ind. p. 41* (1900).

*Hypnum afro-glareosum* *Broth. in Engler's Bot. Jahrb. 1894, p. 214.*

2. *Lignic. — Afr. : Usambara.*

*afro-salebrosum* C. M. in *Rehm. M. Afr. austr. N.* 385 et in *Hedwig.* 1899, p. 136.

Br. *Dicksoni* *Rehm. in Coll. cit. n.* 383 b.

Br. *sulfurescens* C. M. in *sched.*

2. — *Afr.* : Transvaal, Orange.

*albicans* (Neck.) *Bryol. eur. fasc.* 52-54 (Vol. V), *Mon.* p. 19, t. 553 (1853); W. P. Sch. *Syn. II ed.* p. 644 (1876).

*Hypnum albicans* Neck. *Meth. M.* p. 180, n. 36 (1771); *Hedw. Descript. IV*, p. 13, t. 5 (1876); *Brid. Bryol. univ. II*, p. 492 (1827); C. M. *Syn. II*, p. 360 (1851).

H. *albicans*  $\beta$  *flavescens* *Brid. op. cit.* p. 494.

H. *albicans* *Roth Fl. Germ. III*, p. 303 (1800).

H. *rivulare* *Brid. Musc. rec. II, II*, p. 177 (1801).

H. *salebrosum* *Drumm. M. bor. am. I ed.* n. 175 p.p.

H. *sericeum* *gracile albicans*, capsulis subrotundis *Dill. Hist. M.* p. 328, t. 42, f. 63 (1741).

*Maschalocarpus Weissmannianus* *Spreng. ? (fid. Brid.)*.

*Neckera albicans* *Willd. Prodr. fl. Berol. n.* 940 (1787).

*Pterigynandrum cirrosum* *Brid. Bryol. univ. II*, p. 184.

2. In graminos., ad lat. viar. silv. et tecta stram. — *Eur.* : Sept. et intermed., saepe ster. ; in Britann. uberr. fructif. *As.* : *Caucas.* occ. *Am. sept.* : Groenland, tota borealis, Calif.

« *var. dumetorum* *Limpr. Kryptfl. v. Schles. I*, p. 73 (1876) et *Laubm. Deutschl.... III*, p. 125.

In silv. — *Eur.* : Brandeb., Siles., Havar., Wurtenb.

« *var. filiforme* *Th. Jens. Bryol. dan.* p. 178 (1856).

— *Eur.* : Dania.

« *var. flavescens* *ej. op. et l. cc.*

— *Eur.* : Dania, Mecklenb.

« *var. groenlandicum* (C. Jens.) *Par.*

*Hypnum albicans var. groenlandicum* C. Jens. in *Saertryk af Medd. om Groenland*, 1898, p. 437.

♂ et fr. ign. — *Am. sept.* : Fret. Scoresby.

« *var. julaceum* *Warnst. Moossfl. prov. Brandeb.* p. 70 (1885).

Ad terr. humid. — *Eur.* : Pomer., Siles., etc.

« *var. occidentale* *Ren. et Card. in Bot. Centralbl.* 1890, n. 51.

— *Am. sept.* : Washington, Montana.

« *var. pinnatifidum* *Warnst. in Verh. d. bot. Verein d. prov. Brandenb.* 1899, p. 71.

— *Eur.* : Lusatia infer.

« *var. rugulosum* *Warnst. in Allgem. bot. Zeitschr.* 1899, Beih. n. 1, p. 10.

— *Eur.* : Boruss.

*var. Thedenii* *Hartm.* — Br. *Thedenii*.

*ambiguum* de Not. } — Br. *salebrosum* (fid. *Vent. et Bott.*)  
 } — *Scleropodium illecebrum* (fid. *Limpr.*)

*amnicolum* C. M. in *Nuov. Giorn. bot. ital.* 1896, p. 125.

Ster. Saxic. secus rivul. — *As.* : China centr. (prov. Shen-Si).

*amœnum* *Milde in Hedwig.* 1869, p. 51.

Br. *plumosum var. amœnum* *Limpr. Laubm. Deutschl. etc. III*, p. 92.

- Br. plumosum* var. *homomallum* *Milde Bryol. stles. p. 326 (1869).*  
*Br. Venturii* *Warnst. in Fl. 1881, p. 34.*  
 1. Ad rup. quarz. — *Eur.* : Boruss. or., Francon., Tirol., Helvet.  
*antarcticum* *Card. in Rev. bryol. 1900, p. 46.*  
 Ster. — *Am. merid.* : Fret. Gerlache.  
 c var. *cavifolium* *ej. in op. et l. cc.*  
 Ubi typ.  
*aristatum* *Mitt.* — *Rhynchostegium.*  
*Arnoldianum* *Mol.* — *Br. campestre.*  
*asperinum* (*Mitt.*) *Kindb. Eur. and N. Am. Bryin, p. 116 (1897).*  
*Eurhynchium asperinum* *Jaeg. op. cit. p. 421.*  
*Hypnum asperinum* *Mitt. in Journ. of the Linn. Soc. VIII, p. 33, t. 6;*  
*C. M. in Bot. Zeit. 1858, p. 170.*  
 2. Lignic. et terr. — *Am. sept.* : Alaska, Columb. brit., Washington.  
 ins. Vancouver, Idaho, Calif.  
*asperulum* (*Hpe.*) *Jaeg. op. cit. p. 398.*  
*Br. plumosum* (*fid. Mitt.*).  
*Hypnum asperulum* *Hpe. in Linn. 1863, p. 159.*  
 1. Ad rivul. — *Am. merid.* : And. Bogot.  
*asprellum* *Jaeg.* — *Thelia.*  
*atrocladum* *Mitt. in Melliss's S. Helena, 1875, p. 365.*  
 1. — *Afr.* : Ins. S. Helena.  
*atrotheca* (*Duby*) *Besch. Fl. bryol. Réunion, etc. p. 153 (1880-81).*  
*Hypnum atrotheca* *Duby in Mém..... Genève, 1876, p. 8, t. 3, f. 2 et in*  
*Fl. 1877, n. 5.*  
*Isopterygium atrotheca* *Jaeg. op. cit. p. 499.*  
 1. Rupic. — *Afr.* : Usambara, Madagascar, ins. Borbon., Franciaë.  
*aureo-nitens* *Jaeg.* — *Meteorium.*  
*aureum* *de Not.* — *Camptothecium.*  
*auriculatum* (*Lindb.*) *Jaeg. op. cit. p. 406.*  
*Hypnum auriculatum* *Lindb. in Act. Soc. sc. Fenn. 1872, p. 250.*  
 2. — *As.* : Saghalien.  
*austro-alpinum* (*Hpe.*) *Jaeg. op. cit. p. 403.*  
*Hypnum austro-alpinum* *Hpe. in Linn. 1859, p. 610.*  
 1. — *Pac.* : Alp. Austral. or.  
*austro-glareosum* (*C. M.*) *Par. Ind. p. 131 (1893).*  
*Hypnum austro-glareosum* *C. M. in Engler's Bot. Jahrb. 1883, p. 82 et in*  
*Forschungsreise..... Bot. p. 32 (1889).*  
 Ster. — *Afr.* : Ins. Kerguelen.  
*austro-salebrosum* (*C. M.*) *Par. Ind. p. 131 (1894).*  
*Hypnum austro-salebrosum* *C. M. in op. cc. pp. 82 et 31.*  
*H. salebrosum* *Mitt. in Journ. of the Linn. Soc. 1876.*  
 Ster. — *Afr.* : Ins. Kerguelen.  
*Bellii* (*Mitt.*) *Par. Suppl. Ind. p. 41 (1900).*  
*Hypnum Bellii* *Mitt. in hb. Saharampur, n. 3736; Broth. in Act. Soc. sc.*  
*Fenn. 1898, n. 2, p. 35.*  
 2. In silv. *Betul.* — *As.* : N. W. Himalaya (Dehra Doun).

biventrosus (C. M.) Jaeg. *Ad. II*, p. 388 (1876-77).

Br. splendens Aust. in *Coult. Bot. Gaz. II*, p. III.

Hypnum biventrosus C. M. in *Bull. Torr. Bot. Club. V*, p. 49 (1878) et in *Fl. LVIII* p. 90 ( ); *Lesq. et Jam. Man.* p. 388 (1884).

1. In silv. profund. — *Am. sept.* : Carolin. sup., Florid., Ludovic.

Bolanderi (Lesq.) Jaeg. *Ad. II*, p. 390 (1876-77).

Hypnum Bolanderi Lesq. in *Trans. of the Amer. Phil. Soc. XIII*, p. 12 ( ); *Lesq. et Jam. Man.* p. 341 (1884).

2. Terr. — *Am. sept.* : Washington, Calif.

bolivio-plumosus C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 154.

— *Am. merid.* : And. Boliv.

Borgenii (Hpe.) Jaeg. *op. et l. cc.*

Sciuro-Hypnum Borgenii Hpe. in *Linn.* 1874, p. 220.

1. — *Afr.* : Madagascar.

brachycladum (Broth.) Par. *Suppl. Ind.* p. 42 (1900).

Hypnum brachycladum Broth. in *Act. Soc. sc. Fenn.* 1898, n. 2, p. 34.

2. — *As.* : Kashmir.

brevinerve Kindb. — *Plagiothecium brevipungens.*

Brotheri Par.

Br. flexicaule Par. *Suppl. Ind.* p. 44 (1900).

Hypnum flexicaule Broth. in *Hedwig.* 1899, p. 240.

1. — *As.* : Japon.

Bryhnii Kindb. — Br. collinum var. subulaceum.

brunneo-alare (C. M.) Jaeg. *op. et l. cc.*

Hypnum brunneo-alare C. M. in *Linn.* 1875, p. 654.

1. — *Am. sept.* : Mexic.

Buchanani (Hook.) Jaeg. *op. et l. cc.*

Hypnum Buchanani Hook. in *Trans. of the Linn. Soc. IX*, p. 320, t. 28, f. 3 ( ); *C. M. Syn. II*, p. 362 (1851); *Mitt. M. Ind. or.* p. 78 (1859).

2. — *As.* : Japon., Bhotan, Assam, Nepal., Himalaya,

M. Neilgherr.

« var. macrostegium Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1899, II, p. 247.

— *As.* : Sikkim.

caespitans W. P. Sch. in W. Schimp. *M. Abyss.*

— *Afr.* : Abyss.

caespitosum Dixon. — *Scleropodium.*

calcareum Kindb. in *Rev. bryol.* 1895, p. 86.

1. Ad rup. calcar. — *Am. sept.* : Canada (Ottawa).

Calerae (C. M.) Par. *Ind.* p. 131 (1894).

Hypnum Calerae C. M. in *Linn.* 1878-79, p. 452.

2. Ster. — *Am. merid.* : Argentin. Cordob.

californicum Jaeg. — *Scleropodium.*

cameratum (Mitt.) Jaeg. *op. cit.* p. 419.

Hypnum cameratum Mitt. *M. Ind. or.* p. 78 (1859).

2. — *As.* : Japon, Nepal., Himalaya.

campestre (*Bruch*) *Bryol. eur. fasc. 52-54 (Vol. V), Mon. p. 12, t. 545* (1853); *W. P. Sch. Syn. II ed. p. 634* (1876).

*Bryum Arnoldianum* *Mol. in Litz. Moosst. p. 120* (1864).

*Br. Schimperi* *H. v. Klinggr. Hoch. Krypt. Preuss. p. 168* (1858.)

*Br. subalbicans* *de Not. Cron. II, p. 120* (1867) *et Epil. p. 118* (1869).

*Hypnum campestre* *Bruch in sched.*

*H. geniculatum* *Schleich. in sched.*

*H. rutabulum* *var. campestre C. M. Syn. II, p. 368.*

1. In coll. et camp. gramin., agr. et silvat. siccis. — *Eur.* : Suecia, centr. tota, Helvet., Pyren. *As.* : Jéuissei sup., Talysch, Kashmir, Caucas. *Afr.* : Alger. *Am. sept.* : New Brunswick, Nov. Angl., M. Alb., Idaho, Columb. brit., Colorado.

*campestre* *Mitt.* — *Br. salebrosum.*

*campylothallum* *C. M. in Nuov. Giorn. bot. ital. 1896, p. 124 et 1897, p. 268.*

1 ?

— *As.* : China centr. (prov. Schen-Si).

*cavernosum* *Kindb. in Rev. bryol. 1895, p. 81.*

— *Am. sept.* : Canada.

*Chauveti* *Ren et Card. in Bull. Soc. roy. bot. Belg. 1894, II, p. 130.*

2. fr. ign.

— *Afr.* : Ins. Borbon.

*chloropterum* (*C. M. et Kindb.*) *Par. Suppl. Ind. p. 42* (1900).

*Eurhynchium chloropterum* *Kindb. Eur. and N. Am. Bryin. p. 94* (1897).

*Hypnum chloropterum* *C. M. et Kindb. in Mac. Cat. p. 231* (1892); *Par. Ind. p. 620* (1894).

*H. Novae Angliae* *Mao, Canad. M. n. 440* (recte ! *fid. Grout in Bull. Torr. bot. Club. 1899, p. 229.*)

1. Ad rup. et terr. umbr. — *Am. sept.* : Terra nov., New Brunswick.

*chrysocladon* (*Hpe.*) *Jaeg. op. cit. p. 404.*

*Hypnum chrysocladon* *Hpe. in Ann. sc. nat. V ser., p. 341* (1866); *Mitt. M. austr. am. p. 564* (1869).

*Ster.*

— *Am. merid.* : And. Nov. Granat.

*cirrosum* *W. P. Sch.* — *Eurhynchium.*

*var. Funckii* *Jaeg.* — *E. cirrosum* *var. Funckii.*

*var. gracillimum* *Mol.* — *E. cirrosum* *var. Molendoi.*

*cladoneuron* (*C. M.*) *Par. Ind. p. 132* (1894).

*Camptothecium cladoneuron* *Jaeg. op. cit. p. 382.*

*Hypnum cladoneuron* *C. M. in Linn. 1874, p. 652.*

2.

— *Am. sep.* : Mexic.

*clinocarpum* *Jaeg.* — *Eurhynchium.*

*cochleare* *C. M. in Nuov. Giorn. bot. ital. 1897, p. 177.*

*Aquatic. ?* — *Am. merid.* : And. Boliv.

*collinum* (*Schleich.*) *Bryol. eur. fasc. 52-54 (Vol. V), Mon. p. 15, t. 548* (1853); *W. P. Sch. op. cit. p. 647.*

*Br. Holleri* *Mol. Bayerns Laubm. p. 241* (1875).

*Eurhynchium collinum* *Kindb. op. cit. p. 165.*

*Hypnum collinum* *Schleich. Cat. 1815; C. M. op. cit. p. 429.*

1. Ad terr. et in fiss. rup. — *Eur.* : Spitzberg, alp. Norveg., Tatra, Salisb., Stir., Carinth., Rhaet., Helvet., Tirol., Sabaud., Pyren., Arvern. *As.* : Japon, Alatau, Kashmir, Gilgit, Baltistan, Caucas. *Am. sept.* : Grœnland, Terra nov., Athabasc., Assiniboia, reg. occid. usq. ad Calif.

- var. *Bryhnii* Kaur. — Br. collinum var. subjulaceum.
- « var. *saporensae* Besch. in *Ann. sc. nat.* 1893, p. 373.  
— *As.* : Japon.
- « var. subjulaceum Pfeff. *Bryog. Stud.* p. 77 (1869).  
Br. *Bryhnii* Kindb. in *Bot. Notis.* 1886.  
Br. collinum var. *Bryhnii* Kaur. in *Nyt. Mag. for. Nature*, 1888, p. 218.  
Br. nova spec. Kindb. *Enum. n.* 133 (1888).  
*Eurhynchium* *Bryhnii* Kindb. *Eur. and N. Am. Bryin.* p. 106 (1897).  
In schistos. aq. nival. irror. — *Eur.* : Norveg. (Dovre), Helvet. *Am. sept.* : Fret. Scoresby.
- coloradense* Ren. et Card. — *Eurhynchium* *Boscii* var. *coloradense*.  
*colpophyllum* Kindb. — *Eurhynchium*.
- columbico-rutabulum* Kindb. in *Mac. Cat.* p. 198 (1892).  
1. Lignic. — *Am. sept.* : Columb. brit.
- complanatulum* C. M. in *Hedwig.* 1899, p. 138.  
Ster. — *Afr.* : Transvaal.
- comtifolium* (C. M.) Jaeg. op. cit. p. 392.  
*Hypnum* *comtifolium* C. M. in *Linn.* 1876, p. 653.  
2. — *Am. sept.* : Mexic.
- conostomum* (Tayl.) Jaeg. op. cit. p. 392.  
*Hypnum* *conostomum* Tayl. in *Lond. Journ. of bot.* 1848 ; C. M. *Syn. II*, p. 364 (1851).  
H. *Taylori* Mitt. *M. austr. am.* p. 560 (1869).  
*Syn. Terr.* — *Am. merid.* : And. Quitens. et Boliv.
- crassinerve* Kindb. — *Eurhynchium*.
- crocatum* Hpe. *miss.* ; C. M. in *Bull. hb. Boissier*, 1897, p. 218.  
— *Am. sept.* : Guatemala.
- curtum* Lindb. in *Meddel. af Soc. pr. Faun. et Fl. fenn.* 1879, p. 5 ;  
*Limpr. Laubm. Deutschl.*... III, p. 101 (1896).  
Br. *rutabulum* var. *explanatum* (Brid.) Brockm. in *Archiv. Ver. Freund. Naturg. Mecklenb.* 1870, p. 122.  
Br. *Starkii* Rab. *Bryoth. eur. n.* 1188.  
Br. *Starkii* var. *alpestre* *Bryol. eur. fasc.* 52-54 (vol. VI) *Mon.* p. 10, t. 541 (1853) ; *W. P. Sch. Syn. I ed.* p. 540 (1860).  
Br. *Starkii* var. *curtum* Warnst. *Moosfl. Prov. Brandenb.* p. 71 (1885).  
Br. *Starkii* f. *major* Milde *Bryol. siles.* p. 333 in *adn.* (1869) ; Rab. *Bryoth. eur. n.* 1039.  
Br. *Starkii* var. *praelongum* W. P. Sch. *Syn. I ed.* p. 541 et *II ed.* p. 652 (1876) p. p.  
Br. *Starkii* var. *robustum* *Limpr. Kryptfl. v. Schles. I.* p. 75 (1876).  
*Hypnum* *curtum* Lindb. *M. scand.* p. 35 (1879).  
H. *rutabulum* ξ *explanatum* Brid. *Sp. M. II*, p. 184 (1812).  
H. *Starkii* Funck *Deutsch. M.* p. 61, t. 44, n. 41 (1820).  
1. Ad terr. humid. et rad. arb. — *Eur.* : Fenn., German., Siles., Bohem. *As.* : *Jenissei* super., Kashmir.
- curtum* Ren. et Card. — Br. *œdipodium*.
- curvatulum* (Broth.) Par. *Suppl. Ind.* p. 43 (1900).  
*Hypnum* *curvatulum* Broth. in *Act. Soc. sc. Fenn.* 1898, n. 2, p. 38.  
1. Rupic. — *As.* : Kashmir.



- cuspidiferum (Mitt.) Jaeg. op. cit. p. 410.  
 Hypnum cuspidiferum Mitt. M. Ind. or. p. 78 (1859).  
 2 ? — As. : Sikkim.
- cylindricarpum (Mitt.) Far. Ind. p. 133 (1894).  
 Hyoconium cylindricarpum Mitt. in Trans. of the Linn. Soc. 1891, p. 178.  
 2 ? — As. : Alp. Bhotan.
- cyrtophyllum Kindb. in Ottawa Nat. IV, p. 63 (1889) et in Mac. Cat. p. 191 (subsp. Br. acuminati) (1892).  
 Ad Ulmos in silv. — Am. sept. : Ontario, Minnesota.
- declivum Jaeg. — Br. velutinum.
- decurvans (Mitt.) Jaeg. op. cit. p. 410.  
 Hypnum decurvans Mitt. M. Ind. or. p. 78 (1859).  
 — As. : W. Himalaya, Sutledje.
- densum Jur. — Amblystegium.
- Dicksoni Rehm. — Br. afro-salebrosum.
- dicranoides C. M. in Nuov. Giorn. bot. ital. 1898, p. 201.  
 Ster. — As. : China centr. (prov. Schen-Si).
- Diecke Roell in Hedwig. 1897, Beibl. II, p. 41.  
 2 ? Ad arbor. — Eur. : Hispan. (S. de Guadarrama).
- digastrum C. M. et Kindb. in Bidr. p. 453 (nom. 1890) et in Mac. Cat. p. 190 (1892).  
 1. Rupic. — Am. sept. : New Brunswick, Ontario, Chicago.
- Donnellii (Aust.) Ren. et Card. in Bot. Centralbl. 1890, n. 51.  
 Hypnum Donnellii Aust. in Coult. Bot. Gaz. IV, p. 162 ( ); Lesq. et Jam. Man. p. 338 (1884).  
 Stereophyllum Donnellii Card. ap. Grout Rev. N. Am. Isothec. and Brachyth. in Mem. Torr. Bot. Club, VI, p. 207 (1879).  
 1. Lignic. — Am. sept. : Ontario, Florid.
- Durieu de Not. — Orthothecium ?
- duriusculum W. P. Sch. mss.  
 Hypnum duriusculum C. M. Syn. II, p. 426 (1851).  
 H. plumosum Mitt. M. austr. am. p. 565 (1869).  
 1. Ad rup. trachyt. — Am. sept. : Mexic.
- Ehrenbergii Ltz. M. Ehrenb. p. 51, t. 13, f. 1-28 (1868).  
 Ster. — As. : Syria.
- erythropyxis C. M. in Rehm. M. Afr. austr. n. 382 et in Hedwig. 1899, p. 237.  
 — Afr. : Natal.
- erythrorhizon Bryol. eur. fasc. 53-54 (Vol. VI), Mon. p. 14, t. 547 (1853); W. P. Sch. Syn. II ed. p. 646 (1876).  
 H. erythrorhizon Hartm. Skand. Fl. VI ed. p. 355 (1854).  
 1 (Bryol. eur.); 2 (Limpr.). Lignic. et saxic. — Eur. : Scand., Fennia, Karel., Stir., Carinth. As. : Jenissei med. et super. Am. sept. : Washington.
- var. Thedenii Lindb. — Br. Thedenii.

- eupopuleum (C. M.) Par. Ind. p. 134 (1894).  
 Hypnum eupopuleum C. M. in Linn. 1878-79, p. 500.  
 1. — Am. sept. : Venezuela.
- euryodictyon (C. M.) Par. op. et l. cc.  
 Hypnum euryodictyon C. M. in Engler's Bot. Jahrb. 1883, p. 82 et in Forschungsreise.... Bot. p. 82 (1889).  
 Ster. — Afr. : Ins. Kerguelen.
- eustegium Besch. in Ann. sc. nat. 1893, p. 375.  
 2. — As. : Japon.
- fagineum Kindb. — Eurhynchium Vancheri var. tenuicaule.  
 falcatum (Broth.) Par. Suppl. Ind. p. 43 (1900).  
 Hypnum falcatum Broth. in Act. Soc. sc. Fenn. 1898, n. 2, p. 31.  
 1. Cortic. et lignic. — As. : Kashmir.
- fasciculato-caudatum C. M. in Hedwig. 1897, p. 129.  
 Ster. — Am. merid. : Argentin. subtrop. (Chaco).
- fasciculirameum C. M. in Nuov. Giorn. bot. ital. 1897, p. 269.  
 — As. : China centr. (prov. Schen-Si).
- Fendleri (Sull.) Jaeg. Ad. II, p. 393 (1876-77).  
 Hypnum Fendleri Sull. M. of U. St. p. 76 (1856) et Ic. M. p. 189, t. 117 (1864); Lesq. et Jam. Man. p. 340 (1884).  
 H. velutinum  $\delta$  microcarpum C. M. Syn. II, p. 400 (1851).  
 Leskea Fendleri Sull. in Mem. of the Amer. Acad.... 1849, p. 169, t. I.  
 1 et Syn. — Am. sept. : Nov.-Mexic.
- filiforme Jur. in sched.; Limpr. Laubm. Deutschl.... III, p. 138 (1896).  
 Ster. Ad rup. calcar. — Eur. : Austria infer. (m. Schneeberg).
- filirameum C. M. in Hedwig. 1897, p. 127.  
 2. — Am. merid. : Argentin. subtrop. Tucuman.
- Fitz-Geraldi (C. M.) Ren. et Card. in Bot. Centralbl. 1890, p. 51.  
 Hypnum Fitz-Geraldi C. M. in Fl. 1897, p. 224.  
 2. — Am. sept. : Florid.
- flaccum C. M. in Nuov. Giorn. bot. ital. 1897, p. 156.  
 — Am. merid. : And. Boliv.
- flavescens Kindb. — Br. rivulare.
- flexicaule Ren. et Card. in Grout Mem. Torr. bot. Club, VI, p. 167 (1879).  
 — Am. sept. : M. Adirondack, Vermont, New Jersey, Pensilv., Tennessee, Columb. brit.
- flexicaule (Broth.) Par. — Br. Brotheri.
- flexiventrosus (C. M.) Jaeg. Ad. II, p. 387 (1876-77).  
 Hypnum flexiventrosus C. M. in Linn. 1876, p. 653.  
 Ster. — Am. sept. : Mexic.
- frigidum (C. M.) Besch. Prodr. bryol. mexic. p. 104 (1871).  
 Hypnum frigidum C. M. in Bot. Zeit. 1856, p. 456  
 Syn. — Am. sept. : Mexic.
- Funckii } de Not. — Br. glaciale.  
 W. P. Sch. — Eurhynchium cirrosus var. Funckii.

- garovaglioides *C. M. in Nuov. Giorn. bot. ital.* 1897, p. 270.  
— *As.* : China centr. (prov. Schen-Si).
- Geheebii *Milde in Hedwig.* 1869, p. 161; *W. P. Sch. Syn. II ed.* p. 651 (1876).  
Br. ketum *Milde Bryol. siles.* p. 336 p.p. (1869).  
Camptothecium Geheebii *Kindb. Eur. a. N. Amer. Bryin.* p. 90 (1897).
2. Ad rup. basalt. umbr. — *Eur.* : Norveg., Siles., Bohem., Salisb., Rhoengeb., Baden.
- gelidum *Bryhn Descript. M. Norveg.* 1899, p.  
Br. glaciale *Auct. p.p.*  
1. Rupic. in riv. et terr. ad mol. glac. — *Eur.* : Alp. Norveg. sept. et merid.
- gemmascens *C. M. et Kindb. in Bidrag.* p. 453 (nom. 1899) et in *Mac. Cat.* p. 195 (1892).  
1. Lignic. — *Am. sept.* : Columb. brit.
- georgico-glareosum (*C. M.*) *Par. Ind.* p. 134 (1894).  
Hypnum georgico-glareosum *C. M. Bryol. austr. georg.* p. 42 (1890).  
*Ster.* Aquat. — *Am. merid.* : Austr. Georg., fret. Gerlache.
- glabratum (*Broth.*) *Par. Suppl. Ind.* p. 44 (1900).  
Hypnum glabratum *Broth. in Act. Soc. sc. Fenn.* 1898, n. 2, p. 36.  
1. Cortic. — *As.* : Kashmir.
- glaciale *Bryol. eur. fasc. 53-54 (Vol. VI), Mon. f.* 11, t. 542 (1853); *W. P. Sch. Syn. II ed.* p. 652 (1876).  
Br. Funckii *de Not. Cron. II*, p. 21 (1867) et *Epil.* p. 125 (1869).  
Eurhynchium glaciale *Kindb. Eur. a. N. Am. Bryin.* p. 100 (1897).  
? Hypnum Funckii *Hsch. in sched.*; *C. M. Syn. II*, p. 429 (1851).  
H. glaciale *C. Hartm. Skand. Fl. IX ed.* (1864).  
1. Ad terr. et saxa humida juxta niv. — *Eur.* : Ins. Ursor., summ. Alp. Scot., Norveg., Tatrae, Stir., Carinth., Salisb., Julian., Algov., Bavar., Tirol., Helvet., Sabaud.; *Pyren. As.* : Tibet occid. *Am. sept.* : Groenland, prom. Breton.
- « var. dovense *Limpr. Laubm. Deutschl. ... III*, p. 112 (1896).  
Secus rivul. niv. — *Eur.* : Norvegia (Dovre).
- « var. Huntianum (*W. P. Sch.*) *Limpr. op. cit.* p. 113.  
Br. Huntianum *W. P. Sch. mss.*  
Br. glaciale var. subsecundum *Husn. Muscol. gall.* p. 328 (1893).  
Br. Starkii γ praelongum *W. P. Sch. Syn. II ed.* p. 652 p.p. (1876).  
*Ster.* — *Eur.* : M. Scotiae, Sabaud. (M. Blanc).
- « var. majus *Husn. op. et l. cc.*  
— *Eur.* : Alp. Sabaud. (M. Blanc).
- var. subsecundum *Husn.* — Br. glaciale var. Huntianum.
- glareosum (*Bruch*) *Bryol. eur. fasc. 52-54 (Vol. VI), Mon. p.* 19, t. 552 (1853); *W. P. Sch. op. cit.* p. 644.  
Hypnum albicans var. minnidunense *Brid. Bryol. univ. II*, p. 494 (1827).  
H. glareosum *Br. in sched.*  
H. lutescens *Drumm. M. bor. am. I ed.* p. 174.  
H. plumosum *Hedw. M. frond. IV*, p. 37, t. 15, p.p. (1794).  
H. salebrosus *H. et T. Muscol. brit.* p. 166, t. suppl. 5 p.p. (1818).
2. In glareos. et gramin. — *Eur.* : Praccip. interm., saepe ster. *As.* : Jenissei inf., Caucas. occid. et centr. *Am. sept.* : Columb. brit.

- « *var. alpinum* de Not. Cron. II, p. 19 (sub Br. albicante *var. alp.* (1867) et *Epil.* p. 116 (1869).  
Br. glareosum *var. fastigiatum* Warnst. in *Hedwig.* 1885, n. 2.  
— *Eur.* : Alp. Salisb., Carinth., Helvet., Ticin.
- « *var. fastigiatum* Warnst. — Br. glareosum *var. alpinum.*
- « *var. rugulosum* Pfeff. *Bryog. Stud.* p. 77 (sub Br. tauriscorum *var. rugul.*) (1869).  
Ad rup. schist. — *Eur.* : Alp. Salisb., Stir., Tirol., Helvet. As. : Alatau.
- glauco-viride C. M. in *Nuov. Giorn. bot. ital.* 1898, p. 198  
— *As.* : China centr. (prov. Schen-Si).
- glauculum C. M. in *op. cit.* p. 199.  
1. — *As.* : China centr. (prov. Schen-Si).
- gloriosum (C. M.) Par. *Ind.* p. 135 (1894).  
Hypnum gloriosum C. M. in *Fl.* 1890, p. 498.  
Ster. — *Afr.* : Kilima N'Djaro.
- grandirete C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 155.  
1. — *Am. merid.* : And. Boliv.
- « *var. robustum* ej. in *op. et l. cc.*  
Ubi typ.
- graniticum Gumb. — Br. vagans.
- harpidioides C. M. et Kindb. in *Bidrag.* p. 453 (nom., 1890) et in *Mac. Cat.* p. 194.  
Hypnum harpidioides Kindb. *Eur. a. N. Am. Bryin.* p. 133 (1897).  
2. Ster. Lignic. — *Am. sept.* : New Brunswick, Columb. brit.
- helvolum (Mont.) Jaeg. *Ad. II*, p. 402 (1876-77).  
Hypnum helvolum Mitt. *M. austr. am.* p. 564 (1869).  
Leskea helvola Mont. *Syll.* p. 19 (1856).  
Cortic. — *Am. merid.* : And. Quitens.
- Hillebrandi (Lesq.) Sull.  *Ic. M. Suppl.* p. 98, t. 74 (1874).  
Hypnum Hillebrandi Lesq. in *Mem. Calif. Acad. I*, p. 33 ( ); Lesq. *Jam. Man.* p. 340 (1884).  
1. — *Am. sept.* : Calif., Alabama.
- hirtipes W. P. Sch. in *Mandon Pl. Boliv.* n. 1703.  
Rhynchostegium hirtipes C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 157  
— *Am. merid.* : And. Boliv.
- Hollereri Mol. — Br. collinum.
- homocladum C. M. in *Nuov. Giorn. bot. ital.* 1896, p. 126.  
Ster. — *As.* : China centr. (prov. Schen-Si).
- Hopfferi (Welw. et Duby) Par. *Ind.* p. 135 (1894).  
Hypnum Hopfferi W. et Dy. *Choix de Crypt.*..., 1871, p. 7, t. 2, f. 2.  
1. Lignic. — *Afr.* : Ins. Sao Thome.
- horricoma C. M. in *hb. Saharampur* n. 3736 p.p.  
— *As.* : N. W. Himalaya (Dehra Doun).

- Huntianum* W. P. Sch. — Br. glaciale var. *Huntianum*.  
*idahense* Ren. et Card. in Bot. Gaz. 1890, p. 60, t. 9 c.  
 1. — Am. sept. : Idaho.  
*illecebrum* de Not. — *Scleropodium*.  
*implicatum* (Hsch.) Jaeg. Ad. II, p. 388 (1876-77).  
*Hypnum implicatum* Hsch. in Mundt et Maire M. capens. ; C. M. Syn. II  
 p. 362 (1851).  
 2. — Afr. : Prom. B. Sp., reg. lacustr.  
*Inandae* Rehm. M. Afr. austr. n. 381.  
 — Afr. : Natal.  
*inconditum* Besch. Fl. bryol. Réunion, etc. p. 155 (1880-81).  
 1. In gramin. humid. — Afr. : Ins. Bourbon.  
*indistinctum* C. M. in Ren. Prodr. fl. bryol. Mad. etc. p. 230 (1897).  
 — Afr. : Madagascar.  
*intricatum* Kindb. — Br. velutinum var. *intricatum*.  
 var. *declivium* Kindb. — Br. velutinum.  
*jamaicense* C. M. in Bull. hb. Boissier, 1897, p. 567.  
 Ster. — Am. sept. : Jamaic.  
 « var. *album* ej. in op. et l. cc.  
 Ubi typ.  
*jenissense* (Lindb. et Arn.) Par. Ind. p. 135 (1894).  
*Hypnum jenissense* L. et A. M. As. bor. 1890, II, p. 133.  
 2. Terr. in silvul. — As. : Jenissei med.  
*jucundum* de Not. Epil. p. 118 (1869); W. P. Sch. Syn. II ed.  
 p. 643 (1876).  
 Br. salebrosum (fid. Vent. et Bott. Enum. p. 9).  
 1. — Eur. : Ital.  
*julaceum* W. P. Sch. in Mandon Pl. Boliv. n. 1707 bis.  
 — Am. sept. : And. Boliv.  
*kashmirensense* (Broth.) Par. Suppl. Ind. p. 45 (1900).  
*Hypnum kashmirensense* Broth. in Act. Soc. sc. Fenn. 1898, n. 2, p. 37.  
 1. Ad Abiem Webbianam. — As. : Kashmir.  
*kamounense* (Harv.) Jaeg. Ad. II, p. 416 (1876-77).  
*Hypnum kamounense* Harv. in Hook. Ic. Pl. rar. t. 24, f. 10 (1841) et in  
 Lond. Journ. of bot. 1840, p. 19; C. M. Syn. II, p. 365 (1851) Mitt.  
 M. Ind. or. p. 79 (1859).  
 — As. : Nepal., N. W. Himalaya.  
*Knysnae* C. M. in Rehm. M. Afr. austr. n. 387 et in Hedwig. 1899, p. 137.  
 — Afr. : Austral.  
*Krausei* Par. — *Hypnum*.  
*kuroischicum* Besch. in Ann. sc. nat. 1893, p. 373.  
 1. — As. : Japon.  
*labradoricum* (Kindb.) Par. Suppl. Ind. p. 45 (1900).  
*Eurhynchium labradoricum* Kindb. Eur. a. N. Am. Bryin. p. 101 (1897).  
 1. — Am. sept. : Labrador, prom. Breton.

*laetum* (Brid.) *Bryol. eur. fusc.* 52-54 (Vol. VI), *Mon.* p. 20, t. 554 (1853); *W. P. Sch. Syn. II ed.* p. 639 (1876).

*Hypnum laetum* Brid. *Bryol. univ. II*, p. 479 (1827).

*H. luteolum* C. M. *Syn. II*, p. 357 (1851).

*H. salebrosum* *Drumm. M. bor. am. I ed. n.* 175 p.p.

2. Ad saxa calc. rupesq. basalt. — *Eur.* : Central. tota; *Jurass. Am. sept.* : Labrador, Anticosti, Terr. nov., Canada or., reg. sept., centr. et or.; *Ludovic., Calif.*

« *var. dentatum* (Lesq. et Jam.) *Par. Ind.* p. 136 (1894).

*Br. Sullivantii* *Bryol. eur. l. c.* p. 21.

*Hypnum laetum var. dentatum* Lesq. et Jam. *Man.* p. 335 (1884).

*H. salebrosum* Sull. *M. Allegh. n.* 43 (1846).

Ut et ubi typ. in *Am. sept.*

« *var. fallax* Ren. et Card. in *Bot. Centralbl.* 1890, n. 51.

— *Am. sept.* : Indiana, Calif.

« *var. gracillimum* Mol. in *Ltz. Moosst.* p. 106 (1864).

— *Eur.* : Bavar.

« *var. lignicola* Mol. in *Ltz. Beitrage*, p. 85 (1860).

— *Eur.* : Bavar.

« *var. pseudo-acuminatum* Ren. et Card. in *op. et l. cc.*

— *Am. sept.* : Indiana.

« *var. Roellii* eor. in *op. et l. cc.*

— *Am. sept.* : Indiana.

*laetum* Mol. p.p. — *Br. Geheebii*.

*laevifolium* (Mitt.) Jaeg. *Ad. II*, p. 411 (1876-77).

*Hypnum laevifolium* Mitt. *M. austr. am.* p. 565 (1869).

— *Am. merid.* : And. Nov. Granat.

*laevisetum* Kindb. — *Br. salebrosum*.

*lamprocarpum* (C. M.) Jaeg. *op. cit.* p. 387.

*Hypnum lamprocarpum* C. M. *Syn. II*, p. 359 (1851); *Bryol. jav. II*, p. 164, t. 263 (1861-70).

1. Arbor. (ad *Gnaphalium javanicum*). — *As.* : Java.

*lamprochyseum* C. M. et Kindb. in *Bidrag.* p. 453 (nom., 1890) et in *Mac. Cat.* p. 199.

*Camptothecium lamprochyseum* Kindb. *Eur. and N. Am. Bryin.* p. 88 (1897)

1. Saxic. — *Am. sept.* : Ins. Vancouver, Washington, Calif.

« *var. giganteum* Grout in *Bull. Torr. Bot. Club*, 1897, p.

— *Am. sept.* : Absque loco.

*lanceolatum* Warnst. in *Verh. d. bot. Ver. d. prov. Brandenb.* 1899, p. 70 et in *Abh.* ... 1900, p. 212.

1. Ad rad. arb. — *Eur.* : Lusat. infer.

? *lapponicum* W. P. Sch. — *Orthotrichum chryseum f. lapponica*.

*latifolium* (Lindb.) Ren. et Card. *Fl. ins. Miquelon*, p. 51 (1888).

*Br. rivulare var. latifolium* Husn. *Muscol. gall.* p. 322 (1893).

*Hypnum latifolium* Lindb. *M. nonnull. scand. in Medd. af Soc. pr. Faun. et Fl. fenn.* 1879, p. 5 et *M. scand.* p. 35, n. 98 (1879).

2. Ad rup. madid. — *Eur.* : Lapon., Norveg., Alp. Salisb., Stir., Carinth., Helvet., Vales., Longobard. *As.* : Jenissci med. *Am. sept.* : Ins. Miquelon.
- Leibergii *Grout in Bull. Torr. bot. Club.* 1897, p.  
— *Am. sept.* : Idaho.
- lepidopiloides (*C. M.*) *Par. Ind.* p. 137 (1894).  
Hypnum lepidopiloides *C. M. in Linn.* 1878-79, p. 450.  
Ster. — *Am. merid.* : Argent. Cordob.
- lepturum (*Tayl.*) *Jaeg. Ad. II*, p. 395 (1876-77).  
Hypnum lepturum *Tayl. in Lond. Journ. of bot.* 1846, p. 61; *C. M. Syn. II ed.* p. 454 (1851).  
Ster. — *Pac.* : Austral., fret. reg. Georges.
- leskeifolium *C. M. in Rehm. M. Afr. austr. n.* 373.  
Rhynchostegium leskeifolium *ej. in Hedwig.* 1899, p. 139.  
Lignic. — *Afr.* : Pr. B. Sp.
- leucoglaucom *C. M. et Kindb. in Bidrag* p. 453 (nom., 1890) et in *Mac. Cat.* p. 198 (1892).  
1. Terr. — *Am. sept.* : New Brunswick.
- ligusticum (*de Not.*) *W. P. Sch. Syn. I ed.* p. 692 (1866) et *II ed.* p. 642 (1876).  
Hypnum ligusticum *de Not. mss.*  
1. In pasc. humid. — *Eur.* : Ligur., Neap. *Afr.* : Alger.
- longicuspdatum (*Mitt.*) *Jaeg. op. cit.* p. 409.  
Hypnum longicuspdatum *Mitt. M. Ind. or.* p. 77 (1859).  
2? — *As.* : Sikkim.
- longidens (*C. M.*) *Par. Ind.* p. 187 (1894).  
Hypnum longidens *C. M. in Fl.* 1885, p. 427.  
1. Terr. — *Am. merid.* : Fuegia.
- lutescens *de Not.* — *Camptothecium.*  
β *fallax* *Vent. et Bott.* — *Homalothecium fallax.*
- Mac Owani *Rehm. M. Afr. austr. n.* 380.  
— *Afr.* : Natal.
- macrocomum (*C. M.*) *Par. Ind.* p. 137 (1894).  
Hypnum macrocomum *C. M. in Linn.* 1878-79, p. 454.  
1. — *Am. merid.* : Argentin. Salt.
- α *var. robustius* (*C. M.*) *Par. op. et l. cc.*  
Hypnum macrocomum *var. robustius C. M. in op. et l. cc.*  
— *Am. merid.* : Argentin. subtropic.
- mammilligerum *Kindb. in Mac. Cat.* p. 192 (*subsp. Br. salebrosi*) (1892).  
1. Lignic. — *Am. sept.* : Columb. brit.
- Maximilianum* *Gumb.* — *Br. plumosum.*
- melanangium *C. M. in Hildebr. M. Mad. n.* 2078.  
— *Afr.* : Madagascar.
- melanocarpum *Jaeg.* — *Br. oxyrhynchum.*

- membranosum *C. M. in Hedwig*. 1899, p. 133.  
 1. — *Afr.* : Pr. B. Sp.
- meridense (*C. M.*) *Jaeg. Ad. II*, p. 393 (1876-77).  
 Hymn meridense *C. M. Syn. II*, p. 433 (1851).  
 2. — *Am. merid.* : And. Venezuel.
- micrangium *C. M. in Nuov. Giorn. bot. ital.* 1896, p. 127.  
 — *As.* : China centr. (prov. Schen-Si).
- micropus *Bryol. eur. fasc.* 52-54 (*Vol. VI*), *Mon.* p. 9, t. 540 (1853); *W. P. Sch. Syn. II ed.* p. 650 (1876).  
 1. Ad rup. micac. schist. — *Eur.* : Helvet. (Grimsel).
- Mildeanum *W. P. Sch. mss.*; *Milde in Bot. Zeit.* 1862, p. 453.  
 Br. acutum *Sull. Ic. M. Suppl.* p. 99, t. 7 (1874).  
 Br. salebrosum γ palustre *W. P. Sch. Syn. II ed.* p. 641 (1876).  
 Hypnum acutum *Mitt. in Journ. of the Linn. Soc.* 1865, p. 33, t. 6.  
 H. Mildeanum *W. P. Sch. Syn. I ed.*, p. 694 (1860).  
 H. plumosum var. *Mildei Kindb. Laubm. Schwed. u. Norw.* p. 36 (1883).  
 1 vel polyg. Ad terr. argill. humid. — *Eur.* : Lappon., Suec., Dania, zona interm. tota. *Am. sept.* : Fret. Hudson, Scoresby, septentr. tota.  
 « var. robustum *Warnst. in Verh. d. bot. Ver. d. prov. Brandenb.* 1899, p. 71.  
 — *Eur.* : Boruss. or., Lusat. infer.
- minusculifolium *C. M. in Hedwig*. 1897, p. 128.  
 2. — *Am. merid.* : Argentin. subtropic. Salt.
- minutirete *C. M. in Hedwig*. 1899, p. 134.  
 2. — *Afr.* : Pr. B. Sp.
- mirabundum *C. M. et Kindb. in Bidrag*, p. 453 (*nom.*, 1890) *et in Mac. Cat.* p. 199 (1892).  
 1. Lignic. — *Am. sept.* : New Brunswick.
- Miyabei (*Broth.*) *Par. Suppl. Ind.* p. 46 (1900).  
 Hypnum Miyabei *Broth. in Hedwig*. 1899, p. 238.  
 2. Cortic. — *As.* : Japon.
- Molendoi* } *Itz.* — Br. plumosum.  
 } *W. P. Sch.* — Eurhynchium cirrosum var. *Molendoi*.
- molliculum (*Lindb.*) *Par. Ind.* p. 138 (1894).  
 Hypnum molliculum *Lindb. in Broth. Enum. M. Caucas* p. 112 (1892).  
 2. Lignic. — *As.* : Caucas. occ.
- molliculum Reichdt. — *Isopterygium*.
- molirameum *C. M. in Hedwig*. 1897, p. 129.  
 2. — *Am. merid.* : Argentin. subtrop. (Chaco).
- Morenoi *C. M. in op. cit.* p. 128.  
 Ster. — *Am. merid.* : Patagonia.
- norienae *Besch. in Ann. sc. nat.* 1893, p. 375.  
 2. Terr. — *As.* : Japon.
- nanopes *C. M. et Kindb. in Bidrag*, p. 453 (*nom.*, 1890) *et in Mac. Cat.* p. 201 (1892).  
 1. Terr. — *Am. sept.* : Columb. brit.



nematogonium (*C. M.*) *Par. Ind.* p. 138 (1894).

Hypnum nematogonium *C. M. in Linn.* 1878-79, p. 453.

Ster. — *Am. merid.* : Argentin. Cordob.

nigro-viride (*C. M.*) *Par. op. et l. cc.*

Hypnum nigro-viride *C. M. in Fl.* 1890, p. 498.

Ster. — *Afr.* : Kilima N'Djaro.

noësicum *Besch. in Ann. sc. nat.* 1893, p. 377.

2. — *As.* : Japon.

*nova species* Kindb. (Enum.) — *Br. collinum var. subjulaceum.*

Novae Angliae (*Sull. et Lesq.*) *Jaeg. Ad. II*, p. 394 (1876-77).

Eurhynchium Novae Angliae *Kindb. Eur. and N. Am. Bryin.* p. 100 (1897).

Hypnum Novae Angliae *Sull. et Lesq. M. bor. am. I ed. N.* 338; *Sull. M. of U. St.* p. 76 (1856) *et Ic. M. p.* 191, t. 118 (1864); *Lesq. et Jam. Man.* p. 378 (1884).

Scleropodium Novae Angliae *Besch. in Ann. sc. nat.* 1893, p. 78.

2. — *As.* : Japon. *Am. sept.* : Canada (fid. Card.),  
Nov. Angl., ins. Miquelon, Wisconsin, Alaska, Washington.

« *var. Delamarei Ren. et Card. Fl. Miquelon*, p. 50 (1888).

— *Am. sept.* : ins. Miquelon.

novo eboracense *A. J. Grout in the Bryologist*, 1900, n. 3c. u.

1. In silvat. palud. — *Am. sept.* : New-York.

occidentale (*Hpe.*) *Jaeg. op. cit.* p. 392.

Hypnum occidentale *Hpe. in Ann. sc. nat. V ser.*, p. 314 (1866) *Mitt. M. austr. am.* p. 559 (1869).

Saxic. — *Am. merid.* : And. Nov. Granat. et Quitens.

œdipodium (*Mitt.*) *Jaeg. op. cit.* p. 396.

*Br. curtum Ren. et Card. in Rev. bryol.* 1898, p. 19.

Eurhynchium œdipodium *Kindb. Eur. a. N. Am. Bryin.* p. 100 (1897).

Hypnum curtum *Lindb. M. scand.* p. 35 n. 95 (1879)?

H. œdipodium *Mitt. in Journ. of the Linn. Soc. VIII*, p. 35, t. 5 (1864);  
*Lesq. et Jam. Man.* p. 342 (1884).

1. — *Eur.* : Fennia. *Am. sept.* : Terra nov., ins. Princ.  
Edouard, New Brunswick, Ottawa, Montana, M. Rupestr., Columb.  
brit., Washington.

œdistegum (*C. M.*) *Jaeg. op. cit.* p. 410.

Hypnum œdistegum *C. M. Syn. II*, p. 350 (1851).

H. plumosum *Schwaegr. (fid. Mitt. M. Ind. or.* p. 79).

H. pseudo-plumosum? *Mont. in Ann. sc. nat.* 1842, *Crypt. Neilgherr.* n. 12.

1. — *As.* : Yunnan, Himalaya, M. Neilgherr.

*olympicum* Jur. — *Br. venustum.*

*ornellatum* Vent. et Bott. — *Scleropodium.*

orthopyxis *Rehm. M. Afr. austr. n.* 377.

— *Afr.* : Austr.

oxycladon (*Brid.*) *Jaeg. op. cit.* p. 388.

*Br. lætum forma (fid. Card. in Bull. hb. Boissier*, 1899, p. 332).

Camptothecium oxycladon *Kindb. op. cit.* p. 88.

Hypnum attenuatum *Brid. Bryol. univ. II*, p. 448 (1827).

H. oxycladon *cj. Sp. M. II*, p. 128 (1812); *C. M. Syn. II*, p. 368 (1851);  
*Lesq. et Jam. Man.* p. 411 (1884).

1. — *Am. sept.* : Canada, Pensilv.

*oxyrhynchum* (Dz. Mk.) Jaeg. op. cit. p. 387.

Br. melanocarpum Jaeg. op. cit. p. 392.

Hypnum melanocarpum C. M. Syn. II, p. 392 (1851).

H. oxyrhynchum Dz. Mk. in Ann. sc. nat. 1844, p. 308; C. M. op. cit. p. 451; Bryol. jav. II, p. 163, t. 262 (1861-70).

H. velutinum Duby in Pl. Zoll. n. 1809 et in Moritz Syst. Verz. p. 131 ( ).

1. — As. : Java.

*paradoxum* (Hook. f. et Wils.) Jaeg. Ad. II, p. 405 (1875-76).

Hypnum paradoxum H. f. et W. in Lond. Journ. of bot. 1844, p. 554 et Fl. antarct. II, p. 419, t. 155, f. 2 (1847); C. M. Syn. II, p. 324 (1851) et in Fl. 1885, p. 426; Mitt. M. austr. am. p. 563 (1869).

1. Terr. — Afr. : Ins. Kerguelen. Am. merid. : Patagon., Fuegia, ins. Eremitae. Pac. : Austral., Nov. Seeland., Tasman.

*parvulum* Dusen M. Am. austr. n. 763.

Terr. — Am. merid. : Chile austr.

*Payotianum* W. P. Sch. mss.; Boulay, M. Fr. I, p. 135 (1884); Husn. Muscol. gall. p. 426, t. 124 (1894).

Br. trachypodium forma (id. Limpr.)

Eurhynchium Payoti Kindb. Eur. a. N. Am. Bryin. p. 100 (1897).

1 (Limpr.), 2 (Boul.). In fiss. rup. umbr. — Eur. : Alp. Sabaud. (M. Blanc.)

*Peckei* Roumeg. — Eurhynchium praelongum.

*perminusculum* C. M. in Nuov. Giorn. bot. ital. 1898, p. 200.

— As. : China centr. (prov. Schen-Si).

*permolle* C. M. in op. cit. 1896, p. 226.

— As. : China centr. (prov. Schen-Si).

*perpiliferum* C. M. in op. cit. 1896, p. 125 et 1898, p. 199.

— As. : China centr. (prov. Schen-Si).

*piliferum* Kindb. — Eurhynchium.

*pinnirameum* C. M. in op. cit. 1896, p. 126.

Saxic. secus rivul. — As. : China centr. (prov. Schen-Si).

*planiusculum* C. M. in op. cit. 1897, p. 268.

2. — As. : China centr. (prov. Schen-Si).

*platycladum* C. M. et Kindb. Bidrag p. 453 (nom., 1890) et in Mac. Cat. p. 19 (1892).

H. rutabulum Mac. Canad. M. n. 287 p.p.

2. Saxic. — Am. sept. : Washington, Oregon.

*plicatum* Bryol. eur. — Ptychodium.

*plumeum* C. M. n Dusen M. Camer. n. 297.

— Afr. : Cameroon.

*plumosiforme* W. P. Sch. mss.

Hypnum plumosiforme C. M. in Bot. Zeit. 1858, p. 171.

1. — Afr. : Pr. B. Sp.

*plumosum* (Sto.) Bryol. eur. fasc. 52-54 (Vol. VI), Mon. p. 4, t. 597 (1853); W. P. Sch. Syn. II ed. p. 657 (1876).

Br. Maximilianum Gumb. Moossfl. v. Rheimpflz. p. 87 (1857).

Br. pseudo-plumosum Brokm. Laubm. Mecklenb. p. 123 (1870).

Br. Molendoi Ltz. in Sauter Fl. Herzogthum Salzburg. III, p. 60 (1870).

Hypnum alpinum Sm. Fl. brit. III, p. 1330 (teste Wils.).

H. asperulum Brid. Sp. M. II, p. 169 (1812).

- H. asprellum* *Schwaegr. Suppl. I, P. II, p. 246* (1816); *Brid. Bryol. univ. II, p. 480* (1827).
- H. chrysostomum* *Rich. in Mich. Fl. Am. bor. p. 319* (1803); *Brid. Mant. M. p. 173* (1819).
- H. flagellare* *Hedw. Sp. M. p. 282, t. 73 f. 1-3* (1801).
- H. hygrophilum* *Brid. in sched.*
- H. laevisetum* *Crome in Mag. S. Naturfr. V Jahrg. p. 78, t. 3, f. 5-13.*
- H. oedistegum* *C. M. (fid. Mitt. M. Ind. or. p. 79).*
- H. plumosum* *Sic. in Act. Holm. 1795, p. 256.*
- H. polyrhizon* *Brid. Sp. M. II, p. 176* (1812).
- H. pseudo-plumosum* *Brid. Mant. M. p. 170* (1819) *et Bryol. univ. II, p. 472* (1827); *C. M. Syn. II, p. 350* (1851).
- H. pseudo-plumosum*  $\left\{ \begin{array}{l} \beta \text{ asperulum} \\ \gamma \text{ polyrhizon} \\ \delta \text{ chrysostomum} \end{array} \right\}$  *Brid. Bryol. univ. II, p. p. 475 et 489.*
- H. saxatile* *ej. Musc. rec. II, P. II, p. 153* (1798).
- H. secundum* *Mont. Syll. p. 9* (1856) *(fid. Mitt. l. c.).*
- Leskea plumosa* *Sw. M. succ. p. 66* (1).
1. *Ad saxa, rup. et mur. irrig., in et secus rivul. form. quartz. — Eur. : Spitzberg., zon. sept. et interm., raro merid. As. : Japon., Saghalien, China or., Jenissei sup. et med., Kashmir, Tchitral, Himalaya, Nepal., Khasia, M. Neilgherr., Java, Ceylan, Caucas. occ. et centr. Afr. : Alger., Madeir. Am. sept. : Groenland, Terra nov., ins. Miquelon, Canada, Alaska, M. Rupestr., Columb. brit., reg. sept., centr. et occ. Am. merid. : And. Nov. Granat. et Quitens. Pac. : Nov. Seeland.*
- « *var. aquaticum* (*Funck*) *Walther et Mol. Laubm. Ober. Frank. p. 171* (1868).  
*Hypnum aquaticum* *Funck Moostaschb. p. 61, t. 44* (1820).  
*Ad Cataract. — Eur. : Fichbelgeb.*
- « *var. denticulatum* *Del. Fl. cryp. Belg. M. p. 245* (1883).  
*— Eur. : Belg.*
- « *var. homomallum* *Bryol. eur. l. c*  
*Ut et ubi typ. in Eur. et Am. sept.*  
*var. homomallum* *Milde. — Br. amœnum.*
- « *var. julaceum* *Breidl. Laubm. Steierm. p. 185* (1891).  
*Raro fruct. — Eur. : Stir. (ster.), alp. Sabaud. (c. fr. !).*
- « *var. obtusiusculum* (*Broth.*) *Par.*  
*Hypnum pseudo-plumosum var. obtusiusculum* *Broth. M. Turkestan, n. 137*  
*— As. : Turkestan.*  
*var. reflexum* *Aust. — Br. populeum.*
- polygamum* *Warnst. in Allgem. bot. Zeit. n. 10, Sep. Abdr. p. 76* (1899).  
*Polyg. Terr. in dumet. — Eur. : Boruss.*
- populeum* (*Hedw.*) *Bryol. eur. fasc. 52-54 (Vol. VI), Mon. p. 3, t. 535* (1853); *W. P. Sch. op. cit. p. 536.*  
*Br. plumosum var. reflexum* *Aust.*  
*Br. populeum var. angustifolium* *Kindb. in Rev. bryol. 1892, p. 103.*  
*Br. viride* *Brockm. Laubm. Mecklenb. p. 123* (1870).  
*Hypnum concinnum* *Hartm. Skand. Fl. V ed. p. 328* (1849).  
*H. erythropodium* *Brid. Sp. M. II, p. 281 ?* (1812).

(1) De syn. cf. W. P. Sch. Syn. l. c.

- H. implexum* Sw. in *Act. Holm.* 1795, p. 264.  
*H. Laureri* C. M. *Syn.* II, p. 401 (1851).  
*H. plumosum* Sw. *Disp. M. Suec.* p. 66 (1799).  
*H. polysetum* Bland. in *sched.*  
*H. populeum* Hedw. *Sp. M.* p. 270, t. 70, f. 1-6 (1801).  
*H. populeum*  $\gamma$  *erythropodium* Brid. *op. cit.* p. 471 ?  
*H. populeum*  $\beta$  *nutans* ej. *op. et l. cc.* ?  
*H. saxicola* Voit in *Sturm Deutsch. Fl. II. fasc.* 12 c. ic. (1812).  
*H. viride* Lam. *Enc. Meth. bot. III*, p. 181 (1789) *fid. Brockm. Laubm. Mecklenb.* (1869).  
*H. Stereodon Laureri* Funck in *Brid. Bryol. univ. II*, p. 595 (1827).  
 1. Ad arbor. et saxa, rar. ad. terr. — *Eur.* : Fere tota in reg. campestr. *As.* : Japon, Jenissei super., Kashmir, Caucas. *Am. sept.* : Ins. Miquelon, Nov. Scot., New Brunswick, Gaspesia, Ontario, reg. or. ; M. Rupestr.  
*var. amœnum* Limpr. — *Br. amœnum*.  
 \* *var. angustifolium* Besch. in *Ann. sc. nat.* 1893, p. 376.  
     — *As.* : Japon.  
*var. angustifolium* Kindb. — *Br. populeum*.  
 \* *var. attenuatum* Bryol. *eur. l. c. t.* 536  $\epsilon$ ; W. P. Sch. *op. et l. cc.*  
 Ad arbor. — *Eur.* : Ubi typ. *As.* : Japon. *Afr.* : Alger.  
 \* *var. kominaticum* Besch. in *op. et l. cc.*  
     — *As.* : Japon.  
 \* *var. longisetum* Bryol. *eur. l. c. t.* 536  $\gamma$ ; W. P. Sch. *op. et l. cc.*  
 In umbros. humid. ad saxa et terr. — Ubi typ. in *Eur.* et *Am. sept.*  
 \* *var. majus* Bryol. *eur. l. c. t.* 536  $\beta$ ; W. P. Sch. *op. et l. cc.*  
 Ad saxa madida. — Ubi typ. in *Eur.* et *Am. sept.*  
 \* *var. ovatum* Grout in *Bull. Torr. bot. Club*, 1897.  
     — *Am. sept.* : Absque loco spec.  
*var. petrophilum* Ltz. — *Br. populeum var. rufescens*.  
 \* *var. rufescens* Bryol. *eur. l. c. t.* 536  $\zeta$ ; W. P. Sch. *op. et l. cc.*  
*Br. populeum var. petrophilum* Ltz. *Bryol. Notizb.* p. 68 (1865).  
*Hypnum petrophilum* Funck in *sched.*  
 Ad saxa quartz. in silvat. mont. — *Eur.* : Passim.  
 \* *var. subfalcatum* Bryol. *eur. l. c. t.* 536  $\delta$ ; W. P. Sch. *op. et l. cc.*  
 Ad arbor. — *Eur.* : Ubi typ. *Afr.* : Alger. *Am. sept.* : Ubi typ.  
*praelongum* W. P. Sch. in *Mandon Pl. Boliv. n.* 1704; C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 155.  
     — *Am. merid.* : And. Boliv.  
*prasophyllum* (Hpe) Jaeg. *Ad. II*, p. 394 (1876-77).  
*Hypnum prasophyllum* Hpe. in *Ann. sc. nat. V ser.*, p. 315 (1866).  
     — *Am. merid.* : And. Nov. Granat.  
*Preussii* (Broth.) Par.  
*Hypnum Preussii* Broth. in *Engler's Bot. Jahrb.* 1894, p. 214.  
 2. — *Afr.* : Cameroon.  
*procumbens* (Mitt.) Jaeg. *op. cit.* p. 409.  
*Hypnum procumbens* Mitt. *M. Ind. or.* p. 79 (1859).  
 2. — *As.* : Kashmir, Nepal, M. Neilgherr.

- pseudo-albicans* *Kindb. in Bull. Torr. bot. Club. XVII, p. 278 (1890) et in Mac. Cat. p. 194 (1892).*  
*Hypnum albicans* *Mac. Canad. M, n. 235 p.p.*  
 Ster. Arenic. — *Am. sept.* : Ins. Vancouver.
- pseudo-collinum* *Kindb. in Mac. op. cit. p. 196.*  
 1. *Ad puteum.* — *Am. sept.* New Brunswick.
- pseudo-erythrorhizon* *Kindb. in Hedwig. 1896, p. 68.*  
 — *Am. sept.* : Oregon.
- pseudo-laetum* *C. M. in Hedwig. 1898, p. 260.*  
*Hypnum laetum* *Sull. in Proced. Am. Acad. .... 1861, p. 289.*  
 2. *Rupic. in rivul.* — *Am. sept.* : Cuba.
- pseudo-plumosum* *Brockm. — Br. plumosum.*
- pseudo-populeum* *W. P. Sch. mss.*  
*Hypnum pseudo-populeum* *C. M. in Bot. Zeit. 1858, p. 170.*  
 2. — *Afr.* : Prom. B. Sp.
- pseudo-rutabulum* (*Hpe.*) *Jaeg. op. cit. p. 398.*  
*Hypnum pseudo-rutabulum* *Hpe. in Linn. 1863, p. 160; Mitt. M. austr. am. p. 562.*  
 Lignic. et terr. — *Am. merid.* : And. Nov. Granat. et Quitens.
- pseudo-Starkii* *Ren. et Card. in Bot. Centralbl. 1890, n. 51 et in Hedwig. 1893, p. 263.*  
*Br. curtum* (*fid. Limpr.*).  
 2. — *Am. sept.* : Washington.
- pseudo-velutinum* (*Hpe.*) *Jaeg. op. cit. p. 391.*  
*Hypnum pseudo-velutinum* *Hpe. in Bot. Zeit. 1858, p. 171.*  
 1. — *Afr.* : Pr. B. Sp.
- pulchrirete* *C. M. in Rehm. M. Afr. austr. n. 143 et in Hedwig. 1899, p. 138.*  
 2? — *Afr.* : Pr. B. Sp.
- pusillo-albicans* *C. M. in Bull. hb. Boissier, 1897, p. 218.*  
 Ster. Cortic. — *Am. sept.* : Guatemala.
- purum* *Dixon. — Hypnum.*
- reflexum* (*Starke*) *Bryol. eur. fasc. 52-54 (Vol. VI), Mon. p. 8. t. 539 (1853); W. P. Sch. Syn. II ed. p. 650 (1876).*  
*Claopodium laxifolium* *Card. M. Am. sept. p. 150 (fid. ips.).*  
*Eurhynchium laxifolium* *Jaeg. op. cit. p. 426.*  
*E. reflexum* *Kindb. Eur. a. N. Am. Bryin. p. 101 (1897).*  
*Hypnum grimsulanum* *Bryol. eur. l. c. p. II in obs. (fid. C. M.).*  
*H. laxifolium* *Schwaegr. Suppl. II, P. II, p. 159, t. 143 (1826); C. M. Syn. II, p. 449 (1851); Lesq. et Jam. Man. p. 329 (1884).*  
*H. reflexum* *Starke in W. et M. Bot. Taschb. pp. 306 et 476 (1807); Brid. Bryol. univ. II, p. 461 (1827); C. M. Syn. II, p. 448 (1851).*  
*H. subtenue* *Jam. in Proced. Acad. Philad. 1855, p. 447.*  
*Isothecium laxifolium* *Brid. op. cit. p. 359.*  
*Leskea latifolia* *Hook. M. ex. t. 30 (1818-20).*  
*Rigodium reflexum* *Kindb. Laubm. Schewd. u. Norw. p. 14 (1883).*  
*Thuidium laxifolium* *Mac. Cat. p. 187 (1892); Par. Ind. p. 1281 (1894).*  
 1. *Ad Fagorum debil. trunc. et ram.* — *Eur.* : Reg. montana super. et subalp. As. : Saghalien, Amour, Jenissei med. et inf., Kashmir,

Caucas. *Am. sept.* : Groenland, Labrador, Anticosti, Terra nov., ins. Miquelon, Canada, de Alaska ad Calif., Missouri, New Hampshire, Nov. Angl., Maryland.

« *var. Demetrii Ren. et Card. in Bot. Gaz.* 1894, p. 239.

— *Am. sept.* : Labrador.

« *var. pacificum Ren. et Card. in Bot. Centralbl.* 1890, n. 51.

*Eurhynchium pacificum Kindb. Eur. a. N. Am. Bryin.* p. 101 (1897).

— *Am. sept.* : Alaska, Oregon.

« *var. subglaciale Limpr. Laubm. Deutschl.*.... III, p. 115 (1896).

*Saxic.* — *Eur.* : Riesengeb.

*rhynchostegioides C. M. in Rehm. M. Afr. austr.* n. 375.

— *Afr.* : Austral.

*reflexum* × *populeum Rota Contrib.* 1891, p. 212.

— *Eur.* : Ital. super.

*riparium W. P. Sch. in Mandon Pl. Boliv.* n. 1706.

— *Am. merid.* : And. Boliv.

*rivulare (Bruch) Bryol. eur. fasc.* 52-54, (Vol. VI), *Mon.* p. 13, t. 546 (1853); *W. P. Sch. op. cit.* p. 655.

*Br. flavescens Kindb. op. cit.* p. 118.

*Br. rutabulum var. rivularis M. T. Lge. in Bot. Tidskr.* 1869, p. 30.

*Camptothecium megaptitum Mac. Canad. M. n.* 439.

*Hypnum chrysostomum C. M. Syn. II.* p. 368 (1851).

*H. flavescens Brid. Sp. M. II.* p. 185 (1812).

*H. rivulare Br. in sched.* ; *Wils. Bryol. brit.* p. 346, t. 55 (1855).

*H. rutabulum Drumm. M. bor. am. I ed.* n. 178.

*H. rutabulum var. flavescens Brid. Bryol. univ. II.* p. 4<sup>8</sup> ? (1827).

*H. rutabulum var. laxifolium ej. l. c. ?*

*H. rutabulum var. rivulare Th. Jens. Bryol. dan.* p. 177 (1856).

*H. vagans Drumm. Coll. cit.* n. 166.

2. *Ad scaturig. silvat. saxa et mnr. irrig.* — *Eur.* : Ins. Ursor.; reg. campestr. montosa. *As.* : Ins. Kuril., Japon., Jenissei sup., Panjab, N. W. Himalaya, Kashmir, Turkestan, Alatau, Caucas. occ. et centr. *Afr.* : Madeir. *Am. sept.* : Groenland, Labrador, fret. Hudson et Kotzebue, ins. Miquelon, New Brunswick, Ontario de Alaska ad Calif., Wyoming, Minnesota, Idaho, Montana, Wisconsin, Pensilv.

« *var. cataractarum Sauter Fl. d. Herzogthum Salz. III.* p. 60 (1876). *Ad catar. et torr. Alp.* — *Eur.* : Riesengeb., Stir., Salisb., Tirol.

« *var. chrysophyllum Spr. mss.; Dix. et Jam. Stud. Handb.* p. 404 (1896). — *Eur.* : Anglia.

« *var. fallax M. T. Lge. op. et l. cc.*

— *Am. sept.* : Groenland.

« *var. fluitans Lamy et Husn. Muscol. gall.* p. 322 (1893).

*Aquatic.* — *Eur.* : Gall. centr.

*var. latifolium Husn.* — *Br. latifolium.*

« *var. nitidum Sauter in op. et l. cc. (forma ! fid. Limpr.)*

*Terr. in silv.* — *Eur.* : Salisb.

« *var. obtusulum Kindb. in Mac. Cat.* p. 201 (1892).

*Br. rivulare\* Novae Brunswiciae ej. in Ott. Natur.* IV, p. 3 (1889); *Mac. Canad. M. n.* 492.

*Ad rup. madid.* — *Am. sept.* : New Brunswick, Ontario, Columb. brit.

- « *var. rugulosum* Warnst. in *Abh. d. bot. Verein d. prov. Brandenb.* 1900, p. 218.  
In pratis humid. ferrug. — *Eur.* : Boruss. or.
- « *var. subsphaericarpon* de Not. *Cron. II*, p. 18 (1867) *et Epil. p. III* (1869).  
*Hypnum rutabulum var. subsphaericarpon* ej. in *Syll. N.* 32 β (1838).  
— *Eur.* : Lac. Verbanum.
- « *var. tenue* Dixon *Stud. Handb.* p. 404 (1896).  
— *Eur.* : Anglia.
- « *var. turgescens* Warnst. in *Verh. d. K.K. zool. bot. Getellsch. in Wien*, 1900.  
Ad Cataract. — *Eur.* : Tirol. merid.
- « *var. umbrosum* H. Müll. *Westf. M.* n. 315.  
In silv. — *Eur.* : Westphal.
- rivulare* \* *Novae Brunsviciae* Kindb. — *B. rivulare var. obtusulum.*  
*rivulariforme* W. P. Sch. in *Mendon Pl. Boliv. n.* 1705.  
— *Am. merid.* : And. Boliv.
- rivularioides* (C. M.) *Par. Ind.* p. 142 (1894).  
*Hypnum rivularioides* C. M. in *Linn.* 1878-79, p. 451.  
Ster. Lignic. — *Am. merid.* : Argentin. Cordob.
- Roellii* Ren. et Card. in *Bot. Centralbl.* 1890, n. 51 *et in Hedwig.* 1893, p. 263.  
*Eurhynchium Roellii* Kindb. *Eur. and. N. Am. Bryin.* p. 101 (1897).  
2? ♀ et fr. ign. — *Am. sept.* : Ins. Vancouver.
- Rotaeaneum* de Not. *Cron. II*, p. 19 (1867) *et Epil. p.* 117 (1869).  
*Br. sabbrosium var. cylindricum* *Bryol. eur. fasc.* 52-54 (*Vol. VI*), *Mon* p. 16, t. 16 ε (1853); *W. P. Sch. op. cit.* p. 641.  
*Hypnum capillaceum* *Starke in W. et M. Taschenb.* p. 313 (1807).  
*H. salebrosium var. capillaceum* *W. et M. op. et l. cc.*  
l. Ad rad. arb. — *Eur.* : Brandeb., Siles., Stir., Ital. super. (Bergam.)  
*As.* : Japon. *Am. sept.* : Passim.
- « *var. cylindroides* *Limpr. Laubm. Deutschl.*.... *III*, p. 71 (1896).  
Ad muros et saxa. — *Eur.* : Salisb., Tirol. merid.
- ruginosum* (Mitt.) *Par. Ind.* p. 142 (1894).  
*Hycomium ruginosum* *Mitt. in Trans. of the Linn. Soc.* 1891, p. 178.  
2? — *As.* : Japon., Simla.
- rutabuliforme* *Kindb. in Mac. Cat.* p. 198 (1892).  
l. Saxic. — *Am. sept.* : Columb. brit.
- rutabulum* (L.) *Bryol. eur. l. c. Mon. p. II*, t. 543; *W. P. Sch. op. cit.* p. 653.  
*Hypnum crenulatum* *Sm. Fl. brit. II*, p. 1289.  
*H. dentatum vulgatissimum*, operculis obtusis *Dill. Hist. M.* p. 295, t. 38, f. 29 (1741).  
*H. repens triangularibus foliis angustioribus* ej. *Cat. Giss. p.* 218 (1718).  
*H. rutabulum* *L. Sp. pl. I ed. II*, p. 1124 (1753) *non herb. ! Hedic. Descr. IV*, p. 29, t. 12. 1794; *Brid. Bryol. univ. II*, p. 485 (*incl. var. aureo-nitens, crenulatum, exiguum, flaccidum*, p. 487); *C. M. Syn. II*, p. 367 (1851).

l. Ad saxa, terr. umbr., rad. arb. lignaq. putresc. — *Eur.* : Tota in reg. campestr., plan. et montos., praeter arctic. *As.* : Japon., Jenissei super., Kashmir, Tschitral, Himalaya et Tibet. occ., Pers., Caucas., Syr., m. Olymp. *Afr.* : Alger. *Am. sept.* : Terra nov., ins. Miquel., Canada or., ins. Vancouver. Washington, reg. sept., centr. et or. *Am. merid.* : And. Quitens. *Pac.* : Nov. Seeland., Tasman., ins. Campbell.

« *var. apuanum* Vent. et Bott. *Enum. crit.* p. 8 (1884).  
— *Eur.* : Alp. Ital. super.

« *var. aureo-virens* (Brid.) Brockm. *Laubm. Mecklenb.* p. 122 (1870).  
Hypnum aureo-virens Brid. *Sp. M. II.* p. 184 (1812).  
*H. rutabulum var. aureo-virens* ej. op. et l. cc.  
— *Eur.* : Mecklenbourg.

« *var. brevisetum* Fiedl. *Laubm. Mecklenb.* p. II (1844).  
Terr. ad rip. fluv. — *Eur.* : Mecklenbourg.

« *var. canadense* Ren. et Card. in *Rev. bryol.* 1893, p. 19.  
— *Am. sept.* : Ins. Miquel., Canada, Washington.

« *var. cavifolium* Lindb. in *Oesv. af K. Vet.-Akad. Foerh.* 1866, p. 539.  
— *Eur.* : Spitzberg.

*var. dendroideum* M. T. Lge. — Br. rivulare.

« *var. densum* Bryol. *eur. l. c. t.* 544 ε; W. P. Sch. op. cit. p. 654.  
Ad trunc. semiputr. — Hic illic cum typo in *Eur.* et *Am. sept.*

« *var. eurhynchioides* Limpr. op. cit. p. 109, fig. 371.  
Ad rip. fluv. — *Eur.* : Brandeb.

*var. explanatum* Boeckm. — Br. curtum.

« *var. flavescens* Bryol. *eur. l. c. t.* 544 γ; W. P. Sch. op. et l. cc.  
In graminosis ad terr. — *Eur.* : Hic illic cum typo in *Eur.* et *Am. sept.*

« *var. Klinggraeffii* (Klinggr. 1893, in *Leb.-u. Laubm. West.-u. Ostpr.* p. 251 in *adn.*).  
Polyg. Ad saxa. errat. — *Eur.* : Boruss. or.

« *var. longisetum* (Brid.) Bryol. *eur. l. c. t.* 544 β; W. P. Sch. op. et l. c.

Hypnum rutabulum *var. longisetum* Brid. *Bryol. univ. II.* p. 487 (1821).  
In gramin. ad terr. — Hic illic cum typo in *Eur.* et *Am. sept.*

« *var. lutescens* Warnst. in *Verh. d. bot. Verein d. prov. Brandenb.* 1899, p. 73.  
— *Eur.* : Lusat. infer.

« *var. plumulosum* Bryol. *eur. l. c. t.* 544 δ; W. P. Sch. op. et l. cc.  
Ad trunc. semiputrid. — Hic illic cum typo in *Eur.* et *Am. sept.*  
*var. rivularis* M. T. Lge. — Br. rivulare.

« *var. robustum* Bryol. *eur. l. c. t.* 544 ξ; W. P. Sch. op. et l. cc.  
Ad rad. arb. silvat. et saxa umbr. — *Eur.* : Ubi typ. *As.* : Caucas. occ. *Am. sept.* : Ubi typ.

« *var. turgescens* Limpr. op. et l. cc.  
Secus semit. in silv. — *Eur.* : Stiria.



« *var. undulatum* Warnst. in *Allgem. bot. Zeit.* 1899, *Beih. I*, p. 10.  
— *Eur.* : Longobard.

« *var. viviparum* Bryhn De Bryneear. in *Norveg. p.* 21 (1891).  
*Aquat.* — *Eur.* : Norvège.

*var. 4. H. f. et W.* — *Br. salebrosum.*

Ryani Kaur. in *Bot. Not.* 1888, p. 177; *Philib. in Rev. bryol.* 1889, p. 59.  
2 et ps. 1. *Ad terr. gramin.* — *Eur.* : Norveg. merid.

*salebrosum (Hoffm.) Bryol. eur. fasc. 52-54 (Vol. VI), Mon. p. 16,*  
*t. 549 (1853); W. P. Sch. op. cit. p. 641.*

*Br. ambiguum de Not. Cronaca II, p. 20 (1867) et Epil. p. 112 (1869) ftd.*  
*Vent. et Bott. Enum. p. 9 (1884).*

*Br. campestre Mitt. in Fl. Tasman. II, p. 210 (1860) ex ips. !*

*Br. jucundum de Not. op. cc. pp. 20 et 118 (ftd. Vent. et Bott. l. c.).*

*Br. laevisetum Kindb. in Bull. Torr. bot. Club. 1890, p. 279 et in Mac.*  
*Cat. p. 193 (1892).*

*Br. plumosum Brockm. Laubm. Meckl. p. 120 (1870).*

*Hypnum fasciculatum Lam. Encycl. meth. bot. III, p. 177 ( ).*

*H. Lamarckii Brid. Sp. M. II, p. 256 (1812).*

*H. plumosum Huds. Fl. angl. 1 ed. p. 423 n. 18 (1762) ftd. Lindb. ; Brid.*  
*Bryol. univ. II, p. 475 (1827).*

*H. plumosum β salebrosum C. M. Syn. II, p. 359 (1851).*

*H. repens filicinum pennatum Dill. Cat. Giss. n. 218 (1718).*

*H. repens filicinum plumosum ej. Hist. M. p. 200, t. 35, f. 16 (1741).*

*H. salebrosum Hoffm. Deutsch. Fl. II, p. 74 (1796).*

*H. salebrosum*  $\left\{ \begin{array}{l} \gamma \text{ apiculatum} \\ \delta \text{ paraphysigerum} \\ \beta \text{ Torreyanum} \end{array} \right\}$  *Brid. cit. p.p. 478 et 479.*

1. *In terr. humos. silvat., ad saxa, rar. arb. et trunc. putresc.* — *Eur.* :  
*Ins. Ursor.; tota. As. : Japon., Amour, Saghalien, Jenissei, Obi,*  
*Kashmir, Tchitral, Panjab, Himalaya et Tibet occ., Caucas. Afr. :*  
*Alger., Canar., Azor., ins. Kerguelen. Am. sept. : Groenland,*  
*Terra nov., Canada, fret. Kotzebue, Alaska, Columb. brit., ins.*  
*Vancouver, M. Rupestr., reg. sept., occ., centr. et or. Pac. :*  
*Tasman.*

« *var. angustifolium* Warnst. in *Allgem. bot. Zeit. n. 7-8, Beih. n. 1.*  
*Sp. Abdr. p. 9 (1894).*

*In dumetis.* — *Eur.* : Württemberg.

« *var. arcticum* Berggr. *M. Spetsb. n. 16 et in K. Sv. Vet.-Akad.*  
*Handl. XIII, n. 7, p. 79 (1874).*

*Hypnum plumosum var. arcticum Arn. in Oef. af K. Vet.-Akad. Foerh.*  
*p. 116 (1900).*

— *Eur.* : Spitzberg., ins. Parry.

*var. cylindricum* Bryol. eur. — *Br. Rotaezanum.*

« *var. densum* Bryol. eur. l. c. t. 642  $\gamma$ ; *W. P. Sch. op. et l. cc.*  
*Ad arb. rad. et saxa umbr.* — *Eur.* : Cum typo.

« *var. eurhynchoides* Warnst. in *op. et l. cc.*  
*In fiss. rup. basalt.* — *Eur.* : Hassia.

« *var. flaccidum* Bryol. eur. l. c. t. 642  $\delta$ ; *W. P. Sch. op. et l. cc.*  
*In terr. silvat. praecip. zon. sept. in Eur.*

« *var. gracile* Hartm. *Skand. Fl. III ed. p. 309 (1838).*  
*Ad trunc. putrid.* — *Eur.* : Suecia.

- « *var. longisetum* *Bryol. eur. l. c. t. 642* β; *W. P. Sch. op. et l. cc.*  
*H. salebrosum* *Brid. Bryol. univ. II, p. 427* (1827).  
 In salebr. et ad tecta stramin. — Passim cum typ. in *Eur. et Am. sept.*  
*var. palustre* *Bryol. eur.* — *Br. Mildeanum.*
- « *var. paraphylliferum* *de Not. Cronaca II, p. 20* (1867) *et Epil. p. 120*  
 (1869).  
 Ad rad. arb. — *Eur.* : *Lac. Verbanum.*
- « *var. parvicarpum* *Besch. in Ann. sc. nat. 1893, p. 373.*  
 — *As.* : *Japon.*
- « *var. robustum* *Warnst. in Verh. d. bot. Ver. d. Prov. Brandenb.*  
 1899, p. 70.  
 Ad terr. silvat. — *Eur.* : *Roruss.*
- « *var. rostratum* *Besch. in op. et l. cc.*  
 — *As.* : *Japon.*
- « *var. texanum* (*Aust.*) *Par. Ind. p. 144* (1894).  
*Br. texanum* *Aust. in Bull. Torr. bot. Club. VI, p. 445* (1879).  
*Hypnum salebrosum var. texanum* *Lesq. et Jam. Man. p. 337* (1884).  
 — *Am. sept.* : *Texas.*
- « *var. Thomasii* (*Brid.*) *Bryol. eur. l. c. t. 642* ζ; *W. P. Sch. op. et l. cc.*  
*Hypnum capillaceum* β *Thomasii* *Brid. Bryol. univ. II, p. 482* (1827).  
*H. Thomasii* *ej. Mant. M. p. 172* (1819).  
 In loc. aprie. — *Eur.* : *Hic illic.*  
*var. turgidum* *Hartm.* — *Br. turgidum.*
- « *var. Waghornei* *Ren. et Card. in Bot. Gaz. 1894, p. 238.*  
 — *Am. sept.* : *Labrador.*
- « *salicinum* *Bryol. eur. fasc. 52-54 (Vol. VI), Mon. p. 15, t. 548* (1853);  
*W. P. Sch. op. cit. p. 646.*  
*Hypnum salicinum* *Boulay M. Fr. I, p. 134* (1881).  
 1. Ad *Salic.* semi-putrid. et arb. annosior. — *Eur.* : *Vall. Rhen., os*  
*Elbae, Jurass., Delphinat., Corsica; Hispania (Cuença, fid. Roell.).*  
*var. condensatum* *Husn.* — *Br. velutinum var. condensatum.*
- « *saltense* *Hagen in Nyt Mag. for Naturvid. 1900, p. 339.*  
 2. ♂ et fr. ign. — *Eur.* : *Norveg.*
- « *scabripes* *C. M. in Nuov. Giorn. bot. ital. 1897, p. 154.*  
 — *Am. merid.* : *And. Boliv.*
- « *scabrum* (*Mitt.*) *Besch. in hb. Mus. Paris.*  
*Hypnum scabrum* *Mitt. in hb. Griffith. n. 700.*  
 — *As.* : *Assam, Rhotan.*
- « *Schimperi* v. *Klinggr.* — *Br. campestre.*  
*Schlosseri* *Jaeg.* — *Rigodium.*  
*secundum* *Jaeg.* — *Br. plumosum.*  
*semiasperum* *Kindb.* — *Eurhynchium.*
- « *sericeo-virens* (*C. M.*) *Par. Ind. p. 145* (1894).  
*Hypnum sericeo-virens* *C. M. in Fl. 1885. p. 427.*  
 1. *Rupic. et cortic.* — *Am. merid.* : *Fuegia.*

sericeum Warnst. in *Verh. d. bot. Ver. Mark Brandenb.* 1894, p. 24.

Br. salebrosum forma (fid. *Limpr. in op. cit.* III, p. 78).

l. Ad rad. arb. — *Eur.* : Boruss. or., Siles., Tirol. merid.

serpentinum C. M. in *Dusen M. Camer.* n. 276.

— *Afr.* : Cameroon.

silvaticum Warnst. in *Schrift. d. Naturf. Gesells. in Danzig*, 1896, p. 64.

— *Eur.* : Westphal.

*Spegazzinii* Par. — Hypnum.

*splendens* Aust. — Br. biventrosum.

spurio-acuminatum C. M. et Kindb. *Bidrag.* p. 453 (nom., 1890) et in *Mac. Cat.* p. 191 (1892).

Camptothecium spurio-acuminatum Kindb. *Eur. a. N. Am. Bryin.* p. 89 (1897).

l. Lignic. — *Am. sept.* : Ontario.

spurio-albicans C. M. in *Hedwig.* 1897, p. 127.

Ster. — *Am. merid.* : Patagon.

spurio-populeum (Broth.) Par. *Suppl. Ind.* p. 50 (1900).

Hypnum spurio-populeum Broth. in *Act. Soc. sc. Fenn.* 1898, n. 2, p. 36.

l. — *As.* : Kashmir.

spurio-rutabulum C. M. et Kindb. *op. cc. pp.* 453 (nom.) et 197.

l. Ad rad. arb. — *Am. sept.* : Columb. brit., ins. Vancouver.

squalidissimum (C. M.) Par. *Ind.* p. 145 (1894).

Hypnum squalidissimum C. M. in *Linn.* 1878-79, p. 452.

Ster. Rupic. — *Am. merid.* : Argentin. Cordob.

Starkei (Brid.) Lindb. in *Medd. p. Fauna et Fl. fenn.* 1879, p. 5.

Br. Starkei  $\beta$  alpestre *Bryol. eur. l. c. Mon.* p. 10, t. 541  $\beta$ .

Br. Starkei var. robustum W. P. Sch. *Syn. I ed.* p. 540 (1860) et *II ed.* p. 651 (1876).

Eurhynchium Starkei Kindb. *Eur. and N. Am. Bryin.* p. 102 (1897).

Hypnum grimsulanum *Bryol. eur. l. c. Mon. p. II in obs.* (fid. *Limpr.*).

H. reflexum  $\beta$  Starkei Hartm. *Skand. Fl. II ed.* p. 346 (1832).

H. reflexum  $\gamma$  umbratum Myrin *Coroll. fl. ups.* p. 45 pp. (1833).

H. Starkei Brid. *Musc. rec. II, P. II*, p. 107, p. 261 (1801).

H. (Stereodon) Starkei Brid. *Bryol. univ. II*, p. 595, n. 35 (1827).

Stereodon Starkei ej. *op. cit.* p. 825.

l l (nec syn., C. M.). Ad terr. lapid., rad. arb. et trunc. putresc. in

silvat. — *Eur.* : Reg. mont. et subalp. per zon. sept. et intermed.

*As.* : Japon., Amour., Jenissei super., Caucas. occ. et centr. *Am.*

*sept.* : Groenland, Terra nov., ins. Miquel., New. Brunswick. Nov.

Angl., Pensilv., Ontario, Vancouver.

var. alpestre *Bryol. eur.* — Br. Starkei.

« var. complanatum *Limpr. op. cit.* p. 100.

— *Eur.* : Stiria.

var. curtum Wanst. { Br. curtum.

f. major Milde }

« var. nipponense (W. P. Sch.) *Besch. in Ann. sc. nat.* 1893, p. 376.

Hypnum nipponense W. P. Sch. in *Savatier M. Japon.* n. 657.

— *As.* : Japon.

- var. *praelongum* W. P. Sch. } — Br. *curtum* p.p.  
 } — Br. *glaciale* var. *Huntianum* p.p.  
 var. *robustum* } Limpr. — Br. *curtum*.  
 } W. P. Sch. — Br. *Starkei*.
- stereopoma* (Spr.) Jaeg. *Ad. II*, p. 393 (1876-77).  
 Hypnum *stereopoma* Spr. in *Mitt. M. austr. am.* p. 561 (1869).  
 Lignic. et terr. — *Am. sept.* : Jamaic. *Am. merid.* : Venezuela,  
 And. Quitens. et l'eruv.
- stricto-patens* C. M. in *Hedwig*. 1899, p. 137.  
 2. — *Afr.* : Transvaal.
- subalbicans* de Not. *Epil.* p. 118 (1839); W. P. Sch. *op. cit.* p. 645.  
 1. In silv. — *Eur.* : Pedemont. (Taurinum).
- suberythrorhizon* Ren. et Card. in *Bot. Gaz.* 1893, p. 238, t. XXII, f. B.  
 1. — *Am. sept.* : Colorado.
- subfalcatum* Ren. et Card. in *Bull. hb. Boissier*, 1895, p. 241 et in *Bull.*  
*Soc. roy. bot. Belg.* 1895, I, p. 75.  
 — *As.* : Bhotan.
- subfalcatum* Warnst. — Br. *Warnstorffii*.
- subintricatum* Kindb. in *Rev. bryol.* 1895, p. 86.  
 Cortic. — *Am. sept.* : Ins. Vancouver.
- subpilosum* (Hook. f. et Wils.) Jaeg. *Ad. II*, p. 410 (1876-77).  
 Br. *subplicatum* Jaeg. *op. cit.* p. 399.  
 Hypnum *subpilosum* var. 4 H. f. et W. *Fl. antarct. II*, p. 418 (1847).  
 H. *subpilosum* eor. in *Lond. Journ. of bot.* 1841, p. 558 et *Fl. antarct. II*,  
 p. 418, t. 15A, f. 4; C. M. *Syn. II*, p. 366 (1851).  
 H. *subplicatum* Hpe. in *C. M. Syn. II*, p. 363.  
 Terr. — *Afr.* : Ins. Kerguelen, Marion. *Am. merid.* : Patag.,  
 Fuegia, ins. Eremitae, Maclov.
- subplicatum* Jaeg. — Br. *subpilosum*.
- subrutabulum* (C. M.) Jaeg. *op. cit.* p. 398.  
 Hypnum *subrutabulum* C. M. in *Bot. Zeit.* 1856, p. 455.  
 2. — *Afr.* : Pr. B. Sp.
- subtilicaule* Dusen M. *Am. austr. n.* 421 et 422.  
 Ad lign. putrid. — *Am. merid.* : Patagon. occid.
- sulfurascens* C. M. — Br. *afro-salebrosum*.
- sulfureum* (Geh. et Hpe.) Par. *Ind.* p. 146 (1894).  
 Hypnum *sulfureum* Geh. et Hpe. in *Fl.* 1881.  
 1. Ad terr. et rad. arb. — *Am. merid.* : Brasil. austr. or.
- Sullivantii* Bryol. eur. — Br. *laetum* var. *dentatum*.
- tarapenae* Besch. in *Ann. sc. nat.* 1894, p. 47.  
 Hypnum *plumosum* Sull. in *Wilkes Exped.*  
 1. et 2. Secus torr. — *Pac.* : Taïti.
- Tauriscorum* Mol. in *Fl.* 1866, pp. 306 et 533; Pfeff. *Bryog. Stud.*  
 p. 76 (1869); Limpr. *op. cit.* p. 123.  
 Br. *glareosum* f. *depauperata* (fid. Limpr.).  
 2. — *Eur.* : Alp. German., Helvet., Vall. Tellin.

- tenui-pinnatum* (C. M.) *Par. Suppl. Ind.* p. 51 (1900).  
*Hypnum tenui-pinnatum* C. M. in *Ule Bryoth. brasil. n.* 196.  
 Terr. — *Am. merid.* : Brasil. or.
- tenui-prostratum* C. M. in *Hedwig.* 1897, p. 128.  
 2. Cortic. — *Am. merid.* : Argentin. subtropic. Tucuman.
- texanum* Aust. — Br. salebrosum var. *texanum*.
- Thedenii* (Hartm.) *Bryol. eur. fasc.* 52-54 (Vol. VI), *Mon.* p. 18, t. 551 (1853); W. P. Sch. *op. cit.* p. 645.  
 Br. albicans var. *Thedenii* Hartm. *Skand. Fl.* X ed. p. 15 (1871).  
*Hypnum albicans* var. *Thedenii* cf. *op. cit.* IX ed. (1861).  
*H. erythrorhizon* var. *Thedenii* Lindb. *M. scand.* p. 36 (1878).  
*H. Thedenii* Hartm. *Skand. Fl.* VI ed.
2. Ad saxa umbrosa. — *Eur.* : Suecia, Fennia. *As.* : Jenissei sup.  
*Am. sept.* : New Hampshire.
- thraustum* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 270.  
 — *As.* : China centr. (prov. Schen-Si).
- trachypodioides* W. P. Sch. in *Mandon Pl. Boliv. n.* 1707.  
 — *Am. merid.* : And. Boliv.
- trachypodioides* W. P. Sch. (M. Abyss.) — Br. abyssinicum.
- trachypodium* (Funck) *Bryol. eur. fasc.* 52-54 (Vol. VI), *Mon.* p. 7, t. 517 (1853); W. P. Sch. *Syn.* II ed. p. 649 (1876).  
*Eurhynchium trachypodium* Kindb. *Eur. and N. Am. Bryin.* p. 100 (1897).  
*Hypnum populeum* var. *trachypodium* Rab. *Deutsch. Kryptfl.* II, P. III, p. 294 (1848).  
*H. sericeum* Funck Moostasch. p. 62, t. 44 (1820).  
*H. trachypodium* Funck in *Brid. Bryol. univ.* II, p. 766 (1827); C. M. *Syn.* II, p. 367 (1851).  
*H. velutinum* var. *sericeum* Brid. *op. cit.* p. 400 (1).  
*Isothecium trachypodium* (Funck) *Brid. op. cit.* p. 376.
1. In fiss. rup. — *Eur.* : Spitzberg., Alp. Norveg., Salisb., Stir., Carinth., Rhaet., Bavar., Algov., Tirol., Longobard., Delphin., Sabaud.; Arvern. *As.* : Caucas. *Afr.* : Alger. *Am. sept.* : Groenland.
- trochalobasis* C. M. in *Bull. hb. Boissier*, 1897, p. 218.  
 Ster. — *Am. sept.* : Guatemala.
- tromsoënsæ Kaurin* in *Limpr. Laubm. Deutschl.*.... III, p. 93 (1896).  
*Eurhynchium tromsoënsæ* Kindb. *op. cit.* p. 106.  
*Hypnum tromsoënsæ* Arn. et Jens. in *Bihang till K. Sv. Vet.-Akad. Handl.* 1896, n. 10.  
 1. — *Eur.* : Norveg., Stiria.
- truncatum* Besch. in *Ann. sc. nat.* 1893, p. 374.  
 — *As.* : Japon.
- turgidum* (Hartm.) C. Hartm. in *Kindb. Enum.* p. 294 (1888).  
 Br. salebrosum  $\gamma$ ? *turgidum* Hartm. *Skand. Fl.* III ed., II, p. 309 (1838).  
*Hypnum plumosum turgidum* (Hartm.) Lindb. *M. scand.* p. 36 (1879).  
*H. turgidum* Hartm. *Skand. Fl.* V ed. p. 328 (1849).  
 1. rar. fruct. Terr. — *Eur.* : Lappon., Scand., Fennia, Tirol., Tatra? *As.* : Penins. Samoied., Jenissei infer., Alatau. *Am. sept.* : Groenland, Labrador, M. Rupestr.
- umbilicatum* Jur. et Milde *Beitr.*.... p. 600 (1870).  
 1. — *As.* : Kurdistan.

utahense *Jam. in Bot. King's Exped. p.* 409.

*Eurhynchium utahense* *Kindb. op. cit. p.* 105.

*Hypnum utahense* *Lesq. et Jam. Man. p.* 339 (1884).

Syn. Ad rup. arenac. — *Am. sept.* : Montana occ., Utah, Colorado.

vagaus *Milde in Hedwig. 1869, p.* 50 et *Bryol. siles. p.* 331 (1869).

Br. graniticum *Geh. in Fl. 1894, p.* 80.

Syn. Ad rup. arenac. et granit. — *Eur.* : Boruss. or., Tatra, Westphal., Waldeck, Bavar.

Valentini *Besch. Fl. bryol. Réunion, etc. p.* 154 (1880-81).

Ster. — *Afr.* : Ins. Bourbon.

vallium *Sull. et Lesq. M. bor. am. II ed. p.* 84 ( ).

Br. asperimum *Lesq. et Jam. (fid. Jaeg. Ad. II, p.* 398).

Rupic. — *Am. sept.* : Ins. Vancouver. Calif.

Vaucherii *Kindb.* — *Eurhynchium.*

vellereum (*Mitt.*) *Jaeg. Ad. II, p.* 403 (1876-77).

*Hypnum vellereum* *Mitt. in Journ. of the Linn. Soc. 1863, p.* 151.

2. — *Afr.* : Abyss., Cameroon.

velutinum (*L.*) *Bryol. eur. fasc. 52-54 (Vol. VI), Mon. p.* 5, t. 538 (1853); *W. P. Sch. op. cit. p.* 648.

Br. declivum *Jaeg. op. cit. p.* 394.

Br. intricatum var. declivum *Kindb. in Hedwig. 1896, p.* 68.

*Hypnum declivum* *Mitt. in Journ. of the Linn. Soc. VIII, p.* 33, t. 6 (1865).

H. intricatum *Schreb. Spic. Fl. Lips. (1771).*

H. repens trichodes terrestre viridius minus, capitulis tumidioribus cernuis *Dill. Cat. Giss. p.* 216 (1718).

H. velutinum *L. Sp. pl. p.* 1129 (1753); *Hedw. Descr. IV, p.* 70, t. 27 (1797); *Brid. Bryol. univ. II, p.* 457 (1827); *C. M. Syn. II, p.* 399 (1851).

H. velutinum capsulis ovatis cernuis *Dill. Hist. M. p.* 326, t. 42, f. 61 (1711).

H. Stereodon Starkei  $\beta$  tenellus *Brid. op. cit. p.* 596 p.p.

*Leskea velutina* *Schrank Baier. Fl. III, p.* 461 (1789).

l. Ad saxa, terr., rad. arb. truncosq. putresc. — *Eur.* : Tota, praeter arctic. As. : Saghalien, Jenissei super., Kashmi., Pers., Caucas. *Afr.* : Alger. *Am. sept.* : New Brunswick, Gaspe a, Vermont, New-York, Ontario, Idaho, Columb. brit., Washington, Oregon, Calif.

« var. condensatum *Bryol. eur. l. c. t.* 538  $\delta$ ; *W. P. Sch. op. et l. cc.*

Br. salicinum var. condensatum *Husn. Muscol. gall. p.* 330 (1893).

Br. velutinum var. meridionale *W. P. Sch. mss.*; *Venturi in Rev. bryol. 1883, p.* 64.

Ad mur. delaps. et sax. aridiss. — *Eur.* : Praecip. zon. mer. : Sierra Nevada.

« var. fastigiatum *Warnst. in Allgem. bot. Zeit. 1899, Beih. n. 1, Sep. Abdr. p.* 10.

— *Eur.* : Vall. Rhen.

« var. intricatum (*Hedw.*) *Bryol. eur. l. c. t.* 538  $\gamma$ ; *W. P. Sch. op. et l. cc.*

Br. intricatum *Kindb. in Hedwig. 1896, p.* 66.

*Hypnum intricatum* *Hedw. Descr. IV, p.* 73, t. 28 (1797).

H. velutinum var. intricatum *Brid. op. cit. p.* 459; *C. M. Syn. II, p.* 400.

Ad terr. argill. et rup. — *Eur.* : Hic illic cum typo. *Am. sept.* : Canada.

var. *meridionale* W. P. Sch. — Br. *velutinum* var. *condensatum*.

\* var. *praelongum* *Bryol. eur. l. c. t.* 538 β; W. P. Sch. *op. et l. cc.*  
Ad terr. silvat. arenac. — *Eur.* : Cum typo.

var. *venustum* Archangeli. — Br. *venustum*.  
*Venturii* Warnst. — Br. *amœnum*.

*venustum* de Not. *Cronaca II*, p. 21 (1867) et *Epil.* p. 122 (1869); *Vent.*  
in *Rev. bryol.* 1883, pp. 61 et 64.

Br. *olympicum* *Jur. in Unger und Kotschy « die Insel Cypren »* (1865);  
W. P. Sch. *op. cit.* p. 854.

Br. *salicinum* (sec. *Husn. Muscol. gall.*).

Br. *velutinum* var. *venustum* Archangeli in *Nuov. Giorn. bot. ital.* 1889.  
*Hypnum venustum* de Not. *Syll. n.* 42 (1838); *C. M. Syn. II*, p. 397 (1851).

1. (C. M.) nec 2? (de Not.) Lignic. et ad rup. silic. — *Eur.* : Alp.  
Stir., Vall. Tell., Corsica, Pyren., S. de Guadarrama. *As.* :  
M. Olymp., Cypr.

Villardi *Ren. et Card. in Bot. Centralbl.* 1890, n. 51.

*Eurhynchium* Villardi *Kindb. Eur. and N. Am. Bryin.* p. 98 (1897).

1. — *Am. sept.* : Washington.

*vineale* Milde in *Bot. Zeit.* 1864, *Beil.* p. 15, n. 9.

Polyg. Muric. — *Eur.* : Tirol merid.

*viride* Brockm. — Br. *populeum*.

*viridifactum* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 270.

— *As.* : China centr. (prov. Schen-Si).

*Warnstorffii* Par.

Br. *subfalcatum* Warnst. in *Verh. d. Prov. Brandenb.* 1900, p. 72.

-- *Eur.* : Lusat. infer.

*Washingtonianum* Eaton mss; A. J. Grout in *the Bryologist.* 1900,  
n. 3 c. ic.

2? Ad terr. humid. — *Am. sept.* : Washington.

*Weinmanni* (Nees) Par. *Ind. p.* 148 (1894).

*Hypnum Weinmanni* Nees mss.; *C. M. Syn. II*, p. 365 (1851).

Ad arb. — *Eur.* : Rossia merid.

*Wichurae* (Broth.) Par.

*Hypnum Wichurae* Broth. in *Hedwig.* 1899, p. 239.

2. Rupic. — *As.* : China or., Japon.

*Wilkesianum* Jaeg. — Entodon.

*Zickendrathi* Warnst. in *Verh. d. K.K. Zool. bot. Gesells. in Wien*,  
1900, p.

2. Muric. — *Eur.* : Tirol merid.

BRACHYTRICHUM Roehl. in *Ann. d. Wett. Gesells. III*, p. 194 ( ).

Spec. omn. sunt ORTHOTRICHÆ ejusd. nom.

BRAITHWAITEA Lindb. in *Act. Soc. sc. Fenn. X*, 1872, p. 250 in adn.

*arborescens* Mitt. — *Hypnodendron*.

*nematosa* Jaeg. } — *Climacium sulcatum*.  
*sulcata* Lindb. }

**BRAUNFELSIA** Par. Ind. p. 148 (1894). — 5 spec.dicranoides (Dz. et Mk.) Broth. *Natür. Pflanzenfam.* p. 321.Br. Molkenboeri Fleisch. *M. Archip. ind. n.* 9.

Dicranum Molkenboeri Bryol. jav. II, p. 225 (1861-70; Par. Ind. p. 359 1895).

Holomitrium dicranoides Dz. et Mk. *M. frond. ined. Archip. ind. p.* 9, t. 4 (1854; Bryol. jav. I, p. 87 1855-61).

2. Ad rad. arb. — As. : Java.

enervis (Dz. Mk.) Par. op. et l. cc.

Dicranum macrocalyx Bryol. jav. II, p. 225 (1861-70).

Eucamptodon macrocalyx C. M. Syn. I, p. 316 (1849).

Holomitrium enerve Dz. Mk. *M. frond. ined. Archip. ind. p.* 8, t. 3 (1854).Solmsia enervis Hpe. in *Nuov. Giorn. bot. ital.* 1872, p. 273.

2. In humid. — As. : Java.

Molkenboeri Fleisch. — Br. dicranoides.

Petriai (Broth.) Broth. op. et l. cc.

Eucamptodon Petriai ej. in *Oefv. af Finska Vet. Soc. Handl.* 1898, p. 161; Par. Suppl. Ind. p. 144 (1900).

2. fr. ign. — Pac. : Nov. Seeland merid.

plicata (Lac.) Broth. *Natür. Pflanzenfam. l. c.*Dicranum plicatum Lac. Sp. nov. *M. Archip. ind. p.* 9, t. 6 B (1872; Par. Ind. p. 363 (1895).

2. — As. : Java.

scariosa (Wils.) Par. op. et l. cc.

Dicranum elentulum Mitt. *M. Ind. or. p.* 14 (1859).D. scariosum Wils. in *Kew Journ. bot. IX*, p. 294 (1857).Eucamptodon edentulus C. M. in *Linn.* 1869-70, p. 11.

Solmsia scariosa Hpe. in op. cit. p. 274.

2. — As. : Borneo, Ceylan.

« var. angusta C. M. Wright in *Trans. of the Linn. Soc.* 1894-96, p. 256.

— As. : Borneo sept. occ. (m. Kinabalu).

« var. compacta (C. M.) Par. op. et l. cc.

Eucamptodon edentulus var. compactus C. M. in *Jaeg. Ad. I*, p. 151 (1869-70).

Solmsia scariosa var. compacta Jaeg. op. cit. II, p. 652 (1877-78).

Ut et ubi typ.

**BRAUNIA** Bryol. eur. fasc. 29-30 (Vol. III) (1846). — 23 spec.

Spec. omn. non indicatae sunt Hedwigia ejusd. enom.

alopecura (Brid.) Linpr. *Laubm. Deutschl.... I*, p. 824 (1889).Anoetangium sciuroides Bals. et de Not. *Pugill. n.* 23 (1836).

Braunia sciuroides Bryol. eur. fasc. 29-30 (vol. III), Mon. p. 3, t. 275 (1846); W. P. Sch. Syn. II ed. p. 286 (1876).

Gymnostomum ciliatum  $\beta$  nudum Schleich. *Cat.* 1807.Harrisonia sciuroides Rab. *Krypt. Deutsch. Fl. II*, 3, p. 158 (1848).Hedwigia sciuroides de Not. *Syll. M.* p. 95 (1838).Leucodon alopecurus Brid. *Mant. M.* p. 135 (1819) et *Bryol. univ. II*, p. 214 (1827).

Neckera alopecura C. M. Syn. II, p. 104 (1851).

Schistidium nudum Schleich. *Cent. pl. helv.*

1 et Syn. Ad rup. calcar. sicc. — Eur. : Helvet. et Tirol. merid., Pyren., Ital. merid.



- Andrieuxii* Ltz. *Moosst.* p. 164 (1864).  
 l. — *Am. sept.* : Mexic.
- argentinica* C. M. in *Linn.* 1880-82, p. 465.  
 l. Saxic. et terr. — *Am. merid.* : Argentin. subtrop.
- argyrotricha* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 139.  
 — *Am. merid.* : And. Boliv.
- attenuata* (Mitt.) Jaeg. *Ad. II*, p. 88 (1869-70).  
*Hedwigia attenuata* Mitt. *M. Ind. or.* p. 123 (1859).  
 2 ? — *As.* : N. W. Himalaya.
- californica* Lesq. in *Trans. Am. Phil. XIII*, p. 8 ( ): *Sull. Ic. M. Suppl.* p. 41, t. 27 (1874); *Lesq. et Jam. Man.* p. 153 (1884).  
 l. Rupic. — *Am. sept.* : Columb. brit., ins. Vancouver, Idaho, M. Calif., reg. occid.
- « *var. pilifera* Lesq. et Jam. *op. et l. cc.*  
*Hedwigia pilifera* Mitt. in *Journ. of the Linn. Soc. VIII*, p. 45, t. 7 (1864).  
 Ut typ. — *Am. sept.* : Ins. Vancouver, Calif.
- canescens* W. P. Sch. in *Mandón Pl. Boliv. n.* 1641; C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 139.  
 Br. *incana* C. M. in *Linn.* 1878-79, p. 377.  
 l. Cortic. — *Am. merid.* : And. Boliv., Argentin. subtropic.
- cirrifolia* (Wils.) Jaeg. *Ad. II*, p. 87 (1874-75).  
*Hedwigia cirrifolia* Wils. in *Mitt. M. austr. am.* p. 406 (1869).  
 — *Am. merid.* : And. Quitens.
- cochlearifolia* C. M. in *Hedwig.* 1897, p. 105.  
 Ster. — *Am. merid.* : La Plata.
- Delavayi* Besch. in *Ann. sc. nat.* 1892, I, p. 71.  
 l ? Cortic. — *As.* : Yunnan.
- diaphana* (C. M.) Jaeg. *op. et l. cc.*  
*Neckera diaphana* C. M. *Syn. II*, p. 105 (1851).  
 Ster. — *Afr.* : Pr. B. Sp.
- Elliottii* Broth. in *Engler's Bot. Jahrb.* 1897, p. 253.  
 Ster. — *Afr.* : Zambèze.
- entodonticarpa* C. M. in *Fl.* 1880, p. 485.  
 — *Afr.* : M. Kenia, Comor. magna.
- exserta* C. M. in *Linn.* 1878-79, p. 379.  
 l. Cortic. et rupic. — *Am. merid.* : Argentin. subtropic.
- « *var. reflexifolia* C. M. in *op. cit.* p. 380.  
 Ubi et c. typ.
- Humboldtii* Hook. — *Rhacocarpus.*  
*incana* C. M. — Br. *canescens.*
- indica* (Mont.) Par. *Ind.* p. 149 (1894).  
*Anoetangium incanum* Mont. in *sched.*  
*A. secundum* ej. in *Ann. sc. nat.* 1842, n. 32.  
 Br. *macropelma* Jaeg. *Ad. II*, p. 87 (1869-70).  
*Hedwigia indica* Mitt. *M. Ind. or.* p. 123 (1859).  
*Neckera macropelma* C. M. *Syn. II*, p. 104 (1851).  
 l. Ad rup. basalt. — *As.* : M. Neilgherr.

*Mac Owaniana* C. M. et Rehm. in Rehm. M. Afr. austr. N. 597 et in Hedwig. 1899, p. 123.

Ster. In fiss. rup. basalt. — Afr. : Pr. B. Sp., Natai.

*macropelma* Jaeg. — Br. indica.

*Novae Seelandiae* C. M. in Trans. and Proced. of the N. Zeal. Inst. 1893, p. 275.

— Pac. : Nov. Seeland.

*plicata* (Mitt.) Jaeg. op. et l. cc.

*Hedwigia plicata* Mitt. M. austr. am. p. 406 (1869).

— Am. merid. : And. Quitens., Boliv.

*rupestris* (Mitt.) Par. Ind. p. 150 (1893).

*Hedwigia rupestris* Mitt. in Journ. of the Linn. Soc. 1863. p. 160.

1.

— Afr. : Abyss., Cameroon.

*Schimperiana* Bryol. eur. l. c.

*Harrisonia Schimperiana* Hpe. in exc. ad. H. squarrulosam in Ic. M. t. 19 (1844).

*Hedwigia Schimperiana* Br. et W. P. Sch. in W. Sch. M. Abyss. n. 464.

*Neckera Schimperiana* C. M. Syn. II, p. 104 (1851).

1.

— Afr. : Abyss., Choa, Kilima N'Djaro.

*sciuroides* Bryol. eur. — Br. alopecura.

*secunda* (Hook.) Bryol. eur. l. c.

*Anoetangium secundum* Brid. Bryol. univ. II, p. 165 (1827).

*Harrisonia secunda* Spreng. Syst. veg. IV, p. 145 (1827).

*Hedwigia secunda* Hook. M. ex. t. 46 (1818-20).

*Neckera secunda* C. M. Syn. II, p. 103.

1. Saxic. — Afr. : Kilima N'Djaro (?). Am. sept. : Mexic.

*subincana* Broth. in Bihang till. K. Sv. Vet.-Akad. Handl. 1900, p.

1. Cortic. — Am. merid. : Rio Grande do Sul.

*subplicata* D<sup>na</sup> E. G. Britton in Bull. Torr. bot. Club. 1896, p. 98.

— Am. merid. : And. Boliv.

**BREUTELIA** W. P. Sch. Coroll. p. 85 (1856). — 78 spec.

*abyssinica* W. P. Sch. in W. Sch. M. Abyss. n. 424.

2.

— Afr. : Abyss.

*aciphylla* (Wils.) Jaeg. Ad. II, p. 560 (1873-74).

*Bartramia aciphylla* Wils. mss. ; Mitt. in Kew Journ. bot. III, p. 356 et M. austr. am. p. 266 (1869).

Ster.

— Am. merid. : And. Quitens.

*affinis* (Hook.) Mitt. in Kew Journ. bot. 1856, p. 261.

*Bartramia affinis* Hook. M. ex. t. 176 (1820) ; C. M. Syn. I, p. 505 (1849) ; Fl. Tasman. II, p. 194 (1867).

B. Brownei Brid. Bryol. univ. II, p. 49 (1827).

*Glyphocarpus affinis* Jaeg. Ad. I, p. 525 (1873-74).

*Philonotis affinis* ej. op. cit. II, p. 768 (1877-78) ; Par. Ind. p. 916 (1896).

2.

— Pac. : Austral. or. (Nov. Vales), Nov. Seeland.,

Tasman.

afro-scoparia (C. M.) Par. Ind. p. 150 (1894).

Bartramia afro-scoparia C. M. in *M. capens. Mac Owanianis et in Hedwig.* 1899, p. 91.

B. laete virens Rehm. *M. Afr. austr.* n. 206.

— Afr. : Pr. B. Sp.

angustifolia Rehm. in *Coll. cit.* n. 538.

— Afr. : Transvaal.

arcuata (Dicks.) W. P. Sch. op. et l. cc.

Bartramia arcuata Sw. in *Schrad. Journ. d. bot.* II, p. 182 (1801); *Schwaeagr. Suppl. I, P. II, p. 61, t. 62* (1816) et *Suppl. III, P. I, p. 240* (1827-28); *Brid. Bryol. univ.* II, p. 35 (1827); *Bryol. eur.* IV, Mon. p. 15, t. 321 (1842); *C. M. Syn. I, p. 487* (1849).

Br. chrysocoma Lindb. in *Oefv. Vet.-Akad. Foerh.* 1863, n. 7, Sep. Abdr. p. 7.

Hypnum chrysocomum Dicks. *Pl. crypt. brit. fasc. II, p. 12* (1790).

H. palustre erectum, coma lutea basi nigricante Dill. *Hist. M. p. 302, t. 39, f. 36* (1741) et *Herbar.*

Mnium arcuatum ej. *fasc. III, p. 2, t. 7, f. 3* (1793).

Mnium chrysocomum Hedw. *Sp. M. p. 74* (1801).

2. Ad rup. humid. — Eur. : Norveg., Britann., Westphal., Helvet. ; Corsica (ster.)

♂ var. major Besch. *Prodr. bryol. mexic.* p. 60 (1871).

Ut typ. — Am. sept. : Mexic.

aureola Besch. in *Bull. Soc. bot. Fr.* 1885, p. LXVI et in *Miss. sc. Cap Horn, V, Bot. p. 290, t. 4, f. 16* (1889).

Bartramia integrifolia Sull. in *Wilkes Exped.* p. 12.

Fr. ign.

— Am. merid. : Patag. austr., terr. Magellan.

austro-arcuata (C. M.) Par. Ind. p. 151 (1894).

Bartramia austro-arcuata C. M. in *Linn.* 1880-82, p. 406.

Fl. et fr. ign. Rupic. — Am. merid. : Argentin. subtropic.

azorica (Mitt.) Card. in *Eighth Ann. Rep. Missouri bot. Gard.* p. 65 (1897).

Bartramia azorica Mitt. in *Godman's Hist. nat. Azor.* (1870).

Ster.

— Afr. : Ins. Azor.

brachycoma Besch. in *Bull. Soc. bot. Fr.* 1885, p. LXVI et in *Miss. sc. Cap Horn, V, Bot. p. 290, t. 4, p. 15* (1889).

Br. comosa ej. in *Fl.* 1885.

Fr. ign.

— Am. merid. : Chile, Patag.

Brittoniae Ren. et Card. in *Bull. Soc. roy. bot. Belg.* II, p. 161.

2. Ster. — Am. sept. : Costa Rica. Am. merid. : And. Boliv.

carinata (Mitt.) Jaeg. *Ad. I, p. 557* (1873-74).

Bartramia carinata Mitt. *M. austr. am. p. 266* (1869).

Ster.

— Am. merid. : And. Chilens.

chilensis (Ltz.) Par.

Bartramia chilensis Ltz. in *Bot. Zeit.* 1866, p. 186 ; *Par. Ind.* 1893, p. 104.

2 ? Arenic. — Am. merid. : Chile.

chrysea (C. M.) Jaeg. op. cit. p. 560.

Bartramia chrysea C. M. in *Linn.* 1874, p. 598.

Ster.

— Am. merid. : And. Nov. Granat.

*chrysocoma* Lindb. — Br. *arcuata*.

*commutata* (Hpe.) Jaeg. *Ad. II*, p. 702 (1877-78); *Weymouth in Pap. and Proced. of the roy. Soc. of Tasman.* 1893, p. 210 (*descript. locupletiss.*)

*Bartramia affinis Schwaegr. Suppl. III. P. I, II*, t. 237 (1829-30) *fid. Hpe!*  
*B. commutata Hpe. in Linn.* 1876, p. 307.

2. — Pac. : Austral. (Victoria), Tasman.

*comosa* Mitt. in *Journ. of the Linn. Soc.* 1859, p. 82; *Fl. Tasman. II*, p. 195, t. 174, f. 7 (1860).

*Bartramia comosa Handb. of the N. Zeal. Fl.* p. 449 (1867).

2. *Rupic.* — Pac. : Austral. or. (Nov. Vales.), N. Seeland. sept. et med., Tasman.

*comosa* Besch. — Br. *brachycoma*.

*consimilis* (Hook. f. et Wils.) Jaeg. *op. cit.* p. 559.

*Bartramia consimilis Handb. of the N. Zeal. Fl.* p. 49 (1867).

*Hypnum consimile H. f. et W. Fl. antarct. I*, p. 137, t. 60, f. 4 (1844);  
*C. M. Syn. I*, p. 492 (1849) et *II*, p. 617 (1851).

*Ster.* — Pac. : Ins. Auckland.

*crassa* (Hook. f. et Wils.) Jaeg. *op. cit.* p. 556.

*Bartramia crassa H. f. et W. Fl. Tasman. II*, p. 194, t. 174, f. 5 (1860).

2. *Ad rup. madid.* — Pac. : Tasman.

*crassicaulis* (C. M.) Par. *Suppl. Ind.* p. 53 (1900).

*Bartramia crassicaulis C. M. in Fl.* 1896, p. 447.

*Ster.* — Pac. : Ins. Hawai.

*cuspidatissima* (C. M.) Par. *Ind.* p. 152 (1894).

*Bartramia cuspidatissima C. M. in Linn.* 1880-83, p. 408.

*Fl. et fr. ign. Rupic.* — Am. merid. : Argentin. subtropic.

*declivium* (C. M.) Par. *Suppl. Ind.* p. 53 (1900).

*Bartramia declivium C. M. in Bull. hb. Boissier*, 1898, p. 46 et in *Ule Coll. M. brasil. n.* 1817, 1818.

*Ster.* — Am. merid. : Brasil. or. (S. do Itatiaia).

*deflexa* Wils. — Br. *indica*.

*dicranacea* (C. M.) Mitt. *M. Ind. or.* p. 64 (1859).

*Bartramia dicranacea C. M. in Bot. Zeit.* 1853, p. 57.

2. — As. : M. Neilgherr.

*diffracta* Mitt. in *Journ. of the Linn. Soc.* 1863, p. 153.

2. — Afr. : Cameroon.

*divaricata* Mitt. in *Journ. of the Linn. Soc.* 1895, p. 82.

*Bartramia divaricata Fl. Tasman. II*, p. 105 (1860); *Handb. of the N. Zeal. Fl.* p. 449 (1867).

*B. gigantea H. f. et W. Fl. N. Zeal. II*, p. 90 (1855).

2. — Pac. : Nov. Seeland., Tasman.

*dumosa* Mitt. *op. cit.* p. 82.

*Bartramia dumosa ej. M. austr. am.* p. 267 p.p. (1869). *fid. C. M.*

*B. pendula Sull. in Wilkes Expl. Exped.* p. 122 (1859).

2. — Afr. : Ins. Kerguelen, Marion. Am. merid.  
 Ins. Chiloe, Chile, Fuegia, ins. Eremitae.

*Eggersiana* (C. M.) Broth. hb.

Dicranum Eggersianum C. M. in *Hedwig*. 1898, p. 228 ; *Par. Suppl. Ind.* p. 122 (1900).

Ster. Rupic. — *Am. sept.* : Ins. Dominic.

*elegans* Wright. — Br. grandis.

*elongata* Mitt. — Prionodon.

*Eugeniae Aongstr.* in *Oefv. af K. Vet.-Akad. Handl.* 1872, n. 5, p. 120.

Br. intermedia ej. prius.

2. — *Pac.* : Taïti.

*fusco aurea* Broth. in *Oefv. af Finska V. S. Foerh.* 1899, p. 103.

2. — *Pac.* : Austral. or. (Nov. Vales.).

*gigantea* (Brid.). *Bryol. jav.* I, p. 160, t. 130 (1855-61).

Bartramia gigantea Schwaegr. *Suppl. II, P. II*, p. 30, t. 161 (1826) ; *Brid.*

*Bryol. univ. II*, p. 36 (1827) ; *C. M. Syn. I*, p. 489.

Dicranum giganteum Brid. *Sp. M. I*, p. 97 (1806).

Hypnum arundinifolium Duby in *Moritz Verz. d. Zolling. Pfl.* p. 131 ( )

2. Rupic. et terr. — *As.* : Celebes, Java. *Afr.* : Ins. Borbon.

*glabrifolia* Dusen M. *Am. austr.* n. 413.

— *Am. merid.* : Patagon. occid.

*guaphalea* (P. B.) W. P. Sch. hb.

Bartramia arcuata Brid. *Bryol. univ. II*, p. 35 p.p (1827).

B. gnaphalea C. M. *Syn. I*, p. 489 (1849).

B. tomentosa Schwaegr. *Sp. M.* p. 97.

Hypnum? gnaphaleum P. B. *Prodr.* p. 64 (1805).

2. Fr. ign. — *Afr.* : Cameroon, ins. Fernando Po, Borbon.

*grandis* (Geh. et Hpe.) *Par. Ind.* p. 152 (1894).

Bartramia grandis elegans Wright in *Lond. Journ. of bot.* 1892, p. 263.

B. grandis Geh. et Hpe. in *Fl.* 1881, p. 369.

Prionodon robustus Hpe. *Symb.* 1872, p. 55 et *Enum. M. Brasil.* 1879.

Ster. — *Am. merid.* : Brasil. austr. or.

*Hariotiana* Besch. in *Bull. Soc. bot. Fr.* 1885, p. LXVII et in *Miss. sc.*

*Cap Horn, V, Bot.* p. 291, t. 4, f. 17.

Fr. ign. — *Am. merid.* : Patag., terr. Magell., Fuegia, ins. Eremitae.

*Hasskarliana* (Hpe.) *Jaeg. Ad. I*, p. 561 (1873-74).

Bartramia Hasskarliana Hpe. in *Fl.* 1865, n. 37.

— *Am. merid.* : Peruv. or.

*hispida* (Mitt.) *Jaeg. op. cit.* p. 560.

Bartramia hispida Mitt. *M. austr. am.* p. 264 (1869).

— *Am. sept.* : Jamaic.

*incana* (Tayl.) *Jaeg. op. cit.* p. 555.

Bartramia incana Tayl. in *Lond. Journ. of bot.* 1848, p. 189 ; *C. M.*

*Syn. I*, p. 501 (1849) ; Mitt. *M. austr. am.* p. 263 (1869).

2? Saxic. — *Am. merid.* : And. Quitens.

*inclinata* (Hpe. et Ltz.) *Jaeg. op. et l. cc.*

Bartramia inclinata Hpe. et Ltz. in *Bot. Zeit.* 1869, p.

2. Terr. — *Am. merid.* : And. Æquator.

*indica* Mitt. *M. Ind. or.* p. 64 (1859).

*Bartramia deflexa* (forma minor fol. magis deflexis) Wils. *in sched.*

— *As.* : Himalaya, M. Neilgherr.

*integrifolia* (Tayl.) Jaeg. *op. cit.* p. 558.

*Bartramia integrifolia* Tayl. *in Lond. Journ. of bot.* 1846, p. 55; *C. M. Syn. I*, p. 490; *Mitt. M. austr. am.* p. 267.

2. *Ad rup. irror.* — *Am. merid.* : M. Pichincha, And. Boliv.

*intermedia* (Hpe) Besch. *Prodr. bryol. mexic.* p. 60 (1871).

*Bartramia intermedia* Hpe. *Sp. nov. mexic. in Verh. d. K. K. zool. bot. Gesells. in Wien*, 1869, p. 507.

2. — *Am. sept.* : Mexic.

*intermedia* Aongstr. — Br. *Eugeniae*.

*jamaicensis* (Mitt.) Jaeg. *op. et l. cc.*

*Bartramia jamaicensis* Mitt. *op. cit.* p. 265.

— *Am. sept.* : Jamaic.

*Karsteniana* (C. M.) Jaeg. *op. cit.* p. 560.

*Bartramia Karsteniana* C. M. *in Linn.* 1874, p. 599.

2. — *Am. merid.* : Nov. Granat. ? (ster.), Equator.

*kilima N'Djarica* (C. M.) Par. *Ind.* p. 153 (1894).

*Bartramia kilima N'Djarica* C. M. *in Fl.* 1888, p. III.

Ster. *In graminos.* — *Afr.* : Kilima N'Djaro.

*longipes* W. P. Sch. *in Lechler Pl. Chil.* n. 813.

— *Am. merid.* : Chile.

*Lorentzi* (C. M.) Par. *op. et l. cc.*

*Bartramia Lorentzi* C. M. *in Linn.* 1880-82, p. 406.

*Fl. et fr. ign.* — *Am. merid.* : Argentin. subtropic.

*Mac-Leana* Rehm. *M. Afr. austr.* n. 539.

Ster. — *Afr.* : Transvaal.

*macrocarpa* W. P. Sch. *in Mandon Pl. Boliv.*

*Bartramia macrocarpa* C. M. *in Nuov. Giorn. bot. ital.* 1897, p. 45.

*B. mniocarpa* ej. *in hb. New-York*, n. 3137.

2. — *Am. merid.* : And. Boliv.

*macrotheca* Jaeg. — Br. *tomentosa*.

*mniocarpa* (W. P. Sch.) Par. *Suppl. Ind.* p. 53 (1900).

*Bartramia mniocarpa* C. M. *in op. cit.* p. 44.

*Philonotis mniocarpa* W. P. Sch. *in Mandon Pl. Boliv.* n. 1677.

— *Am. merid.* : And. Boliv.

*Mülleri* Jaeg. *Ad. I*, p. 561 (1873-74).

*Bartramia divaricata* C. M. *in Linn.* 1874, p. 591.

Ster. — *Am. sept.* : Costa Rica.

*pendula* (Hook.) Jaeg. *op. cit.* p. 558.

*Bartramia pendula* Hook. *M. ex. t.* 21 (1818-20); *C. M. Syn. I*, p. 491; *Fl. Tasm. II*, p. 191 p.p. (1860); *Handb. of the N. Zeal. Fl.* p. 418 (1867).

*Bryum pseudulum* Brid. *Mant. M.* p. 120 (1819).

- Mnium pendulum* Sm. in *Trans. of the Linn. Soc. VII*, p. 262.  
*Philonotis pendula* Brid. *Bryol. univ. II*, p. 27 (1827).
2. — *Afr.* : Ins. Kerguelen? *Am. merid.* : Fuegia, ins. Eremitae. *Pac.* : Austral. or. (Nov. Vales.), Nov. Seeland., Tasman., ins. Campbell.
- persquarrosa* Dusen M. *Am. austr. n.* 376 et 477.  
 Palud. et terr. — *Am. merid.* : Chile austr., Patagon. occid.
- perruncinata* ej. *Coll. cit. n.* 697.  
 — *Am. merid.* : Chile austr.
- pinnata* (Hpe.) Jaeg. *op. cit. p.* 559.  
*Bartramia pinnata* Hpe. in *Ann. sc. nat. V ser. III*, p. 373 (1866); Mitt. *M. austr. am. p.* 265 (1869).  
 Ster. — *Am. merid.* : And. Novo-Granat.
- plicata* Mitt. in *Journ. of the Linn. Soc.* 1859, p. 83.  
*Bartramia plicata* ej. *M. austr. am. p.* 266.  
 Paludic. — *Am. merid.* : And. Chilens.
- Readeriana* (Col.) A. Gepp in litt.  
*Bartramia Readeriana* Col. in *Trans. and Proceed. of the N. Zeal. Inst.* 1884, p. 258.  
 — *Pac.* : Nov. Seeland.
- rivalis* (C. M.) Par. *Suppl. Ind. p.* 53 (1900).  
*Bartramia rivalis* C. M. in *Bull. hb. Boissier*, 1898, p. 46 et in *Ule Coll. M. brasil. n.* 1820.  
 Ster. Ad. rivul. — *Am. merid.* : Brasil. or. (S. do Itatiaia).
- robusta* (Hook. f. et Wils.) Jaeg. *op. cit. p.* 560.  
*Bartramia robusta* H. f. et W. *Fl. antarct. I*, p. 133, t. 59 (1844) et *Handb. of the N. Zeal. Fl. p.* 447 (1867); C. M. *Syn. I*, p. 495 (1849) et *II*, p. 619 (1851).  
 Ster. — *Am. merid.* : Fuegia. *Pac.* : Ins. Auckland. Campbell.
- rupestris* (Mitt.) Jaeg. *op. et l. cc.*  
*Bartramia rupestris* Mitt. in *Handb. of the N. Zeal. Fl. p.* 447 (1867).  
 ♀ et fr. ign. Ad rup. marit. — *Am. merid.* : Fuegia.
- scabrifolia* Dusen M. *Am. austr. n.* 413.  
 — *Am. merid.* : Patagon. occid.
- scoparia* (Schwaegr.) W. P. Sch. hb.  
*Bartramia scoparia* Schwaegr. *Suppl. III, P. I, II, t.* 211 (1829-30); C. M. *Syn. I*, p. 490 (1849).  
 2. — *Am. sept.* : Antill. gall.
- secundifolia* (C. M.) Par. *Suppl. Ind. p.* 54 (1900).  
*Bartramia secundifolia* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 46.  
 2. — *Am. merid.* : And. Boliv.
- Sieberi* (Hsch.) Mitt. in *Journ. of the Linn. Soc.* 1859, p. 83.  
*Bartramia pendula* Spreng. *Syst. veg. IV, P. II, p.* 323 (1827).  
 B. *scoparia* Schwaegr. *Sp. M. p.* 98.  
 B. *Sieberi* Hsch. in *Sieber M. Nov. Holl. n.* 13; C. M. *Syn. I*, p. 491; *Fl. Tasman. II*, p. 194, t. 174, f. 6 (1860); *Handb. of the N. Zeal. Fl. p.* 449 (1867).  
 2. — *Pac.* : Austral., Nov. Seeland. sept. et merid., Tasman.

- Spielhausi (C. M.) *Par. Suppl. Ind.* p. 54 (1900).  
*Bartramia Spielhausi* C. M. in *Hedwig*. 1899, p. 91 ; *Par. Ind.* p. 112 (1894).  
 Ster. — *Afr.* : Pr. B. Sp.
- squarrosa (Mitt.) *Jaeg. op. et l. cc.*  
*Bartramia squarrosa* Mitt. *M. austr. am.* p. 265 1869.  
 Ster. — *Am. merid.* : And. Nov. Granat.
- Stuhlmanni *Broth. in Engler's Bot. Jahrb.* 1894, p. 192.  
 2. ♂ ign. — *Afr.* : Reg. lacustr.
- subarcuata W. P. Sch. *ms.* ; *Besch. Prodr. bryol. mexic.* p. 60 (1871).  
*Bartramia subarcuata* C. M. *Syn. II*, p. 617 ; Mitt. *op. cit.* p. 262.  
 2. — *Am. sept.* : Mexic. *Am. merid.* : And. Nov. Granat.
- subdisticha (Hpe.) *Jaeg. op. cit.* p. 556.  
*Bartramia subdisticha* Hpe. *Symb.* 1872, p. 50.  
 ♂ et fr. ign. — *Am. merid.* : Brasil. austr. or.
- subelongata *Broth. in Dusen M. Am. austr. n.* 444.  
 Ad rup. irror. — *Am. merid.* : Patagon. occid.
- subgnaphalea (C. M.) *Par. Ind.* p. 154 (1894).  
*Bartramia subgnaphalea* C. M. in *Fl.* 1890, p. 480.  
 2. — *Afr.* : Kilima N'Djaro, reg. lacustr.
- submniocarpa (C. M.) *Par. Suppl. Ind.* p. 54 (1900).  
*Bartramia submniocarpa* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 28.  
 2. — *Am. merid.* : And. Boliv.
- subtomentosa (Hpe.) *Jaeg. op. et l. cc.*  
*Bartramia subtomentosa* Hpe. *Symb.* 1872, p. 49 et 1877, p. 723.  
 Ster. — *Am. merid.* : Brasil. austr. or.
- tenuifolia (Mitt.) *Par. Suppl. Ind.* p. 54 (1900).  
*Bartramia tenuifolia* Mitt. in *Meliiss's S. Helena*, 1875.  
 Ster. Terr. — *Afr.* : Ins. Tristao da Cunha.
- thysanomitrioides *Dusen M. Am. austr. n.* 623.  
 — *Am. merid.* : Chile austr.
- tomentosa (Sw.) W. P. Sch. *hb.*  
*Bartramia arcuata* *Schwaegr. Suppl. I, P. II*, p. 62 (1816).  
*B. macrocarpa* Hpe. in *Linn.* 1863, p. 141.  
*B. macrotheca* ej. in *Ann. sc. nat. V ser. III*, p. 373 (1866) *fid. Mitt.*  
*B. tomentosa* C. M. *Syn. I*, p. 488 (1849).  
*Br. macrotheca* *Jaeg. Ad. I*, p. 556 (1873-74).  
*Bryum tomentosum* Sw. *Fl. Ind. occ. III*, p. 1837 (1806).  
*Mnium tomentosum* ej. *Prodr.* p. 139 (1795).  
 ? *Philonotis tomentosa* Brid. *Bryol. univ. II*, p. 26 (1826).  
 2. — *Am. sept.* : Costa Rica, Guadal. Jamaic. *Am. merid.* : Venezuela, And. Nov. Granat., Quitens. et Boliv.
- Trianae (Hpe.) *Jaeg. op. cit.* p. 558.  
*Bartramia Trianae* Hpe. in *Ann. sc. nat. V ser. III*, p. 374 (1866).  
 Ster. — *Am. merid.* : And. Nov. Granat.



*Ulei* C. M. in *Ule Bryoth. brasil.* n. 36

*Bartramia Ulei* *ej. in Bull. hb. Boissier*, 1898, p. 16 et in *Ule Coll. M. brasil.* N. 662 et 1819.

— *Am. merid.* : Brasil. subtropic. or.

*Wainioi Broth. in Bot. Centralbl.* 1888, p. 86.

2. — *Am. merid.* : Brasil. or.

*yunnanensis Besch. in Ann. sc. nat.* 1892, I, p. 63.

Ster. — *As.* : Yunnan.

**BROTHERA** C. M. in *Kindb. Enum.* 1891, p. 105. — 1 spec.

*Leana (Sull.) C. M. mss.*

*Campylopus Leanus Sull. M. of U. St.* p. 19 (1856) et  *Ic. M.* p. 29, t. 18 (1864); *Lesq. et Jam. Man.* p. 78 (1884).

*Dicranum albicans Bryol. eur. (fid. Hpe. in Jaeg. Ad. II,* p. 656).

*Leucobryum Leanum Kindb. Eur. and N. Am. Bryin.* p. 176 (1891).

*Leucophanes? Leanus Sull. M. Allegh. I ed. n.* 172 (1816); *C. M. Syn. I.* p. 446 (1849).

*Syrrophodon Leanus Sull. l. c. (fid. Lesq. et Jam. l. c.).*

2. *Lignic.* — *As.* : Japon, Simla. *Am. sept.* : Minnesota, Ohio, reg. centr.

**BRUCHIA** Schwaegr. *Suppl. II*, p. 91 (1823) (1). — 25 spec.

*acuminata Broth. in Bihang till K. Sv. Vet.-Akad. Handl.* 1900, p.

1. *Ad terr. uligin.* — *Am. merid.* : Paraguay.

*amoena C. M. in Fl.* 1888, n. 1.

1 ? — *Pac.* : Australia.

*aurea Besch. Not. M. Paraguay*, p. 259 (1877).

*Br. ligulata C. M. in Fl.* 1888, n. 1.

1. *Ad terr. apric.* — *Am. merid.* : Paraguay.

*Beyrichiana (Hpe) C. M. Syn. I*, p. 18 (1849).

*Phascum Beyrichianum Schwaegr. Suppl. IV*, p. 301 (1842).

*Sporledera Beyrichiana Hpe. in Linn.* 1837, p. 279; *Par. Ind.* p. 1229 (1897).

1. — *Am. sept.* : New Jersey, Maryland, Illinois.

*Beyrichiana Sull., Lesq. et Jam.* — *Br. flexuosa (fid. D<sup>na</sup> E. G. Britton).*

*Bolanderi Lesq. in Mem. Calif. Acad. I*, p. 5 ( ); *Sull. Ic. M. Suppl.* p. 23 (1874); *Lesq. et Jam. Man.* p. 46 (1884).

— *Am. sept.* : M. Calif.

*brevicollis Lesq. et Jam. Man.* p. 47 (1884).

*Br. flexuosa (fid. D<sup>na</sup> E. G. Britton).*

*Syn.* — *Am. sept.* : Carol. infer.

*brevifolia Sull. M. of U. St.* p. 17 (1856) et  *Ic. M.* p. 25, t. 15 (1864); *Lesq. op. et l. cc.*

*Br. vogesiaca n. 2 H. f. et W. in Drumm. M. bor. am. II ed. n.* 15 p.p.

1. *Ad terr. arenos.* — *Am. sept.* : Carol. infer., Ludovic., Texas.

---

(1) De *Bruchiis* N. Amer. cf. D<sup>nam</sup> E. G. Britton in *Bull. Torr. bot. Club*, 1894, pp. 343-372 et tt. 213-217.

*brevipes* Hook.  *Ic. pl. rar. III*, p. 231, p.p. (1844); *Sull. Ic. M.* p. 24, t. 14, p.p. (1864); *Lesq. et Jam. op. et l. cc.*

Br. Drummondii Hpe. in sched.; *Jaeg. M. Cleistoc.* p. 36; *D<sup>na</sup> E. G. Britton in op. cit.* p. 261.

Br. flexuosa var. minor C. M. *Syn. I*, p. 20 (1849).

1. Ad terr. argill. arenac. — *Am. sept.*: Carol., Virgin., Florid., Ludovic.

*brevipes* Hook. p.p. — Br. elegans.

Caroliniae Aust. in *Bull. Torr. bot. Club*, VI, p. 144 (1879).

Br. Ravenelii var. mollis Lesq. et *Jam. Man.* p. 49 (1884).

1. Terr. — *Am. merid.*: Florid.

*curviseta* Lesq. et Jam. — Br. texana.

Donnellii Aust. in *Bull. Torr. bot. Club*, VI, p. 144 (1879); *Lesq. et Jam. Man.* p. 48 (1884).

Br. flexuosa Sull. et Lesq. *M. bor. am. II ed. n.* 41 p.p. (fid. *D<sup>na</sup> E. G. Britton.*).

1. — *Am. sept.*: Florid.

*Drummondii* Hpe. — Br. brevipes.

Eckloniana C. M. *Syn. I*, p. 19 (1849).

*Spoledera* Hpe mss.; *Par. Ind.* p. 1229 (1897).

2. — *Afr.*: Prom. B. Sp.

elegans (Hsch.) *Jaeg. M. Cleistoc.* p. 35 ( )

Br. brevipes Hook. op. et l. cc. p.p.; C. M. *Syn. I*, p. 18.

*Phascum elegans* Hsch. in *Linn.* 1841, p. 114.

1. — *Afr.*: Pr. B. Sp.

*exigua* C. M. — *Pleuridium Krauseanum*.

flexuosa (Schwaegr.) C. M. in *Bot. Zeit.* 1847 et *Syn. I*, p. 20; *Lesq. et Jam. Man.* p. 46.

*Phascum flexuosum* Schwaegr. *Suppl. II, P. I*, p. 101, excl. fig. 3 et 8 (1823); *Brid. Bryol. univ. I*, p. 38 (1827).

2 (C. M.), 1 vel syn. (Lesq. et Jam.). — *Am. sept.*: Minnesota, Reg. merid.

“ var. microcarpa (Wils.) *Lesq. et Jam. op. et l. cc.*

Br. microcarpa Wils. mss.

Br. vogesiaca *Drumm. M. bor. am. II ed. n.* 14.

*Sporledera setifolia* *Jaeg. M. cleistoc.* p. 35 ( ) et *Ad. II*, p. 376 (1876-77)

Ut et cum typ.; Ludovic.

flexuosa Sull. — Br. Sullivantii.

var. minor C. M. — Br. brevipes.

var. nigricans Sull. — Br. Sullivantii var. nigricans.

fusca *D<sup>na</sup> E. G. Britton in op. cit.* p. 361, t. 216.

1. Inter lapid. quartz. in arenac. — *Am. sept.*: Maryland, Carol. super.

Hallii Aust. in *Bull. Torr. bot. Club*, V, p. 21 (1878); *Lesq. et Jam. op. cit.* p. 47.

— *Am. sept.*: Texas.

Hampeana C. M. *Syn. I*, p. 18 (1849).

Br. Schwaegrichenii *Jaeg. M. cleistoc.* p. 36 ( ) et *Ad. I*, p. 376 (1871-72).

*Phascum brevipes* Schwaegr. *Suppl. IV*, p. 303 b.

- Sporledera Hampeana* Mitt. *M. austr. am.* p. 25 (1869); *Par. Ind.* p. 122 (1897).  
 1. — *Am. sept.*: Ludovic. *Am. merid.*: Chile.
- ligulata* C. M. — Br. aurea.
- longicollis* Eaton in *Bull. Torr. bot. Club.* 1890, p. 100, t. 101.  
 1. — *Am. sept.*: New Hampshire.
- microcarpa* Wils. — Br. flexuosa var. microcarpa.
- minuta* Mitt. in *Journ. of the Linn. Soc.* 1859, p. 65 et in *Fl. Tasman.* II, p. 165, t. 171, f. 4 (1860).  
*Sporledera minuta* ej. in *Handb. of the N. Zeal. Fl.* p. 223 (1867; *Par. Ind.* p. 1229 (1897).  
 1. Terr. — *Pac.*: Nov. Seeland., Tasman.
- nigricans* Aust. — Br. Sullivantii var. nigricans.
- palustris* C. M. — *Sporledera*.
- Ravenelii* Wils. mss.; *Sull. M. of U. St.* p. 17 (1856) et *Ik. M.* p. 26, t. 16 (1864); *Lesq. et Jam. op. cit.* p. 49.  
*Sporledera Ravenelii* Jaeg. *M. cleist.* p. 35.  
 1. Terr. — *Am. sept.*: Carolin. infer., Alabama.  
 var. *mollis* Lesq. et Jam. — Br. Caroliniae.
- Rehmanni* C. M. — *Sporledera*.
- repens* Hach. — *Platygyrium*.
- Schwaegrichenii* Jaeg. — Br. Hampeana.
- Sullivantii* Aust. in *Bull. Torr. bot. Club*, VI, p. 143 (1879); *Lesq. et Jam. Man.* p. 46 (1884).  
 Br. flexuosa *Sull. Ik. M.* p. 22, t. 13 excl. descript. (1864); *Sull. et Lesq. M. bor. am. n.* 41 p.p. (fid. D<sup>na</sup> E. G. Britton).  
 1. Ad terr. argill. — *Am. sept.*: Ohio, reg. merid.  
 « var. *nigricans* *Lesq. et Jam. op. et l. cc.*  
 Br. flexuosa var. *nigricans* *Sull. M. of U. St.* p. 17 (1856); *Sull. et Lesq. M. bor. am. I ed.* n. 42.  
 Br. *nigricans* Aust. in *op. et l. cc.*  
 Ut typ. — *Am. sept.*: Alabama.
- texana* Aust. in *Bull. Torr. bot. Club*, 1878, p. 21.  
 1. — *Am. sept.*: Texas.
- trobasiana* de Not. in *Erb. crittog. ital.* n. 719 (1862) et *Epil. briol. ital.* p. 729 (1869); *W. P. Sch. Bryol. eur. Suppl. I et II*, t. 1 et *Syn. II ed.* p. 29 (1876).  
 i. In pratis arid. — *Eur.*: Stiria, lac. Maj.
- Uleana* C. M. in *Ule Bryoth. brasil.* n. 3.  
 1. Terr. — *Am. merid.*: Brasil. subtropic. or.
- uruguensis* C. M. in *Linn.* 1878-79, p. 231.  
 2. — *Am. merid.*: Paraguay, Uruguay.
- vogesiac* *Schwaegr. Suppl. II, P. II*, p. 91, t. 127 (1824); *Bryol. eur.* I, t. II; *C. M. Syn. I*, p. 20; *W. P. Sch. Syn. ed. II*, p. 28.  
*Saproma vogesiaceum* Brid. *Bryol. univ.* I, p. 53 (1827).  
*Saproma cyrtophyllum* ej. *op. et l. cc.*

*Voitia vogesiaca* Hsch. in Moug. et Nestl. Stirp. crypt. Voges. Rhen. fasc. VIII, n. 706.

1. In decliv. gramin. humid. et in turfac. — Eur. : Voges. (M. Hohnneck), Palatin. super., Salisb.

*vogesiaca* { Drumm. — Br. flexuosa var. microcarpa.  
                   { H. f. et W. p.p. } — Br. brevifolia.  
                   { — Br. curviseta.

*Whiteleggei* C. M. — Sporledera.

**BRYELLA** Berk. Handb. of brit. M. 1863.

*recta* Berk. — Phascum.

**BRYHNIA** Kaur. in Bot. Notis. 1892, Heft. II. — 2 spec.

*graminicolor* Grout. — Eurhynchium.

*Novae Angliae* Grout. — Brachythecium.

*scabrida* (Lindb.) Kaur. in op. et l. cc.; Joergersen in Bergen's Mus. Adsb. f. 1894-95, p. 25 c. ic.

Brachythecium Novae Angliae (fid. Grout in Bull. Torr. bot. Club, 1898, p. 229).

Eurhynchium scabridum Kindb. Eur. and N. Am. Bryin. p. 94 (1897).

Hypnum scabridum Lindb. in Bot. Notis. 1887, p. 41.

2. Rupic. — Eur. : Norveg. merid. Am. sept. : Terra nov.

*Tokubuchii* (Broth.) Par.

Hypnum Tokubuchi Broth. in Hedwig. 1899, p. 241.

2. — As. : Japon.

**Bryoidium** C. M. in Nuov. Giorn. bot. ital. 1897, p. 6 (nom., sect. FISSIDENTIS).

**Bryotis** C. M. in Linn. 1875, p. 380 (sect. BRYI).

**Bryotypus** Hagen in Tromsøe Mus. Aarsheft. 1898-99, p. 120 (sect. BRYI).

**BRYOXIPHIMUM** (1) Mitt. M. austr. am. pp. 24 et 580. — 3 spec.

*mexicanum* Besch. in Journ. de bot. 1892, p. 180.

Br. norvegicum Mitt. M. austr. am. p. 580.

Eustichium norvegicum W. P. Sch. in Besch. Prodr. bryol. mexic. p. 29 (1871).

♀ et fr. ign. — Am. sept. : Mexic.

*norvegicum* (Brid.) Mitt. in Journ. of the Linn. Soc. XII, p. 580 ( ).

Eustichia imbricata Kindb. Eur. and N. Am. Bryin. p. 153 (1897).

Eustichia norvegica C. M. Syn. I. p. 42 (1819); W. P. Sch. Syn. II ed. p. 148 (1876).

Eustichium norvegicum Bryol. eur. III, p. 195.

Fissidens imbricatus Desv. in Brid. Bryol. univ. II, p. 671 (1827).

Phyllogonium Eustichia norvegicum Brid. op. et l. cc.

Ph. norvegicum Sull. Contrib. on Bryol. of N. Amer. occ. ( ).

2. Rupic. — Eur. : Norveg. ? Island. Am. sept. : de Ohio centr. ad Kentucky (ster.); Wisconsin (c. fr.) (2).

(1) Sub *Bryosiphys*, nom. a cl. Bescherelle in Journ. de bot. 1892, p. 180 correcto.

(2) Cf. O. Knight in Bull. Torr. bot. Club, 7 bri 1883 et in Rev. bryol. 1884, p. 15 (descript. fruct.).

*norvegicum* } Mitt. — *Br. mexicanum*.  
                   } Bryol. eur. — *Bryoxiphium*.

Savatieri (*Husn.*) Mitt. *M. and Hepat. from Japan*, 1891, p. 154.

*Eustichia japonica* Berggr. in *Fl.* 1881, p. 290 (*fid. Geh. in op. cit.* 1883, n. 31).

*E. norvegica* W. P. Sch. in *Savatier M. Japon.* n. 148.

*E. Savatieri* Husn. *Rev. bryol.* 1878, p. 85.

2. Ad rup. madid. et in spelunc. — *As.* : China centr. (prov. Schen-Si, Japon.

**BRYUM** Dill. Cat. Giss. p. 222 (1718) et Append. p. 85 (1719). — 831 spec. (1).  
 Spec. omn. non indicatae sunt RHODOBRYA ejusd. nom.

abbreviatum *Hpe. Enum. M. Brasil.* p. 31 (1879).

2 ? — *Am. merid.* : Brasil. austr. or.

abduanum *Rota Cat. M. Bergom.; de Not. Epil.* p. 734 (1869).

*Br. Mühlenbeckii* ? (*fid. Limpr.*).

Ster. Ad rup. et aren. humid. — *Eur.* : Ital. super. (Vall. Tellin.).

aberrans *Hpe. Symb.* 1877, p. 724.

Terr. — *Am. merid.* : Brasil. austr. or.

abruptinervium *C. M. in Hedwig.* 1898, p. 102.

*Rhodobryum abruptinervium* Par. *Suppl. Ind.* p. 298 (1900).

2. — *Pac.* : Austral. or. (Nov. Vales.)

« var. vel spec. propria ? *C. M. in op. et l. cc.*

— *Pac.* : Austral. or. (Victoria).

*abyssinicum* *C. M.* — *Brachymenium*.

*acanthoneuron* Aongstr. in *Oefv. af K. Vet.-Akad. Foerh.* 1876, n. 4, p. 20.

Ster. — *Am. merid.* : Brasil. austr. or.

*acaulon* { *Ericae tenuifoliae* *Geranii folio* Dill. — *Aloina ambigua*.  
                   *foliis teneris confertis, capsulis conicis* Dill. — *Tortella caespitosa* (*fid. Hedw.*).  
                   *heteromallum, setis longis pallidis* Dill. — *Ditrichum pallidum*.

*aciculare* L. — *Rhacomitrium*.

*aciculare montanum* Stroem. — *Rhacomitrium* ?

*aciculariforme* Hoffm. — *Rhacomitrium sudeticum* ?

*aciculinum* Kindb. *Eur. and N. Am. Bryin.* p. 315 (1897).

— *Am. sept.* : Columb. brit.

*aciphyllum* Voit. — *Mnium affine*.

var. ej. — *Mnium cuspidatum*.

*aculeatum* Joergens. in *Christiania Vid. Selsk.* 1894, p. 80.

Syn. — *Eur.* : Norveg.

*acuminatissimum* (*C. M.*) Broth. *Natür. Pflanzenf.* p. 612.

*Mnium acuminatissimum* *C. M. in Hedwig.* 1900, p. 248.

Ster. Terr. — *Am. merid.* : Rio de Janeiro.

*acuminatulum* *C. M.* — *Webera*.

(1) De *Bryis* Novo Seelandicis Brownianis (41) cl., p. 34 in *adn. et Broth. Natür. Pflanzenfam.* p. 598.

- acuminatum* { Bryol. eur. — Webera.  
 Rab. — Webera elongata var. acuminata.  
 Sw. — Barbula agraria.  
 Wils. — Webera himalayana.  
 var. arcuatum Bryol. eur. — Webera acuminata var. arcuata.  
 var. minus Bryol. eur. — Webera acuminata var. minor.  
 var. polysetum Bryol. eur. — Webera acuminata var. polyseta.  
 var. tenellum Bryol. eur. — Webera acuminata var. tenella.
- acutiforme* Limpr. in litt. ad Ryan, 1894; Hagen in Tromsø Mus. Aarsheft. p. 156 (1898-99).  
 1. Arenic. — Eur. : Spitzberg ? Norveg. bor. Am. sept. : Groenland.
- acutiusculum* C. M. in Fl. 1887, p. 220.  
 2. — Am. sept. : Alaska.
- acutum* Lindb. in Medd. Soc. pro Faun. et Fl. fenn. XIII Hefst. p. 239 (1886).  
 B. calophyllum Berggr. M. Spetsb. n. 73.  
 1 (Limpr., Philib.), polyg. (Lindb.). Ad rip. aren. fluv. — Eur. : Lapon. ross., Norveg. bor., Alp. Vales. As. : Jenissei infer. Am. sept. : Groenland.
- acutum* Huds. — Blindia.
- aequabile* C. M. in Linn. 1880-83, p. 368.  
 Ster. Muric. — Am. merid. : Argentin. subtropic.
- aeruginosum* C. M. in Hedwig. 1898, p. 95.  
 2. — Pac. : Austral. or. (N. Vales.).
- aerugo* Dusen M. Am. austr. n. 517.  
 Paludic. — Am. merid. : Patag. occid.
- aestivum* L. — Hymenostylium curvirostre.
- affine* { Brid., Wils. — Mnium.  
 Gmel. — Orthotrichum.  
 Lindb. — Br. cuspidatum.  
 var. cylindricum Arn. — Br. cuspidatum var. cylindricum.  
 var. urnigerum Arn. — Br. cuspidatum var. urnigerum.
- afro-alpinum* Rehm. M. Afr. austr. n. 248 b; C. M. in Hedwig. 1899, p. 73.  
 Ster. — Afr. : Orange.
- afro-crudum* C. M. — Webera.
- afro-dimorphum* Par.  
 Br. dimorphum C. M. in Rehm. M. Afr. austr. n. 243.  
 — Afr. : Austr.
- afro-julaceum* C. M. — Anomobryum ?
- afro-littorale* C. M. in Engler's Bot. Jahrb. 1883, p. 88 et in Forschungsreise... Bot. p. 62 (1889).  
 Ad terr. humid. — Afr. : Monrovia, Cameroon., Lagos.
- afro-nutans* C. M. — Webera.
- aggregatum* C. M. — Webera.
- agrarium* Sw. — Barbula.
- alandense* J. O. Bomanss. in Act. Soc. p. Faun. et Fl. fenn. 1899, n. 4.  
 1. In glareos. — Eur. : Ins. Aland.
- alandicum* Bomanss. — Br. Arnellii.

*alaskanum* Kindb. *Eur. a. N. Am. Bryin.* p. 381 (1897).  
— *Am. sept.* : Alaska.

*albicans* Mitt. } — *Mniobryum*.  
(1876). — *Mniobryum austro-albicans*.  
var. *glaciale* C. M. — *Mniobryum albicans* var. *glacialis*.

*albidum* { L. — *Octoblepharum*.  
et *glaucum, fragile majus* Dill. — *Leucobryum glaucum*.  
et *glaucum, fragile minus* Dill. — *Leucobryum minus*.

*albo-pulvinatum* C. M. in *Fl.* 1886, p. 507.  
Ster. — *Afr.* : Angola.

*albulum* Mitt. *M. Ind. or.* p. 69 (1859).  
2? — *As.* : India.

*Alexandri* Philib. in *Rev. bryol.* 1900, p. 57.  
— *As.* : M. Alexandre.

*algovicum* Sendtn. — *Br. pendulum* var. *compactum*.

*alpiniforme* Kindb. in *Mac. Cat.* p. 271 (1892).  
Ster. Rupic. — *Am. sept.* : Ontario.

*alpinulum* Besch. *Fl. bryol. Réunion, etc.* p. 97 (1880-81).  
2. Ster. Terr. — *Afr.* : Ins. Nossi-Be, Nossi-Comba, S. Maria de Mad.

*alpinum* Huds. *Fl. angl.* p. 415 (1762); *L. Mant. M. II*, p. 309 (1767) et *hb. fasc. V*; *Brid. Bryol. univ. I*, p. 663 (1826); *Bryol. eur. fasc. 6-9 (Vol. IV), Mon.* p. 76, t. 380 (1839); *C. M. Syn. I*, p. 285 (1849).

*Br. hypnoides pendulum sericeum, coma insigni atro-rubente* Dill. *Hist. M.* p. 394, t. 50, f. 61 (1741).

*Br. pachypoma* Wils. in *Journ. of bot.* 1840, *II*, p. 364 p.p.

*B. teretiusculum* Hook. in *Wall. Cat. n.* 7591; *Harc. in Journ. of bot. l. c.* p. 9 et in *Hook. Ic. pl. rar. t. 20, f. I* (1841) *fid. Mitt.*; *C. M. op. cit.* p. 303.

*Mnium alpinum* Sw. *Meth. M.* p. 356 (1787).

2. Ad rup. silic. madid. — *Eur.* : E reg. campestr. usq. in subalp. per zon. intermed. et sept., raro merid.; S. d'Estrella, de Guadarrama. *As.* : Khasia, Nepal., Himalaya, Tibet. occ., M. Neilgherr., Caucas. *Afr.* : Ins. Canar., Madeir., Cameroon, Kilima N'Djaro, ins. Kerguelen. *Am. sept.* : Columb. brit., reg. or.

« var. *angustifolium* Husn. *Muscol. gall.* p. 247 (1889).

*Br. atlanticum* Solms Laub.

Ut typ. — *Eur.* : Gall. occid. et merid., Astur.

var. *brevifolium* Myr. — *Br. Mühlenbeckii*.

« var. *denticulatum* Card. et Ther. in *Bot. Gaz.* 1900, p. 124.  
— *Am. sept.* : Montana.

« var. *distantifolium* Dixon Stud. *Handb.* p. 338 (1896).  
Rupic. ad torr. mont. — *Eur.* : Anglia.

var. *gemmiparum* Husn. { — *Br. gemmiparum*.  
var. *mediterraneum* de Not. }

« var. *meridionale* W. P. Sch. *op. cit.* p. 441.

*Praecip. ad rup. marit.* — *Eur.* : Cornwall., Lusit. *Afr.* : Ins. Azor., Canar., Madeir.

« *subspec. subappressum* Kindb. in *Hedwig*. 1896, p. 66.

Br. appressum (erratum !) ej. *Eur. a. N. Am. Bryin.* p. 356 (1897).

— *Am. sept.* : Washington.

« *var. viride* Husn. *op. et l. cc.*

Ut et cum typ.

*alpinum* { Vill. — Br. Schleicheri.  
Wulf. — Bartramia Halleriana.  
viridissimum et majus, capitulis crassissimis Hall. — Bartramia ithyphylla.

*alte-operculatum* Besch. *Fl. bryol. Réunion, etc.* p. 93 (1880-81).

2. — *Afr.* : Ins. Bourbon., Franciae.

*alticaule* C. M. — *Mniobryum*.

*var. robustior* C. M. — *Mniobryum alticaule var. robustior*.

*altisetum* C. M. in *Hedwig*. 1898, p. 96.

Syn. — *Pac.* : Austral. or. (Victoria).

« *var. humilisetum* C. M. in *op. et l. cc.*

Ubi typ.

*altissimum* C. M. — *Bryum Preussii*.

*alto-limbatum* (Hpe.) Mitt in *Trans. and Proceed. of the roy. Soc. of Victoria*, 1883, p. 73.

*Rhodobryum alto-limbatum* Hpe. in *Linn.* 1869-70, p. 517; *Par. Ind.* p. 1114 (1897).

— *Pac.* : Austral. occid.

*ambiguum* { Duby. — Br. plumosum.  
Kindb. — Webera.

*amblyacis* C. M. in *Hedwig*. 1898, p. 104.

*Rhodobryum amblyacis* Par. *Suppl. Ind.* p. 299 (1900).

2. — *Pac.* : Austral. or. (Victoria).

*Amblyodon* C. M. in *Linn.* 1878-79, p. 293.

Syn. — *Am. merid.* : Argentin. Tucuman.

*amblyolepis* Card. in *Rev. bryol.* 1900, p. 45.

Ster. — *Am. merid.* : Fret, Gerlache.

*amblyphyllum* Philib. in *Rev. bryol.* 1898, p. 263.

1. — *As.* : M. Alatau.

*Amblystegium* Ryan in *Hagen M. Norveg. bor.* p. 164 (1898-99).

1. vel syn. — *Eur.* : Norveg. bor.

*amœnum* Broth. in *Act. Soc. sc. fenn.* 1898, n. 2, p. 25.

2. Terr. — *As.* : Kashmir.

*ammophilum* Ruthe in *Hedwig*. 1897, p. 384.

Br. lacustre  $\beta$  angustifolium ej. in *Baenitz Herb. eur.* 1892, Lief. 67, n. 6724.

In arenos. — *Eur.* : Pomeran.

*amplirete* Rehm. *M. Afr. austr. n.*

— *Afr.* : Orange.

*amplirete* C. M. — *Mniobryum*.



- et ampullis angustioribus* Dill. — Tetraplodon.  
 mnioides.  
*ampullaceum, foliis* { *Serpilli pellucidis, collo crassiore* Dill. — Spla-  
 chnum sphaericum.  
*Thymi pellucidis, collo strictiore* Dill. — Splachnum  
 ampullaceum.
- andicola* Kunth *Syn. pl. Æquinoct. I, p. 58* (1822-28); *Brid. Bryol. univ. I, p. 667* (1826); *C. M. Syn. I, p. 256* (1849).  
*Rhodobryum andicola* Par. *Ind. p. 1114* (1897).  
 2. — *Am. sept.* : Mexic. *Am. merid.* : And. Quitens.  
 et Chilens.
- androgynum* Web. — Aulacomnium.  
 β *prolixum* Roth. — Aulacomnium palustre *var. polycephalum*.
- angeiothecium* C. M. in *Hedwig. 1898, p. 108*.  
*Rhodobryum angeiothecium* Par. *Suppl. Ind. p. 299* (1900).  
 2. — *Pac.* : Ins. Norfolk.
- angermannicum* Arn. in *Rev. bryol. 1898, p. 6*.  
*Syn. Ad. rup. sicc.* — *Eur.* : Suecia centr. or.
- angolense* W. et Dy. — Brachymenium.  
*angulosum* Menz. — Conostomum australe.
- angustifolium* *Brid. Bryol. univ. I, p. 686* (1826); *C. M. Syn. I, p. 309* (1826); *Mitt. M. austr. am. p. 300* (1869).  
 2. — *Am. sept.* : Haïti.
- angustifolium* { *Kaur.* — Br. saltense.  
 { *W. P. Sch.* } — Br. bolivianum.  
 { } — Br. semense.
- angustirete* Kindb. in *Bull. Torr. bot. Club. XVI, p. 94* (1879).  
*Syn. Terr. et ad rup. humid.* — *Am. sept.* : M. Rupestr., Columb.  
 brit.
- angustis* { *pallidis foliis, capitulis erectis longioribus pediculis insiden-*  
 { *tibus* Dill. — *Barbula unguiculata*.  
 { *viridibus foliis, capitulis erectis brevibus pediculis insidentibus*  
 { *Dill.* — *Barbula unguiculata*.
- anisodontacum* Dusen *M. Am. austr. n. 733*.  
*Terr.* — *Am. merid.* : Chile.
- annotinum* { *Brid.* — *Mielichhoferia Jamesoni*.  
 { *Hedw.* — *Webera*.  
 { *lanceolatum pellucidum, capsulis oblongis pendulis* Dill.  
 { — *Webera annotina* (fid. C. M.).  
 { *lanceolatum pellucidum, capsulis oblongis pendulis* Dill.  
 { — *Bryum pallens* (fid. Limpr.)  
 { *palustre, capsulis ventricosus pendulis* Dill. — Br. pseudo-  
 triquetrum.
- annulatum* Hook. f. et Wils. *Fl. antarct. I, p. 134, t. 60* (1844); *C. M. Syn. I, p. 577* (1848); *Handb. of the N. Zeal. Fl. p. 443* (1867).  
 2. *Ad terr. argill.* — *Pac.* : Nov. Seeland. sept. et med., ins. Campbell.
- anoetangiaceum* C. M. et Kindb. *Bidrag. p. 454 (nom., 1890) et in Mac. Cat. p. 130* (1892).  
 1 ? *Ad rup. torr.* — *Am. sept.* : Columb. brit., M. Rupestr.

*Anomodon* Mont. in *Ann. sc. nat.* 1857, p. 151.

2. — *Afr.* : Ins. Sao Anton. prom. Viridis.

*antarcticum* H. f. et W. — Webera.

*apalodictyoides* C. M. in *Bot. Zeit.* 1853, p. 21; *Mitt. M. Ind. or.* p. 74 (1859).

— *As.* : M. Neilgherr.

*apiculatum* Schwaeggr. *Suppl. I, P. II*, p. 102, t. 72, (1816); C. M. *Syn. I*, p. 308 (1849); *Mitt. M. austr. am.* p. 301 (1869) (1).

2. — *Am. merid.* : Brasil. austr. or., Monte-Video.

*apiculatum* { Bryol. eur. (1849). — Br. marginatum.  
Dz. et Mk. — Br. plumosum.  
Wils. — Br. Mildeanum.

*apocarpum* L. — Grimmia.

var. *aureum* Retz. —

var. *incanum* Ehrh. — } Hedwigia ciliata.

var.  $\beta$  L. —

var. *virens* Ehrh. — Grimmia apocarpa.

*apocaulon* Hoffm. — Grimmia apocarpa.

*apophysatum* C. M. in *Nuov. Giorn. bot. ital.* 1896, p. 24.

— *Am. merid.* : And. Boliv.

*appressifolium* Broth. in *Oefv. af Finska Vet.-Akad. Foerh.* 1898, p. 175.

2. Ad rivul. — *Pac.* : Nov. Seeland.

*appressum* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1893, I, p. 116; *Ren. M. Masc. Mad.* n. 27.

2. — *Afr.* : Madagascar.

*appressum* { Kindb. — Br. alpinum *subsp.* subappressum.  
Wils. — Br. turbinatum.

*aptychodes* C. M. — Mniobryum.

*aquaticum* { Gmel. — Cinclidotus.

Hoffm. — Dichodontium pellucidum.

*arachnoideum* C. M. in *Fl.* 1879, p. 378.

Ster. — *Afr.* : M. Kenia, Usambara.

*Araucariae* C. M. in *Ule Bryoth. brasil.* n. 140.

— *Am. merid.* : Brasil. or.

*archangelicum* Bryol. eur. fasc. 32 (Vol. IV), Mon. p. I, t. 333 (1846); C. M. *Syn. I*, p. 271 (1849) et II, p. 569 (1851); W. P. Sch. *Syn. II* ed. p. 416 (1876).

Br. inclinatum plano-operculatum Breidl. in sched.

? Br. subnitidulum Kindb. in *Bot. Not.* 1896, p. 5.

Br. Tauriscorum Limpr. in *LX Jahresb. d. schles. Gesell.* p. 241 (1883).

Syn. Terr. — *Eur.* : Spitzberg, mar. Alb., Dovre, Alp. Salisb., Stir., Carinth., Helvet. (Gemmi). *As.* : Jenissei inf. *Am. sept.* : Groenland., fret. Scoresby, ins. Sabine, Gaspes., M. Rupestr.

*arctico-pallens* Philib. — Br. flavescens.

(1) Cl. Mitten procul dubio errat, quum in *M. Ind. or.* p. 70, *Br. apiculatum* Schwaegr. ad *Br. nitens* Hook. ut syn. ducit.

*arcticum* (R. Br.) *Bryol. eur. l. c., Mon. p. 2, t. 335*; *C. M. Syn. I ed.* p. 269; *W. P. Sch. op. cit. p. 409.*

*Br. nutans* var. *Drumm. M. bor. am. I ed. n. 263 B.*

*Hemisynapsium arcticum* *Brid. Bryol. univ. I, p. 606* (1826).

*Pohlia arctica* *R. Br. Suppl. App. ad iter Parray. p. 197.*

*Syn. Ad terr. humid. et in fiss. rup. — Eur. : Spitzberg, Septem Ins., ins. Ursor., Scotia, Alp. Scand., Tatra, Austr. inf., Salisb., Stir., Carinth., alp. Julian., Tirol., Helvet., Jurass. As. : Penins. Tschutschica, Jenissei. Am. sept. : Fret. Smith, ins. Melville, Groenland, Labrador, Gaspes., reg. sept., occid., M. Rupestr., Calif.*

« *var. helveticum* (Phil.) *Husn. Musc. gall. p. 235* (1889).

*Br. helveticum* *Philib. in Rev. bryol. 1886, p. 83.*

*Ut typ. — Eur. : Helvet.*

« *var. kungeanum* *Philib. in op. cit. 1899, p. 62.*

— *As. : M. Alatau.*

*var. tomentosa* *Jaeg. — Br. tomentosum.*

*arcticum* *Lindgr. — Br. Lindgrenii.*

*arcticum purpurascens* *Aongstr. — Br. purpurascens.*

*arctogaeum* *Hagen in Tromsø Mus. Aarsheft. 1898-99, p. 230.*

*Syn. Ad terr. steril. — Eur. : Norveg. bor.*

*arcuatulum* *Kindb. — Br. arcuatulum.*

*arcuatulum* *Limpr. Einig. neue Art.... 1883, p. 237 et Laubm. Deutschlands.... II, p. 304* (1892).

*Br. arcuatulum* *Kindb. Eur. and N. Am. Bryin. p. 373* (1897).

*Syn. Ad rup. madid. umbros. — Eur. : Norveg. Am. sept. : Fret. Scoresby.*

*arcuatulum* *H. f. et W. — Eccremidium.*

*ardonense* *Breidl in Österr. bot. Zeitschr. 1889, n. 4.*

1. — *As. : Caucas.*

*Arenae* *C. M. in Fl. 1885, p. 402.*

2. *Paludic. — Am. merid. : Fret. Magell.*

*arenarium* *Jur. Laubmfl. Österr.-Ung. p. 273* (1882).

*Br. atro-purpureum* } *Garov. Enum. p. 28* (1840).

*var. arenarium* *Husn. Musc. gall. p. 244* (1889).

2. *Limic. — Eur. : Austr. infer., Salisb., Gall. centr. (Bourbonnais).*

*arenarium* } *C. M. — Br. Pootenianum.*

} *Saut. — Br. Blindii.*

*areoblastum* *C. M. in Fl. 1886, p. 280.*

2. *Terr. — Afr. : Cameroon, ins. Fernando Po, Sao Thomé.*

*argentatum* *C. M. in Engler's Bot. Jahrb. 1883, p. 83 et in Forschungsreise.... Bot. p. 41* (1869).

*Ster. — Afr. : Ins. Ascension.*

*argenteum* *L. Sp. pl. p. 1120* (1753); *Brid. Bryol. univ. I, p. 657* (1826); *Bryol. eur. fasc. 6-9 (Vol. IV), Mon. p. 78, t. 384* (1839); *C. M. Syn. I, p. 314* (1849); *W. P. Sch. op. cit. p. 448.*

*Argyrobryum argenteum* *Kindb. Laubm. Schw. u. Norw. p. 78* (1883).

*Br. capitulis subrotundis reflexis, cauliculis teretibus argenteis* *Dill. Cat. Giss. p. 226* (1718).

- Br. pendulum* julaceum, argenteum et sericeum *Dill. Hist. M.* p. 392, t. 50, f. 62 (1741) *et hb.*  
*Hypnum argenteum* *Schrank, Bayer. Fl.* p. 469 (1789).  
*Mnium argenteum* *Hoffm. Deutsch. Fl. II*, p. 51 (1796).
2. Fere ubique, except. arbor., per tot. fere terrar. orbem, a Spitzberg. et Groenland. ad terr. antarct.
- var. *argyrotrichum* Mitt. — *Br. argyrotrichum*.  
 var. *australe* C. M. — *Br. oranicum*.
- « *f. ceylonensis* *Fleisch. M. Archip. ind. n.* 144.  
 Terr. — *As.* : Ceylan.
- var. *corrugatum*. — *Br. corrugatum*.
- « var. *costa-ricense* *Ren. et Card. in Bull. Soc. roy. bot. Belg.* 1892, I, p. 167.  
 Ad mur. et terr. — *Am. sept.* : Costa-Rica.
- var. *julaceum* Bland. — *Br. argenteum* var. *majus*.
- « var. *lanatum* (P. B.) *Bryol. eur.*, C. M. et W. P. Sch. II, cc.  
*Br. lanatum* *Brid. Sp. M. III*, p. 20 (1877).  
*Mnium lanatum* P. B. *Prodr.* p. 75 (1805).  
 In mont. editor. ut in reg. campestr. zon. merid. — *Eur.* : Alp. Tirol, etc. *Afr.*, utraque. *Amer. et Austral.*
- var. *lanatum* W. P. Sch. (Pl. Boliv.). — *Br. capillipes* var. *dimorphum*.
- « var. *majus* *Schwaeagr. Suppl. I. P. II*, p. 88 (1816); *Bryol. eur.*, C. M. et W. P. Sch. II, cc.  
*Br. argenteum* var.  $\left\{ \begin{array}{l} \text{julaceum Bland. Uebers. d. Mecklenb. M. p. (1809).} \\ \text{muticum W. P. Sch. mss.; Lindb. Spetsb. M. p. 516 (1866).} \end{array} \right.$   
*Br. julaceum* *Schrad. Spic. Fl. germ.* p. 70 (1794).  
 Loc. humidior. — *Eur.* et *Am. sept.* : Cum typ.
- var. *muticum* W. P. Sch. — *Br. argenteum* var. *majus*.
- « var. *submuticum* *Besch. Fl. bryol. Réunion, etc.*, p. 93 (1880-81).  
 In carbonar. — *Afr.* : Ins. Bourbon.
- argentisetum* C. M. in *Fl.* 1890, p. 478.  
 Ster. — *Afr.* : Kilima N'Djaro.
- argillicola* *Broth. in Oefv. af Finska Vet.-Akad. Foerh.* 1899, p. 102.  
 2. Ad terr. argill. — *Pac.* : Tasman.
- argyreum* C. M. in *Linn.* 1880-83, p. 380.  
 2. — *Am. merid.* : Uruguay.
- argyroglyphodon* *Philib. in Rev. bryol.* 1900, p. 51.  
 1. Terr. — *As.* : M. Alatau.
- argyrotrichum* (Mitt.) C. M. in *Linn.* 1875, p. 385.  
*Br. argenteum* var. *argyrotrichum* Mitt. in *Journ. of the Linn.* 1886, p. 307.  
 Terr. — *Afr.* : Niam-Niam, Usagara, Usambara.
- « var. *laxum* C. M. in *op. et l. cc.*  
 Ut typ. — *Afr.* : Niam-Niam.
- aristatum* *Hagen in Tromsøe Mus. Aarsheft.* p. 183 (1898-99).  
 Syn. Humic. — *Eur.* : Norveg. bor.

*aristatum* Dicks. — *Barbula unguiculata*.

*Arnellii* Par. Ind. p. 163 (1894).

Br. alandicum *J. O. Bomanss. in Rev. bryol.* 1896, p. 90.

Br. flexisetum *Lindb. et Arn. M. As. bor.* 1890, II, p. 44.

Syn. Ad terr. humid. — *Eur.* : Ins. Aland. *As.* : *Jenissei* infer.

*Arnellii* Bomanss. — Br. halophytum.

*articulatum* Sendn. — Br. Funckii.

*Arvenii* Arn. in *Bot. Not.* 1897, p.

2. Ad terr. calcar. — *Eur.* : Suecia.

*arvorum* Stroem. — *Ditrichum tortile* var. *pusilla* (fid. *Hagen*).

*Aschersoni* C. M. in *Fl.* 1894, p. 31.

2. fr. ign. In foss. aquat. — *Afr.* : Oasis Dachel.

*astorense* Broth. — *Anomobryum*.

*aterrimum* C. M. in *Rehm. M. Afr. austr.* n. 235.

— *Afr.* : Prom. B. Sp.

*atlanticum* Solms Laub. — Br. alpinum var. *angustifolium*.

*atro-purpureum* *Wahlenb. in Web. et Mohr Ind. M.* (1803) p.p.; *Bryol. eur. fasc.* 6-9 (*Vol. IV*), *Mon.* p. 73, t. 378 (1839); *C. M. Syn. I*, p. 306 (1849); *W. P. Sch. Syn. II ed.* p. 438 (1876).

*Bryum balanoides* *Tayl. in Drumm. M. Swan River* (fid. *Mitt. in Kew Journ. bot.* 1856, p. 261).

Br. bicolor *Dicks. Pl. crypt. fasc. IV*, p. 16 (1801) fid. *Lindb.*

Br. caespitium  $\beta$   $\left\{ \begin{array}{l} \text{atro-purpureum } \textit{Bland. Mechl. Laubm. fasc. V,} \\ \text{n. 224 (1808).} \\ \text{bicolor } \textit{Brid. Sp. M. III, p. 27 (1817).} \end{array} \right.$

Br. carneum  $\beta$   $\left\{ \begin{array}{l} \text{atro-purpureum } \textit{Schwaegr. Sp. M. p. 67 (1830).} \\ \text{pusillum } \textit{Schultz. Suppl. Fl. Starg. p. 84 (1819).} \end{array} \right.$

Br. dichotomum *Wils. Bryol. brit. p. 244, t. 50 (1855)*; *H. f. et W. Fl. of the N. Zeal. II*, p. 85 (1855).

Br. erythrocarpum  $\left\{ \begin{array}{l} \textit{Brid. Sp. M. III, p. 18 (1817).} \\ \beta \text{ bicolor } \textit{ej. Bryol. univ. I, p. 655 (1826).} \end{array} \right.$

Br. gracilentum *Tayl. mss.*; *C. M. Syn. II*, p. 576 (1850).

Br. pendulum ovatum, caespitium et pilosum, seta bicolore *Dill. Hist. M.* p. 396, t. 50, fig. 66 A, F et G (1741).

Br. pendulum surculis teretibus viridibus *ej. op. cit. t. 50, f. A, B et E, et hb.*

Br. trichodes, capitulis reflexis, pediculis ima medietate rubris, summa luteo-viridibus *ej. Cat. Giss. p. 226 (1718)*.

Br. trichodes, capitulis reflexis, viridissimum holosericeum *ej. op. et l. cc.* *Mnium bicolor* *P. B. Prodr. p. 74 (1805)*.

*Mn. bulbiliosum* *Mont. in Ann. sc. nat. II ser. XVI, p. 268 (1841)*.

*Mn. dichotomum* *P. B. op. et l. cc.*

2. Muric., terr. et ad. aren. marit. — *Eur.* : Zon. merid., rar. intermed., rariss. septentr. *As.* : *Caucas.*, Minor. *Afr.* : *M. Sinai*, *Alger.*, ins. *Canar.*, *Madeir. Am. sept.* : *Niagara*, *Nevada*, *Illinois*, *Tennessee*, *New-York*, *Pensylv.*, *Virgin.*, *Florid.*, *Alabama*. *Pac.* : *Austral.*, *Nov. Seeland*.

var. *arenarium* *Husn.* — Br. *arenarium*.

« var. *macrothecium* *Besch. Cat. M. Alger. p. 25 (1882)*.

— *Afr.* : *Alger*.

*atro-purpureum*  $\left\{ \begin{array}{l} \text{de Not. — Br. murale.} \\ \text{Garov. — Br. arenarium.} \\ \text{Wahlenb. p.p. — } \textit{Mniobryum.} \end{array} \right.$

*atro-sanguineum* C. M. in *Linn.* 1880-83, p. 386.

2. Ad mur. et terr. — *Am. merid.* : *Argent.* subtropic.

*atrothecium* C. M. — *Webera*.

*atro-virens* Vill. — *Br. erythrocarpum*.

*attenuatum* Dicks. } (fid. C. M.) — *Br. pallens* var. *abbreviatum*.  
 } (fid. Lindb.) — *Entosthodon Templetoni*.

*Atwateriae* C. M. in *Fl.* 1873, p. 76; *Lesq. et Jam. Man.* p. 234 (1884).

*Br. Macounii* Aust. in *Coult. Bot. Gaz.* II, p. 110.

2. Ad rup. humid. — *Am. sept.* : Ontario, Oregon, Washington, Columb. brit., ins. Vancouver, Texas, Calif.

*Auberti* (*Schwaegr.*) *Brid. Mant. M.* p. 119 (1819); *C. M. Syn. I*, p. 262 (1849).

*Polla Auberti* *Brid. Bryol. univ. I*, p. 711 (1826).

*Mnium Auberti* *Schwaegr. Suppl. I, P. II*, p. 132, t. 80 (1816).

*Syn.* — *Afr.* : Ins. Borbon., *Franciae*.

*Auberti* } *Hsch.* — *Br. schistoneuron* var. *decumbeus*.

} *Schwaegr.* — *Br. schistoneuron* var. *densifolium*.

*aulacomnioides* C. M. in *Hedwig.* 1899, p. 72.

*Ster.* In turf. — *Afr.* : Prom. B. Sp.

*auratum* Mitt. — *Anomobryum*.

*aureum* } *Huds.* — *Leptobryum* *piriforme*.  
 } *capitulis reflexis piriformibus, calyptra quadrangulari, foliis*  
*in bulbi formam congestis* Dill. — *Funaria hygrometrica*.

*auricomum* *Besch. Fl. bryol. Réunion, etc.* p. 99 (1880-81).

2. — *Afr.* : Ins. Borbon.

*auriculatum* C. M. in *Linn.* 1875, p. 379.

*Ster. Terr.* — *Afr.* : Niam-Niam.

*australe* *Hpe. Ic. M.* t. 26 (1844); *C. M. Syn. I*, p. 285 (1849).

2. — *Pac.* : Austral. or.

♂ var. *minus* *Hpe. in Linn.* 1852, p. 714.

— *Pac.* : Austral. or. (Victoria).

*austro-albicans* C. M. — *Mniobryum*.

*austro-alpinum* C. M. in *Hedwig.* 1898, p. 99.

*Ster.* — *Pac.* : Austral. or. (Victoria).

*austro-bimum* *Broth. in Oefo. af Finska Vet.-Akad. Foerh.* 1893, p. 177.

2. In fiss. rup. sicc. — *Pac.* : Nov. Seeland.

*austro-caespitium* C. M. in *Engler's Bot. Jahrb.* 1883, p. 77 et in *Forschungsreise... Bot.* p. 13 (1889).

*Syn.* — *Afr.* : Ins. Kerguelen.

*austro-crudum* C. M. — *Webera*.

*austro-elongatum* C. M. — *Webera*.

*austro-nutans* C. M. (Ins. Kerg.) — *Webera pseudo-nutans*.

*austro-pallescent* *Broth. in op. et l. cc.*

1. Ad rivul. — *Pac.* : Nov. Seeland.

*austro-polare* *Card. in Rev. bryol.* 1900, p. 45.

*Ster.* — *Am. merid.* : Fret. Gerlache.

*austro-polymorphum* C. M. — *Webera*.

- austro-turbinatum *C. M. in Linn.* 1880-83, p. 368.  
 Ster. — *Am. merid.* : Argentin. Cordob.
- austro-ventricosum *Ren. Prodr. fl. bryol. Mad. etc.* p. 294 (1897).  
 Ster. — *Afr.* : Madagascar.
- autoicum *Arn. in Rev. bryol.* 1898, p. 2.  
 1. — *Eur.* : Alp. Norveg.
- autumnale *Limpr. Einige neue Art...* 1883, p. 239 et *Laubm. Deutschl...*  
*II*, p. 312 (1892).  
 Syn. In aren. humid. — *Eur.* : Norveg. bor.
- Axel-Blyttii *Kaur. in Bot. Not.* 1889, n. 2 ; *Philib. in Rev. bryol.* 1889,  
 p. 61.  
 1. — *Eur.* : Norveg.
- axillare *Philib. in op. cit.* 1898, p. 61.  
 1. — *As.* : M. Alatau.
- badium *Bruch ut syn. in Brid. Bryol. univ. I*, p. 850 (1826) ; *W. P. Sch. op. cit. p.* 444.  
 Br. caespitici. m  $\left\{ \begin{array}{l} \text{C. M. Syn. I, p. 284 p.p. (1849).} \\ \gamma \text{ badium } \textit{Brid. Bryol. univ. l. c.} \\ \beta \text{ gracillimum } \textit{Bryol. eur. fasc. 659 (Vol. IV), Mon.} \\ \text{p. 71, t. 375 } \beta \text{ (1839).} \end{array} \right.$   
 Br. tectorum *de Not. in sched. fig. Jaeg. Ad. I*, p. 632 (1873-74).  
 2. In arenos. humid. — *Eur.* : Zon. intermed. *As.* : Caucas. occid. et centr.
- Haenitzii *C. M. in Fl.* 1888, p. 417 (1).  
 Syn. — *Eur.* : Norveg. bor.
- Baileyi *Broth. in Bail. Syn. of the Queensl. Fl. IV (nom., 1890) et in Oefv. af Finska Vet.-Akad. Foerh.* 1890, p. 100.  
 2. In lapidos. — *Pac.* : Austral. or. (Victoria).
- balanocarpum *Besch. in Bull. Soc. bot. Fr.* 1894, p. 82.  
 1. Terr. — *As.* : Tonkin.
- balanoides *Tayl.*  $\left\{ \begin{array}{l} \text{— Br. atro-purpureum (fid. Mitt.).} \\ \text{— Br. pachythea (fid. H. f. et W.).} \end{array} \right.$
- baldense *Vent. in de Not. Epil.* (1869) et in *Rev. bryol.* 1881, p. 31.  
 2. Ad rivul. — *Eur.* : M. Baldo, Valesia.
- barbatum  $\left\{ \begin{array}{l} \text{Curt. — } \textit{Barbula unguiculata.} \\ \text{Wils. } \left\{ \begin{array}{l} \text{— Br. capillare var. carinthiacum (fid. Arn.).} \\ \text{— Br. Stirtoni (fid. Limpr.).} \end{array} \right. \end{array} \right.$
- barbulaceum *C. M. in Linn.* 1875, p. 389.  
 Br. coronatum *Wils. in Gardn. M. brasil. n.* 35 ; *C. M. Syn. I*, p. 307 p.p. (1849).  
 2. — *Am. merid.* : Brasil.
- barbuloides *C. M.* — *Brachymenium.*
- Barnesi *Wood in W. P. Sch. Syn. II ed. p.* 471 (1876).  
 Br. argenteum *(fid. Braithw. Brit. Moossf. II, p.* 176).  
 Br. lacu-tre forma bulbillifera *Limpr. Laubm. Deutschl... II*, p. 317 (1892).  
 Ster. In coll. arenac. — *Eur.* : Anglia (Westmoreland).

(1) « Forsan *Br. subglobosum* Schlieph. » *Limpr. op. cit. p.* 356. — « *Br. pallescens* var. » *fid. Hagen.*

- Baroni* Wright. — *Brachymenium*.
- Bartramia* { Gmel. — *Bartramia* Halleriana.  
                   { Schwaegr. — *Cryptopodium bartramioides*.
- bartramioides* Rehm. *M. Afr. austr.* n. 257.  
                                   — *Afr.* : Orange.
- bartramioides* Hook. — *Cryptopodium*.
- bartramiopsis* C. M. in *Linn.* 1880-83, p. 372.  
     Ster.                     — *Am. merid.* : Uruguay.
- Bateae* C. M. in *Linn.* 1898, p. 93.  
     Ster.                     — *Pac.* : Austral. or. (Nov. Vales.).
- Baueri* Hpe. in *Linn.* 1869-70, p. 457.  
                                   — *Am. sept.* : S. Nevada Calif.
- Bauerleni* C. M. in *Hedwig.* 1898, p. 98.  
     Ster.                     — *Pac.* : Austral. or. (Nov. Vales.)
- Beccarii* C. M. — *Webera*.
- Bellii* C. M. in *Hedwig.* 1898, p. 93.  
     2. Ad rad. arb. — *Pac.* : Nov. Seeland.
- Benguetiae* (C. M.) Par. *Suppl. Ind.* p. 57 (1900).  
     *Sclerbryum Benguetiae* C. M. in *Micholitz Pl. Philipp.* n. 176.  
                                   — *As.* : Ins. Philippin. (Luzon).
- bergoense* J. O. Romanss. in *Rev. bryol.* 1899, p. 12 et in *Act. Soc. p. Faun. et Fl. fenn.* 1899, n. 4.  
     Syn. Ad. litt. mar. — *Eur.* : Suecia, ins. Aland.
- Bernoullii* C. M. in *Bull. hb. Boissier*, 1897, p. 183.  
                                   — *Am. sept.* : Guatemala.
- Bescherellei* Jaeg. *Ad. I*, p. 627 (1873-74).  
     *Br. erythrocarpoides* W. P. Sch. in *Besch. Fl. bryol. Nov. Caled.* p. 214 (1873).  
     *Br. torulosicolum* C. M. in *Hedwig.* 1898, p. 90 nom.  
     Syn.                     — *Pac.* : Nov. Caledon.
- Bescherellei* R. C. — *Br. Rodriguezii*.
- betulinum* Kaur. ap. Joergens. in *Christiania Vid. Selsk. Foerh.* 1894, p. 86.  
                                   — *Eur.* : Norveg.
- Beyrichianum* C. M. { — *Rhodobryum*.  
                                   { — *Rhodobryum Swartzianum*.
- bicolor* C. M. in *Fl.* 1890, p. 476.  
     2. In palud. edit. — *Afr.* : Kilima N'Djaro.
- bicolor* Dicks. { — *Br. atro-purpureum* (fid. Limpr.).  
                                   { — *Br. erythrocarpon* (fid. Milde).
- Biddlecomiae* Aust. in *Coult. Bot. Gaz.* II, p. 110; *Lesq. et Jam. Man.* p. 226 (1884).  
     *Br. Edwardsianum* C. M. et Kindb. in *Bidrag* nom. 1890 et in *Mac. Cat.* p. 120 (1892).  
     *Br. warneum* *Mac. Canad. M.* n. 383.  
     Syn. Ad terr. aren. humid. — *Am. sept.* : Ins. prine. Edouard, Colorado.



*Bigelowii* Sull. — *Webera*.

*bigibbosum* Besch. in *Bull. Soc. bot. Fr.* 1898, p. 65.

2. — *Pac.* : Taiti.

*Billeti* Besch. in *Rev. bryol.* 1898, p. 73.

2. Ad terr. argill. humid. — *As.* : Tonkin.

*Billardieri* *Schwaeagr. Suppl. I, P. II, p. 115, t. 76* (1816); *Brid. Bryol. univ. I, p. 667* (1826); *C. M. Syn. I, p. 253* (1849).

*Br. rufescens* H. f. et W. ? (*fid. Mitt.*).

Ad arb. et rup. humid. — *Pac.* : Austral., Nov. Belg., ins. Lord Howe, Tasman.

*Billardieri* } *Bryol. eur.* — *Br. provinciale*.

              } Hpe. (M. Preiss.). — *Rhodobryum pallenticoma* (*fid. C. M.*).  
*bimo aff.* Wils. — *Br. pseudo-triquetrum*.

*bimoideum* de Not. *Epil. p.* 383 (1869).

2. ♂ ign. — *Eur.* : Vall. Tell.

*bimum* Schreb. *Spic. fl. lips. p.* 83, n. 1047 (1771); *Brid. Bryol. univ. I, p. 678* (1826); *Bryol. eur. fasc. 6-9 (Vol. IV), Mon. p. 50, t. 363* (1839); *C. M. Syn. I, p. 257* (1849); *W. P. Sch. op. cit. p. 430*.

*Br. lanceolatum bimum*, setis et capsulis longis pendulis *Dill. Hist. M. p. 405, t. 51, fig. 73 F* (1741) et *hb.*

*Br. pseudo-triquetrum* β *bimum* *Schwaeagr. Suppl. I, P. II, p. 211* (1816).

*Br. turbinatum* *Drumm. M. bor. am. I ed. n. 266*.

*Br. ventricosum* β *Hook. et Tayl. Musc. brit. p. 124* (1818).

*Hypnum palustre erectum breve*, foliis brevibus, tenuibus, angustis, minus confertis *Dill. Cat. Giss. p. 220* (1718).

*Mnium bimum* *Brid. Musc. rec. II, P. III, p. 93* (1803).

*Syn.* In pratis uligin. et ad rup. madid. — *Eur.* : Reg. campestr. et montan., rar. alp. zon. intermed. et sept.; ins. Melit. *As.* : Amour, Caucas. *Afr.* : Ins. Kerguelen. *Am. sept.* : Franz Joseph Fjord, Terra nov., Canada merid.; Alaska, ins. Vancouver, Columb. brit., Washington, Oregon, Nevada, Illinois, Wisconsin, Kansas, reg. centr. et or. *Am. merid.* : And. Nov. Granat. et Quitens. *Pac.* : Nov. Seeland., Tasman.

« *var. amœnum* Warnst. in *Verh. d. bot. Ver. Prov. Brandenb.* 1899, p. 60.

— *Eur.* : Lusat. infer.

« *var. angustifolium* *Kindb. in Mac. Cat. p. 123* (1892).  
*Rupic.* — *Am. sept.* : Ontario.

« *var. atrotheca* *Ren. et Card. in Bot. Gaz.* 1894, p. 238.  
— *Am. sept.* : Terr. nov.

« *f. aulacodon* *Hagen in Tromsø Mus. Aarsheft.* 1898-99, p. 168.  
In sphagnet. — *Eur.* : Norveg. bor.

« *var. brevisetum* *Warnst. in op. et l. cc.*  
— *Eur.* : Lusat. infer.

*var. cuspidatum* *Bryol. eur.* — *Br. cuspidatum*.

« *var. longicollum* *Warnst. Moossft. Prov. Brandenb. p. 52* (1885).  
— *Eur.* : Brandeb.

« *var. subnivale* *Mol. Alg. Alp. p. 78* (1865).  
— *Eur.* : Alp. Algov.

*bipartitum* Dicks. — *Ceratodon purpureus*.

- blandum* Hook. f. et W. in Journ. of bot. 1844, p. 546 et Fl. antarct. I, p. 134, t. 60 (1844); C. M. Syn. I, p. 348 (1849) et II, p. 577 (1851); Handb. of the N. Zeal. Fl. p. 441 (1867).  
2. Paludic. — Pac. : Nov. Seeland. sept. et med., ins. Campbell.  
« var. *luridum* Hook. f. et Wils. Fl. Tasman. p. 190 (1860).  
— Pac. : Nov. Seeland., Tasman., ins. austr.
- Blindii* Bryol. eur. fasc. 32 (Vol. IV), Mon. Suppl. I, p. II, t. 383 (1846); C. M. Syn. I, p. 298 (1849); W. P. Sch. op. cit. p. 447.  
*Argyrobryum Blindii* Kindb. Art. d. Laubm. Schwed. u. Norveg. p. 78 (1883).  
Br. *arenarium* Saut. mss. (1836) in Fl. Salz. III, p. 46 (1870).  
Br. *Funkii* var. ? Bryol. eur. Mon. Bryt. p. 77 1839.  
Br. *Funkii* var. *subsphaericum* W. P. Sch. in Husn. Muscol. gall. p. 243 (1889).  
Br. *Kiaerii* Lindb. M. scand. p. 41 (1879).  
2. In terr. arenos. vel argill. aren. et tophac. — Eur. : Salisb., Stir., Carinth., Pinzgov., Tirol., alp. Rhaet., Helvet. Am. sept. : M. Rupestr. et Selkirk.  
var. *Payoti* Husn. — Br. Payoti.
- Bohnhofii* C. M. in Fleisch. M. Archip. ind. n. 145.  
Ster. Ad scaturig. — As. : Ceylan
- Boissieri* de Not. — *Webera longicolla*.  
*Bolanderi* Lesq. — *Webera*.
- bolivianum* Par. Ind. p. 167 (1894).  
Br. *angustifolium* W. P. Sch. in Mandon Pl. Boliv. n. 1605.  
— Am. merid. : And. Roliv.
- Bommanssonii* Lindb. in Act. Soc. pr. Faun. et Fl. fenn. 1884, p. 183.  
— Eur. : Fennia.
- boreale* Schwaegr. — Br. *pallens* var. *boreale*.
- boreum* Hagen in Tromsø Mus. Aarsheft. p. 228 (1898-99).  
Syn. — Eur. : Norveg. bor.
- bornholmense* Winkelm. et Ruthe in Hedwig. 1899, Beibl. p. 120.  
2. In sphagnet. — Eur. : Ins. Bornholm.
- Pottorii* C. M. in Linn. 1874, p. 622.  
2. Lignic. — Am. sept. : Mexic.
- brachyacris* C. M. mss.; Mitt. in Trans. and Proced. of the roy. Soc. of Victoria, 1882, p. 73.  
— Pac. : Absq. loco (Austral. vel Tasman.).
- brachycarpum* J. O. Bomanss. in Act. Soc. sc. p. Faun. et Fl. fenn. 1899, n. 4.  
Syn. Ad litt. mar. — Eur. : Ins. Aland.
- brachycarpum* Garov. — *Webera polymorpha* var. *brachycarpa*.
- brachycladulum* C. M. in Hedwig. 1898, p. 98.  
2. — Pac. : Nov. Seeland. sept. occ.
- brachydontium* Hpe. — *Webera*.
- brachymeniaceum* C. M. in Rehm. M. Afr. austr. n. 378 et in Hedwig. 1899, p. 71.  
1. Inter frutic. — Afr. : Prom. B. Sp.
- brachymeniopsis* C. M. — *Anomobryum*.

- brachyneuron* *Kindb. in Ott. Nat. V, p. 180.*  
 Syn. — *Am. sept.* : Fret. Behring.
- brachyphyllum* *Mitt. M. austr. am. p. 302 (1809).*  
 2. — *Am. merid.* : And. Quitens.
- brachypodium* *C. M. — Brachymenium.*
- brachystegium* *C. M. in Ule Bryoth. bras. n. 144.*  
 Terr. — *Am. merid.* : Brasil. or.
- brachyteciella* *C. M. in Hedwig. 1898, p. 91.*  
 2. — *Pac.* : Austral. or. (Nov. Vales.).
- brasiliense* *Hpe. Symb. 1877, p. 725.*  
 2. — *Am. merid.* : Brasil. austr. or.
- Breidleri* *Mac. — Webera Ludwigii.*
- Breutelii* *C. M. — Aplodontium.*
- brevicaule* *W. P. Sch. nss. ; Mitt. op. cit. p. 303.*  
 Br. Liebmannianum *C. M. Syn. II, p. 573 (1851).*  
 2. — *Am. sept.* : Mexic.
- brevicaule* } *Hpe.* } *Brachymenium.*  
 } (Pl. Austr.) *Br. curtum.*  
 } *Vill. — Grimmia commutata.*
- brevicoma* *Hpe. Enum. M. Brasil. p. 31 (1879).*  
 — *Am. merid.* : Brasil. austr. or.
- breviflorum* *Gmel. — Ceratodon purpureus.*
- brevifolium* *Dicks. —* } *Barbula fallax var. brevifolia.*  
 } *Didymodon tophaceus var. brevifolius.*
- brevigemmatum* *Dusen M. Am. austr. n. 586.*  
 Saxic. — *Am. merid.* : Patag. occid.
- breviramulosum* (*Hpe*) *Mitt. in Trans. a. Proced. of the roy. Soc. of Victoria, 1883, p. 73.*  
*Rhodobryum breviramulosum* *Hpe in Linn. 1876, p. 311 ; Par. Ind. p. 115 (1897).*  
 — *Pac.* : Austral. or. (Victoria), Tasman.
- Bridelianum* *C. M. — Webera.*
- Brownianum* *Dicks. — Tetradontium.*
- Brownii* (*R. Br.*) *Bryol. eur. fasc. 32 (Vol. IV), Mon. Suppl. p. 3, t. 337, excl. f. 14 (1846) ; C. M. Syn. I, p. 273 (1849) ; W. P. Sch. op. cit. p. 411.*  
*Hemisynapsium bryoides* *Brid. Bryol. univ. I, pp. 605 et 840 (1826).*  
*Pohlia bryoides* *R. Br. in Suppl. App. ad iter Parray. p. 196.*  
 Syn. In aren. sæpe inund. terraq. turfac. — *Eur.* : Alp. Norveg.  
*Am. sept.* : Franz Joseph Fjord, ins. Melville, Labrador, Terr. nov.,  
 M. Rupestr.
- var. *minor* *Kaur. — Br. stenocarpum.*
- « var. *terskeanum* *Philib. in Rev. bryol. 1899, p. 70.*  
 — *As.* : M. Alatau.
- brunneidens* *C. M. in Hedwig. 1898, p. 106.*  
*Rhodobryum brunneidens* *Par. Suppl. Ind. p. 219 (1910).*  
 2. — *Pac.* : Austral. or. (Victoria).

- brunnescens* } Ren. — *Br. torquescens* f. *monoica*.  
 } Schleich. — *Br. subrotundum*.  
*bulbiferum* Neck. — *Phaescum cuspidatum*.  
*bulbifolium* Lindb. mss.; Norrlin in *Bot. Not.* 1873, n. 9; *Not. Saellsk. pr. Faun. et Fl. fenn. Foerh. N. ser., X Heft*, p. 297 (1874).  
 Syn. In argill. et aren. humid. — *Eur.*: Lappon. Tornens.  
*bulbiforme aureum, calyptra quadrangulati, capsulis piriformibus nutantibus* Dill. — *Funaria hygrometrica*.  
*bulbillicaule* C. M. in *Fl.* 1879, n. 24.  
 Ster. — *Afr.*: M. Kenia.  
*bulbilliferum* Rehm. *M. Afr. austr.* n. 252.  
 — *Afr.*: Prom. B. Sp.  
*bulbillinum* C. M. in *Linn.* 1880-83, p. 384.  
 2. Saxic. — *Am. merid.*: Argentin. Tucuman.  
*bulbillosum* Mont. in *Ann. sc. nat. XVI, II*, p. 268 et *Syll.* p. 33 (1856);  
 C. M. *Syn. I*, p. 301 (1849).  
*Br. dichotomum* (fid. Mitt.).  
 2? — *Am. merid.*: Brasil. austr. or. *Pac.*: Nov.  
 Seeland.?  
*bullatum* C. M. — *Anomobryum*.  
*bullosum* C. M. in *Dusen M. Camer. n.* 44; *Broth. in Engler's Bot. Jahrb.* 1897, p. 246.  
 2. fr. ign. Rupic. in rivul. — *Afr.*: Cameroon.  
*caesium* Vill. — *Ditrichum glaucescens*.  
*caespiticiforme* de Not. *Epil.* p. 398 (1869).  
 1. — *Eur.*: Summ. M. Vall. Aoste.  
*caespiticioides* C. M. in *Hedwig.* 1898, p. 89.  
 2. — *Pac.*: Tasman.  
*caespitium* L. *Sp. pl.* p. 1121 (1753) non hb.! *Brid. Bryol. univ. I*, p. 669 (1826); *Bryol. eur. fasc. 6-9 (Vol. IV), Mon.* p. 70 excl. var.  $\beta$ , t. 374 et 375 f.  $\gamma$  (1839); C. M. *Syn. I*, p. 284 p.p.; W. P. Sch. op. cit. p. 443.  
*Br. caespitium* var. { *elongatum* Schultz in *Fl. Syll. II*, p. 188 (1828).  
 { *nigricans* Brid. *Sp. M. III*, p. 27 (1817)?  
 { *tenerimum* Schultz op. et l. cc.  
*Br. flaccidum* H. f. et W. *Fl. of the N. Zeal. II*, p. 85? (1855).  
*Br. nigricans* Crome *Nachl. I*, n. 74 (1805?).  
*Br. pachypoma* Wils. in *Kew Journ. of bot. IX*, p. 361 (1857).  
*Br. pendulum ovatum, caespitium et pilosum, seta bicolori* Dill. *Hist. M.* p. 396, t. 50, f. 66 B-D (1741) et hb.  
*Br. trichodes, capitulis reflexis, pediculis ima medietate rubris, summa luteo-viridibus* Dill. *Cat. Giss.* p. 226 (1718) (1).  
*Hypnum caespitium* Schrank, *Bayer. Fl. II*, p. 473 (1789).  
*Mnium caespitium* Hedw. *Fund. M. II*, p. 94 (1782).  
*Mn. nigricans* Crome op. et l. cc. excl. syn.?  
 2. Ad mur., sax. et terr. — *Eur.*: Vulgatiss. *As.*: Japon., Saghalien, Taimyr, Obi, Jenissei med., Jakoutsch, Khasia, Himalaya occid., Kashmir, Gilgit, Caucas., Minor. *Afr.*: Occid. et merid. *Am. sept.*: Tota a Groenland. et Columb. brit. ad Florid. et Ludovic. *Pac.*: Ins. Hawaï, Nov. Seeland., Tasman.

(1) *Br. atro-purpureo* jam a cl. Limpricht ut syn. datum!

- *var. atlanticum* Card. in *Ann. Rep. of the Missouri Bot. Gard.* 1897, p. 62.

— *Afr.* : Ins. Azor., Madeir.

*var. atro-purpureum* Bland. — *Bryum*.  
*var. badium* Bryol. eur. — *Bryum*.  
*var. bicolor* Brid. — *Br. atro-purpureum*.  
*var. comense* Husn. — *Br. comense*.

- *var. crinitum* H. f. et W. *Fl. Tasman.* II, p. 191 (1860).  
 — *Pac.* : Tasman.

*var. gracillimum* Bryol. eur. — *Br. badium*.  
*var. hemisphaericum* Brid. — *Br. caespitium* ?

- *var. imbricatum* Bryol. eur. et W. P. Sch. II, cc.

*Br. caespitium var. Kunzei* Warnst. *Moossfl. Prov. Brandenb.* p. 53 (1885).

*Br. Kunzei* H. et H. in *Fl.* 1819, I, p. 90; *Brid. Bryol. univ.* I, p. 661 (1826).

*Br. tectorum de Not. f. d. Jaeg. Ad. I*, p. 632 (1873-74).

In mur. aprie. sicc. — *Eur.* : Cum typ.

*var. Kunzei* Warnst. — *Br. caespitium var. imbricatum*.

- *var. laxum* Warnst. in *Abh. d. bot. Ver. d. Prov. Brandenb.* 1900, p. 206.

In lapidos. — *Eur.* : Boruss. or.

*var. nigricans* Brid. — } *Br. caespitium* ?  
*var. rupestre* Brid. — }

- *var. strangulatum* Warnst. in *op. et l. cc.*  
 Muric. — *Eur.* : Boruss. or.

*var. subsphaericarpum* } Brid. — *Br. caespitium* ?  
*var. tenerum* }

*caespitium* { C. M. p.p. — *Br. badium*.  
 Drumm. — *Br. pendulum*.  
 Hsch. -- *Br. congestum*.  
 Schwaegr. { — *Br. cirratum*.  
 Sull. — *Br. micro-capillare*.  
 Wils. { — *Br. subrotundum*.  
 — *Br. tibetanum*.

*calcareum* Vent. in *Rev. bryol.* 1880, p. 26.

1. In tophac. — *Eur.* : Tirol. (Alp. Trident., 2050 mt. alt.)

*calcareum* Dicks. — *Seligeria*.

*calcicola* Arn. in *Rev. bryol.* 1898, p. 5.

Syn. Ad rup. calcar. — *Eur.* : Rossia or. (Perm). *As.* : Jenissei inf.

*californicum* Sull. in *Bot. Pac. R.R. Survey (Whipple)*, p. 168, t. 6; *Lesq. et Jam. Man.* p. 237 (1884).

2. Terr. — *Am. sept.* : Calif.

*callistomum* Philib. in *Rev. bryol.* 1885, p. 76.

Syn. — *Eur.* : Norveg.

*collistomum* Dicks. — *Dicranella varia var. callistoma*.

*calobolax* C. M. in *Bot. Centralbl.* 1883, n. 2-4.

2. — *As.* : Penins. Tschutschica.

- Digitized by Google

- canescens* W. P. Sch. in *Lechler Pl. Magell.* (hb. Boissier, absq. no).  
— *Am. merid.* : Fret. Magell.
- canescens* { Hoffn. — *Rhacomitrium*.  
Vill. — *Tetraplodon mnioides* var. *Brewerianum*.
- capensi-argenteum* C. M. in *Hedwig*. 1899, p. 67.  
— *Afr.* : Prom. B. Sp.
- capillaceum* { Dicks. — *Distichium*.  
Hull. { — *Ulotia Bruchii* ?  
— *Ulotia crispa* ?  
— *Ulotia crispula* ?  
*breve, pallide et late-virens, capsulis ovatis* Dill. —  
*Weisia viridula*  
*capsulis sphaericis* Dill. — *Bartramia pomiformis*.
- capillare* L. *Sp. pl.* p. 1586, n. 30 (1753) et hb. p.p.; *Bridl. Bryol. univ.* I, p. 665 (1826); *Bryol. eur. fasc.* 6-9 (Vol. IV), *Mon.* p. 60, t. 368 et 369 excl. var.  $\gamma$  (1839); *C. M. Syn.* I, p. 281 (1849); W. P. Sch. *op. cit.* p. 449.  
*Br. foliis latiusculis congestis, capsulis longis nutantibus* Dill. *Hist. M.* p. 398, t. 50, f. 67 (1741).  
*Br. nitidum, foliis Serpylli pellucidis, capitulis reflexis et surculis annotinis et marcidis egredientibus* Dill. *Cat. Giss.* p. 227 (1718).  
*Br. squalidum* Vill. *Pl. Dauph. III*, p. 889, t. 54 (1786).  
*Br. stellare roseum pendulum minus, capsulis et setis longioribus* Dill. *Hist. M.* p. 412, t. 52, f. 78 (1741) *fid. Linpr.*  
*Hypnum capillare* *Weis Pl. crypt. fl. Gotting.* p. 205 (1770).  
*Mnium capillare* L. *Syst. veg. XIV ed.* p. 947 (1784).
2. *Ad mur., ligna, saxa terramq. per totum fere terr. orbem.*
- var. *affine* Brid. — *Br. cuspidatum*.  
var. *angustatum* Boul. — *Br. capillare* var. *flaccidum*.
- « var. *brevicuspidatum* Kindb. in *Mac. Cat.* p. 131 (1892).  
*Terr.* — *Am. sept.* : Ins. Vancouver.
- « var. *capense* C. M. *op. et l. cc.*  
— *Afr.* : Prom. B. Sp.
- « var. *carinthiacum* *Bryol. eur. et W. P. Sch. II. cc.*  
*Br. capillare* var. *compactum* C. M. *op. et l. cc.* (excl. syn. *Br. Ferchelii*).  
*Ut typ.* — *Eur.* : Suecia, alp. Carinth. *As.* : *Caucas.*
- var. *cochlearifolium* *Bryol. eur.* — *Br. elegans*.  
var. *compactum* C. M. — *Br. capillare* var. *carinthiacum*.  
var. *corsicum* Brid. — *Br. Donianum*.  
var. *cuspidatum* W. P. Sch. — *Br. capillare* var. *macrocarpum*.  
? var. *cymbifolium* H. f. et W. — *Br. cavum*.
- « var. *domingense* C. M. *op. et l. cc.*  
*Br. domingense* C. M. in *Linn.* 1843, p. 594.  
*Br. flaccidum* *Bridl. op. cit.* p. 667.  
*Br. tomentosum* *ej. op. cit.* p. 668.  
— *Am. sept.* : Haïti.
- var. *elegans* Boul. — *Br. elegans*.
- « *subsp. erythroloma* Kindb. *Eur. a. N. Am. Bryin.* p. 358.  
— *Am. sept.* : Ins. Vancouver.

- « *var. Ferchellii Bryol. eur. et W. P. Sch. II. cc.*  
*Br. elegans var. Ferchellii Bredl. Laubm. Steierm. p. 132.*  
*Br. Ferchellii Funck in Brid. Bryol. univ. p. 848 (1826).*  
*Br. Funckii var. Ferchellii Schwaegr. Sp. M. p. 69 (1830).*  
*Ad mur. et rup. in subalp. et alp. praecip. calcar. — Eur. et Am. sept.*
- « *var. flaccidum Bryol. eur. l. c.*  
*Br. capillare var. angustatum Boul. M. Est Fr. p. 396 (1872).*  
*Terr. et lignic. in loc. umbr. — Eur. et Am. sept.*  
*var. laete-virens Hüb. — Br. elegans.*
- « *var. macrocarpum Hüben. Muscol. germ. p. 442 (1831)*  
*Br. capillare*  $\beta$   $\left\{ \begin{array}{l} \text{cuspidatum W. P. Sch. Coroll. p. 74 (1856).} \\ \text{majus Bryol. eur. II. cc.} \end{array} \right.$   
*Ut et cum typ. in Eur. et Am. sept.*  
*var. majus Bryol. eur. — Br. capillare var. macrocarpum.*
- « *var. meridionale W. P. Sch. Coroll. p. 75.*  
*Br. ligusticum de Not. Epil. p. 395 (1869).*  
*Ad mur. et terr. petros. zon. merid. — Eur. et Am. sept.*  
*var. norvegicum K. et Arn. — Br. crispulum.*  
*var. obconicum Husn. — Br. obconicum.*
- « *var. platyloma (Schwaegr.) W. P. Sch. Syn. I ed. p. 371 (1860) et II ed. l. c.*  
*Br. platyloma Schwaegr. Suppl. I, P. II, p. 116, t. 76 (1816); Brid. Bryol. univ. I, p. 668 (1826).*  
*Ad mur. et rup. reg. merid. — Eur. et Afr. : Alger., ins. Azor., Canar., Madeir.*
- « *var. radiculosum Piré in Del. Fl. crypt. Belg. M. p. 170 (1883).*  
*— Eur. : Belgia.*  
*var. restitutum Kindb. — Br. restitutum.*  
*var. rhizophilum Sanio. — Br. capillare var. rosulatum.*
- « *var. rosulatum Mitt. in Ann. a. Mag. of nat. hist. 1851, p. 367.*  
*Br. capillare var. rhizophilum Sanio in sched. ; Braithw. Brit. Moossfl. II, p. 191.*  
*Br. elegans var. rosulatum Arn. in Bot. Not. 1896, p. 108.*  
*— Eur. : Anglia, Suec., Dania, Boruss. occid., Stir.*
- « *subsp. ruraliforme Kindb. op. et l. cc.*  
*— Am. sept. : Ins. Vancouver.*
- « *var. sanguineum (Arn.) Par.*  
*Br. elegans var. sanguineum Arn. in Oefv. af K. Vet.-Akad. Foerh. 1900, n. I, p. 118.*  
*— Eur. : Spitzberg.*
- « *subsp. spelungense Kindb. op. et l. cc.*  
*Stir. — Eur. : Alp. Helvet. (Splügen).*
- « *var. subelimbata (Joerg.) Par.*  
*Br. elegans var. subelimbata Joerg. in Berg. Mus. Aarsh. 1894-95, n. 13, p. 21.*  
*— Eur. : Norveg.*



- « *subsp. suecicum* Kindb. *op. et l. cc.*  
Fr. ign. — *Eur.* : Suecia.
- « *var. tectorum* Warnst. in *Verh. d. bot. Ver. d. Prov. Brandenb.*  
1899, p. 62.  
— *Eur.* : Lusat. infer.
- var. torquescens* Husn. — Br. *torquescens*.
- « *var. triste* (de Not.) Limpr. *Laubm. Deutschl.... II*, p. 377 (1892).  
Br. *triste* de Not. in *Cron. briol. ital. in Comm. II*, n. 14 (1866).  
Ad rup. granit. — *Eur.* : Ital. super. (lac. Maj.)
- var. Lesq. et Jam.* — Br. *sanguilentum*.
- capillare* Stroem. — *Ditrichum flexicaule*.
- capillareoides* C. M. in *Breut. M. capens.*  
— *Afr.* : Prom. B. Sp.
- capillifolium* C. M. — *Brachymenium capillare*.
- capillipes* C. M. in *Nuov. Giorn. Bot. ital.* 1897, p. 25.  
— *Am. merid.* : And. Boliv.
- « *var. dimorphum* C. M. in *op. et l. cc.*  
Br. *argenteum var. lanatum* W. P. Sch. in *Mandon Pl. Boliv.* n. 1664.  
Ut typ.
- « *var. subulatum* C. M. in *op. et l. cc.*  
Ut typ.
- capillisetum* C. M. in *Linn.* 1880-83, p. 382.  
2. In aren. rivul. — *Am. merid.* : Argentin. Cordob.
- capitellatum* C. M. et Kindb. in *Mac. Cat. p.* 127 (1892).  
Br. *miniaturum* Mac. *Canad. M. n.* 175 p.p.  
Ster. Ad foss. — *Am. sept.* : Ins. Vancouver.
- capitellatum* C. M. (China). — Br. *subcapitellatum*.
- capitulatum* Mitt. — *Brachymenium*.
- capitulis subrotundis reflexis, caudiculis teretibus argenteis* Dill. — Br.  
*argenteum*.
- capsulis longis subulatis* Dill. — *Barbula subulata*.
- Cardoti* Kindb. — *Webera*.
- carinatum* Boul. — *Webera cucullata var. carinata*.
- Carionis* C. M. — *Brachymenium*.
- carneum* { Drumm. — *Mniobryum albicans*.  
C. Hartm. — *Mniobryum atro-purpureum*.  
L. — *Mniobryum*.
- var. atro-purpureum* Schwaegr. — Br. *atro-purpureum*.
- var. pulchellum* { Drumm. p.p. } — *Mniobryum atro-purpureum*.  
H. f. et W. — *Webera Drummondii*.  
Hartm. — *Mniobryum atro-purpureum*.  
— *Webera pulchella*.
- var. pusillum* Schultz. — Br. *atro-purpureum*.
- catenulatum* W. P. Sch. *Syn. II ed. p.* 471 (1876).  
*Webera commutata* (fid. Card.).  
Ster. — *Eur.* : Scotia.

Catharinae C. M. in *Ule Bryoth. brasil. n. 27.*

— *Am. merid.* : Brasil. austr. tropic.

caule { *erecto, ramis lateralibus brevibus fertilibus.* L. — *Rhacomitrium microcarpum.*  
*inclinata, foliis arrectis subulatis, capitulis erectiusculis* L. — *Dicranum scoparium.*

cavifolium W. P. Sch. in *Besch. Fl. bryol. Antill. fr. p. 32* (1876).

Ster. — *Am. sept.* : Guadalup.

cavifolium { Rehm. — *Br. subcavifolium.*  
 Wils. — *Br. neodamense.*

cavum C. M. in *Bot. Zeit.* 1845, p. 94 et *Syn. I, p. 23* (1849); *Mitt. M. M. austr. am. p. 308* (1869).

*Br. capillare?* var. *cymbifolium* H. f. et W. in *Journ. de bot.* 1844, p. 145.

2. Ad mur. et terr. — *Am. merid.* : Brasil. austr. or., Monte-Video.

cellulare Hook. in *Schwaegr. Suppl. III, I, p. 214* (1827); C. M. *Syn. I, p. 291* (1849); *Mitt. M. Ind. or. p. 72* (1859).

*Brachymenium cellulare* Jaeg. *Ad. I, p. 573* (1873-74); *Par. Ind. p. 120.*  
*Bryum megalodictyon Sull. et Lesq.?*

2. Terr. — *As.* : Tonkin., Nepal., N. W. Himalaya.

Celsii L. — *Ceratodon purpureus.*

ceramiocarpum C. M. — *Anomobryum.*

cernuum { Brid. — *Br. lacustre.*  
 Bryol. eur. — *Br. pendulum.*  
 Lindb. — *Br. uliginosum.*  
 Wils. — *Br. subrotundum.*

cerviculatum Dicks. — *Dicranella.*

chalarhodon C. M. in *Dusen M. Camer. n. 207 et 660; Broth. in Engler's Bot. Jahrb.* 1897, p. 248.

*Rhodobryum chalarhodon* Par. *Suppl. Ind. p. 299* (1900).

2. Rupic. — *Afr.* : Cameroon.

chilense Reichdt. in *Novara Exped. Bot. I, p. 175, t. 30.*

Cortic. — *Am. merid.* : Chile.

chionophilum C. M. in *hb. Saharampur, n. 3738.*

— *As.* : N. W. Himalaya.

chlorosum C. M. — *Anomobryum.*

chlororhodon C. M. in *Hedwig.* 1898, p. 107.

*Rhodobryum chlororhodon* Par. *Suppl. Ind. l. o.*

2. Ad terram aren. silvat. — *Pac.* : Austral. or. (Victoria).

chryseum Mitt. *M. austr. am. p. 304* (1869).

2. — *Am. sept.* : Guatemala.

chrysoblastum C. M. in *Linn.* 1878-79, p. 477.

2. — *Am. merid.* : Venezuela.

chrysochaete C. M. in *Sikora Pl. Mad. n. 14292.*

— *Afr.* : Madagascar.

chrysoneuron C. M. in *Bot. Zeit.* 1851, p. 549; *Handb. of the N. Zeal Fl. p. 443* (1867).

*Br. duriusculum* H. f. et W. *Fl. of the N. Zeal. II, p. 84* (1855).

2. Terr. — Pac. : Alp. Austral. or., Nov. Seeland. sept. et med., Tasman.

*ciliare* { Dicks. — *Encalypta streptocarpa*.  
Gmel. — *Encalypta ciliata*.  
Grev. — *Mnium*.  
*ciliatum* Dicks. — *Hedwigia*.  
*cinclidioides* C. M. — *Mnium*.

*cirratum* Hoppe et Hsch. in *Fl.* 1819, I, p. 90; *Brid. Bryol. univ.* I, p. 672 (1826); *Bryol. eur. fasc.* 32 (Vol. IV), *Mon. Suppl.* p. 8, t. 357 (1846); *C. M. Syn.* I, p. 278 (1849) et II, p. 570 (1851); *W. P. Sch. op. cit.* p. 429.

*Br. caespitium* Schwaegr. *Sp. M.* p. 59 (1830).

*Br. intermedium* var. *aristatum* Fiedl. *Syn. Laubm. Mecklenb.* p. 69 (1844).

*Br. intermedium* β *cirratum* *Bryol. eur. fasc.* 6-9, *Mon.* p. 47 (1839).

*Br. paradoxum* *cirratum* *Jur. Laubmfl. (Esterr.-Ung.)* p. 266 (1882).

*Webera intermedia* var. *aristata* Schultz *Syll. Fl. II*, p. 179 (1828).

*Syn.* In humid. subuligin. — *Eur.* : Spitzberg, mar. Alb., Alp. Scandin., German., Stir., Helvet., Sabaud., Delphinat., etc.; *Ligur. As.* : Jenissei, Baltistan, Kashmir, Luristan, Caucas., *Minor. Am. sept.* : Groenland., Terra nov. ? ins. Miquel., Gaspes., Wyoming, Washington, Columb. brit., Colorado, M. Calif.

var. *cuspidatum* Husn. — *Br. cuspidatum*.

« var. *megalosporum* Kindb. in *Muc. Cat.* p. i22 (1892).

Ad rup. humid. — *As. sept.* : Columb. brit.

« var. *pseudo-pendulum* Warnst. in *Abh. d. bot. Ver. d. Prov. Brandenburg.* 1900, p. 203.

Terr. — *Eur.* : Boruss. or.

« var. *sulcatum* Joerg. *Om Floraen i Nord. Reisen*, p. 86 (1894).

— *Eur.* : Norveg. arctic.

<i>cirratum</i>	{	Huds. — <i>Dicranoweisia</i> .	{	<i>et stellatum tenuioribus foliis</i> Dill. — <i>Dicranoweisia</i> .	{	<i>Cynodontium polycarpum</i> ? ( <i>fid.</i> W. P. Sch.)
		<i>setis et capsulis brevioribus et pluribus</i> Dill. —		{		<i>Dicranoweisia</i> ( <i>fid.</i> C. M.)
						<i>Ptychomitrium polyphyllum</i> ( <i>fid.</i> W. P. Sch.)
		<i>setis et capsulis longioribus</i> Dill. --		{		<i>Dicranum scottianum</i> f. E.-H. ( <i>fid.</i> Lindb.)
						<i>Tortella tortuosa</i> ( <i>fid.</i> C. M. et Limpr.)

*cirriferum* (de Not.) Limpr. *Laubm. Deutschl. .... II*, p. 317 (1892).

*Pohlia cirrifer* de Not. *Mant. n.* 9 (1835) et *Syll. n.* 38 (1838).

*Syn.* Ad rivul. glacial. — *Eur.* : Simplon; alp. Helvet. ?

*clathratum* Amann in *Rev. bryol.* 1889, p. 89.

2. Ad rivul. — *Eur.* : Norveg. bor., Rhaet.

« f. *holodon* Hagen in *Tromsø Mus. Aarsheft.* 1898-99, p. 185.

— *Eur.* : Finmark.

*clavaeforme* Hpe. — *Webera*.

*clavariaeforme* C. M. — *Brachymenium*.

*clavatum* (W. P. Sch.) C. M. *Syn. I*, p. 292 (1849); *Mitt. M. austr. am.* p. 307 (1869).

*Pohlia clavata* W. P. Sch. in *Ann. sc. nat. II ser., VI*. p. 148, t. 2.

2. — *Am. merid.* : Chile.

*clavatum* H. f. et W. — *Br. curvicolium*.

*clavicaule* C. M. — *Anomobryum*.

*claviger* Kaur. in *Bot. Not.* 1884, p. 4.

*Br. veronense* (*fid. Limpr.*).

— *Eur.* : Norveg.

*Climacodontium* Par. — *Anomobryum cygnicollum*.

*coarctatum* } C. M. — *Brachymenium*.  
 } Mitt. — *Brachymenium melanothecium* ?  
 } Schultz. — *Mniobryum carneum*.

*cognatum* Mitt. *M. Ind. or.* p. 69 (1859).

*Syn.* — *As.* : Ceylan.

*coloratum* C. M. in *Linn.* 1878-79, p. 296.

*Rhodobryum coloratum* Par. *Ind.* p. 1115 (1897).

2. *Arenic.* — *Am. merid.* : Columb.

*columbicum* Kindb. — *Mniobryum*.

*comatum* Besch. *Prodr. bryol. mexic.* p. 55 (1872).

*Rhodobryum comatum* Par. *Ind.* p. 1115 (1897).

2. — *Am. sept.* : Mexic.

? *Combæ de Not. Cron. briol. ital. in Comm. II*, n. 32 (1866) et *Epil.* p. 409 (1869).

*Br. Geheebii* C. M. (*fid. de Not. l. c.*).

*Ster.* — *Eur.* : M. Sardin. centr. ; Helvet. (Aargau, quoad *Br. Geheebii.*)

*comense* W. P. Sch. *Syn. II ed.* p. 413 (1876).

*Br. caespiticius* var. *comense* Husn. *Musc. gall.* p. 212 (1889).

*Br. caespiticius* forma *Vent. et Bott. Enum. n.* 317 in *adn.* (1884).

*Er. imbricatum* Garov. *Cat. III*, n. 22 de *Not. Epil.* p. 402 (1869);  
*W. P. Sch. Syn. op. cit.* p. 418 (*fid. Philib. in Rev. bryol.* 1880, p. 36).

*Br. Philiberti* Amann in *Rev. bryol.* 1889, p. 87.

2. ♂ ign. *Muric.* — *Eur.* : Ital. super. (Como), Helvet., alp. Rhaet. ;  
*Pyren.* (*ster.*).

\* *f. brevimucronata* Bryhn in *Saertryk af Meddel. om Groenland*, 1898, p. 397.

— *Eur.* Norveg. bor. *Am. sept.* : Fret. Scoresby.

*commutatum* { Boul., Wats. — *Webera*.  
 } C. M. — *Brachymenium*.

var. *elongatum* Boul. — *Webera filum*.

β *gracile* Boul. — *Webera commutata* var. *gracilis*.

*compactile* C. M. — *Webera*.

*compactulum* C. M. in *Linn.* 1880-83, p. 383.

*Ster. Terr.* — *Am. merid.* : Argentin. Cordob., Chile.

*compactulum* C. M. (China). — *Webera subcompactula*.

*compactum* Dicks. — *Mniobryum carneum*.

- compressidens *C. M. Syn. I*, p. 490 (1849).  
*Pohlia turbinata Schwaegr. Suppl. II, P. II, II*, p. 150, t. 194 (1827).  
 2. In argill. et ad rup. madid. — *As.* : Nepal., Java, ins. Bali.
- compressulum*. — *Anomobryum*.  
*concaevum Mitt. M. austr. am.* p. 309 (1869).  
 2. — *Am. merid.* : And. Quitens.
- concinatum* Spr. — *Anomobryum*.  
*condensatum Hpe. mss.*; *C. M. in Bot. Zeit.* 1858, p. 55.  
 Syn. — *Afr.* : Pr. B. Sp.
- confertum Limpr. Laubm. Deutschl.... II*, p. 362 (1892).  
 Syn. fr. ign. — *Eur.* : Alp. Stir.
- confluens* Joerg. — Br. Kaurini.  
*congestiflorum Philib. in Rev. bryol.* 1898, p. 66.  
 1. — *As.* : M. Alatau.
- congestum Mitt. M. austr. am.* p. 309 (1869).  
 Br. caespitium *Hsch. Fl. brasil. I*, p. 44 (1840).  
 2. Terr. — *Am. merid.* : And. Quitens., Brasil.
- conicum* } *Hsch.* — *Anomobryum*.  
 } *Dz. et Mk.* — Br. Decaisnei.
- conoideum* Dicks. — *Zygodon*.  
*consimile* Mitt. — *Brachymenium*.  
*constrictum* Br. — Br. Klinggraeffii.  
*contextum* H. et H. — Br. pallescens var. *contextum*.
- contortum J. Stirton in Proced. of the Nat. hist. Soc. of Glasgow*, 1876,  
 pp. 187-188.  
 2. — *Pac.* : Nov. Seeland.
- contortum* } *Griff.* — *Brachymenium nepalense*.  
 } *Wulf.* — *Encalypta streptocarpa*.
- contractum J. O. Bomanss. in Rev. bryol.* 1899, p. 9 et in *Act. Soc. pr. Faun. et Fl. fenn.* 1899, n. 4.  
 Br. insulosum *Philib.* ?  
 Syn. Ad rup. granit. et litt. aren. mar. — *Eur.* : Ins. Aland.
- contractum C. M.* — *Webera*.  
*controversum Hoffm.* — *Weisia viridula*.
- convolutaceum C. M. in Linn.* 1875, p. 388.  
 Br. coronatum *Hpe. Prodr. Fl. Nov. Granat.* p. 50.  
 2. — *Am. merid.* : Nov. Granat.
- convolutum* Dicks. — *Harbula*.
- Corbieri Philib. in Rev. bryol.* 1887, p. 23.  
 Syn. In. turf. — *Eur.* : Normann. marit.
- coronatum Schwaegr. Suppl. I, P. II*, p. 103, t. 71 (1816); *Brid. Bryol. univ. I*, p. 650 (1826); *C. M. Syn. I*, p. 307 (1849); *Bryol. jav. I*, p. 145, t. 118 (1861-70).  
 2. Calcic. — *As.* : Nepal. ?, Ceylan, Java, ins. Banca, penins. Malay., Amboine. *Afr.* : Guinea gall., Usagara. *Am. sept.* : Florid., Cuba, Jamaic., Haïti; Guadalup. ? *Am. merid.* : And. Quitens., Guian., Brasil. or., Peruv. *Pac.* : Nov. Caledon.

var. *doliolum* Jaeg. — Br. *doliolum*.

var. *minus* Hpe. — Br. *pachytheca*.

« var. *orientale* Hpe. in *Nuov. Giorn. bot. ital.* 1872, p. 281.

— As. : Borneo.

var. Wils. — Br. *nitens*.

*coronatum* { Besch. — Br. *Hogbergii*.  
C. M. p.p. — Br. *Phallus*.  
Hpe. — Br. *convolutaceum*.  
Hpe. et Ltz. — Br. *gracilifolium*.  
H. f. et W. — Br. *curtum*.  
Wils. — Br. *barbulaceum*.

*corrugatum* Hpe. *Symb.* 1870, p. 276.

Br. *argenteum* var. *corrugatum* Besch. *Prodr. bryol. mexic.* p. 53 (1871).

Br. *argenteum* β *lanatum* Hsch. *Fl. brasil.* I, p.

2. Rupic. et terr. — As. : M. Neilgherr. *Afr.* : Ins. Mayotte. *Am. sept.*  
et *merid.* : Tropic.

« var. *niveum* Hpe. *Symb.* 1872, p. 52.

— *Am. merid.* : Brasil. austr. or.

*corsicum* Kindb. *Eur. and N. Am. Bryin.* p. 357 (1897).

Ster. — *Eur.* : Corsica.

*corsicum* W. Arn. — Br. *Donianum*.

*crassicaule* Dusen *M. Am. austr.* n. 482.

Ad rup. irror. — *Am. merid.* : Patag. occid.

*crassicostatum* C. M. — Webera.

*crassidens* Kindb. — Webera.

*crassimucronatum* Philib. in *Rev. bryol.* 1899, p. 31.

— As. : M. Alatau.

*crassinerve* Hook. f. et Wils. *Fl. of the N. Zeal.* p. 83 (1855).

— *Pac.* : Austral. or. (Victoria), Nov. Seeland.,

Tasman.

*crassinervium* Ltz. in *Bot. Zeit.* 1866, p. 186.

2. Rupic. — *Am. merid.* : Chile austr.

*crassinervium* Besch. — Br. *Pancheri*.

*crassirameum* Ren. et Card. in *Bot. Gaz.* 1890, p. 57, t. 8 B.

2. — *Am. sept.* : Oregon, Calif.

« var. *Covillei* eor. in op. cit. 1900, p. 21.

— *Am. sept.* : M. Rupestr.

*crassum* Hook. f. et Wils. in *Fl. of the N. Zeal.* II, p. 86, t. 86,

f. I (1855) et *Handb.* p. 442 (1867).

2. — *Pac.* : Austral. or., Nov. Seeland., Tasman.

*creberrimum* Tayl. in *Lond. Journ. of bot.* 1846, p. 54 ; C. M. *Syn.* I,  
p. 283 (1849) ; *Handb. of the N. Zeal. Fl.* p. 442 (1867).

2. Ad terr. humid. — *Pac.* : Austral. or., Nov. Seeland., Tasman.

*cribrosum* Hoffm. — *Coscinodon pulvinatus*.

*crinitum* Mitt. — *Brachyemium systylium*.

*crispatum* Dicks. — *Rhabdoweisia denticulata*.

*crispulum* Hpe. mss. ; C. M. in *D. II Deutschl. Nordpolarf.* II, 1873 (?)  
p. 69 (false ut syn. Br. *teretis*) ; Hagen in *Tromsøe Mus. Aarsheft.*  
1898-99, p. 205.

Br. *capillare* var. *norvegicum* Kaur. et Arn. mss. ; Arn. in *Bot. Not.* 1896,  
p. 107.

- Br. neodamense Auct. *scand. p. p. (nec Itzig.)*.  
 Br. pseudo-triquetrum { *var. cavifolium* } *Berggr. M. et Hep. Spetsb. p. 63*  
                                   { *var. compactum* } (1875).  
                                   { *var. rotundifolium* *Zeit. in K. Sv. Vet.-Akad.*  
                                   *Handl. 1876, n. 13, p. II?*
2. ♂ ign. In argillos. — *Eur.* : Spitzberg, Norveg. bor. *Am. sept.* :  
 Groenland, occid. et or.
- crispum* { (L.) Gmel. — Ulota.  
                   { Schreb. — Dicranella Schreberi.
- cristatum* *Philib. in Rev. bryol. 1897, p. 23.*  
 Syn. Terr. — *Eur.* : Alp. Delphinat.
- cristatum* C. M. — Brachymenium.  
*crudoides* Sull. et Lesq. — Webera.
- crudum* { Mitt. } (M. austr. am) — Webera synoico-cruda?  
                   { (1876). — Webera austro-cruda.  
                   { Schreb. — Webera.
- Crügeri *Hpe. in C. M. Syn. I, p. 300 (1849) et II, p. 574 (1851); Mitt.*  
*M. austr. am. p. 298 (1869).*  
 Br. Lansbergii *Dz. et Mk. Prodr. fl. surinam. p. 40, t. 4 (1854).*  
 Br. ovalifolium *Sull. in Proceed. of the Amer. Acad. 1861.*
2. Ad rup. et terr. — *Am. sept.* : Cuba, Jamaic., S. Vincent. *Am.*  
*merid.* : Ins. Trinitat., Venezuela, Guian. batav., Amazon., And.  
 Quitens. et Peruv.
- cubitule* Dicks. — Br. pseudo-triquetrum.  
*cucullatiforme* Kindb. — Webera.  
*cucullatum* Schwaegr. — Webera.
- Culmannii *Limpr. Laubm. Deutschl. .... II, p. 357 (1892).*  
 Syn. — *Eur.* : Norveg., Helvet. (Gemmi). *Am. sept.* :  
 Fret. Scoresby.
- cuneifolium* { Dicks. — Tortula.  
                   { Hoffm. — Tortula muralis *var. aestiva.*
- cupulatum C. M. in *Linn. 1871-73, p. 149.*  
 2 ? — *Pac.* : Austral. or.
- curtum *Par. Ind. p. 176 (1894).*  
 Br. brevicaulis *Hpe. in Linn. 1869-70, p. 528.*  
 Br. coronatum *H. f. et W. Fl. of the N. Zeal. II, p. (1855).*  
 Br. iteratum *Par. op. cit. p. 189.*  
 Br. macropelma C. M. in *Linn. 1872, p. 149.*  
 1 (Hpe.), 2 (C. M.) — *Pac.* : Austral.
- curvatum *Kaur. et Arn. in Bot. Not. 1897, p. 67 et in Rev. bryol. 1898,*  
*p. 3. (subsp. Br. Hagenii).*  
 Syn. Ad terr. arenac. — *Eur.* : Lappon. lulens., Norveg. bor.
- curvicollum *Mitt. in Handb. of the N. Zeal. Fl. p. 442 (1867).*  
 Br. clavatum *H. f. et W. in Fl. of the N. Zeal. II, p. 84, t. 85, f. 2 (1855).*
2. Terr. — *Afr.* : Tristao da Cunha. *Pac.* : Nov. Seeland., Tasman.
- « *var. extenuatum* *Mitt. op. et l. cc.*  
                                   — *Pac.* : Nov. Seeland.
- « *var. γ* *Mitt. op. et l. cc.*  
                   Ubi praeced.
- curvisetum Garov. — Webera polymorpha *var. brachycarpa.*

*cuspidatum* W. P. Sch. *Syn. II* ed. p. 430 (1876).

Br. affine *Lindb. M. scand.* p. 16 (1879).

Br. binum *var. cuspidatum* *Bryol. eur. fasc. 6-9 (Vol. IV), Mon.* p. 50 (1839).

Br. capillare γ affine *Brid. Bryol. univ. I*, p. 848 (1826).

Br. cirratum *var. cuspidatum* *Husn. Musc. gall.* p. 248 (1889).

Br. hybridum *Br. mss.; Bryol. eur. l. c. in adn.*

Br. paradoxum (*non Schwaegr.*) *Jur. Laubmfl. Oesterr.-Ung.* p. 266 *excl. var. β* (1882).

*Pohlia paradoxa* *Hüb. Musc. germ.* p. 479 (1833).

*Webera affinis* *Br. mss. in Brid. op. et l. cc.*

*Syn. Ad mur. et rup. humidiusc. — Eur. : Zon. sept. et intermed. As. : Japon., Baltistan, Kashmir, Caucas. Am. sept. : Ins. Vancouver, Washington, Oregon, Montana, Idaho, Wisconsin, Illinois.*

« *var. cylindricum* (*Arn.*) *Par. Suppl. Ind.* p. 61 (1900).

Br. affine *var. cylindricum* *Arn. in Bot. Not.* 1897, p. c.ic.

— *Eur. : Scandin.*

« *var. obtusiusculum* (*Lindb. et Arn.*) *Par. Ind.* p. 177 (1894).

Br. affine *var. obtusiusculum* *L. et A. M. As. bor. II*, 1890, p. 43.

*Ut typ. — As. : Jenissei infer.*

« *var. urnigerum* (*Arn.*) *Par. Suppl. Ind. l. c.*

Br. affine *var. urnigerum* *Arn. in op. et l. cc.*

— *Eur. : Suecia.*

*cuspidatum* { *Drumm., Turn. — Mnium.*  
Gmel. — *Pottia lanceolata.*  
Schreb. — *Mnium.*  
Wils. p.p. { — *Br. Thomsoni.*  
— *Br. turbinatum.*

*cyclophyllum* (*Schwaegr.*) *Bryol. eur. fasc. 6-9 (Vol. IV), Mon.* p. 63, t. 339 (1839); *C. M. Syn. I*, p. 287 (1849) *et II*, p. 571 (1851); *W. P. Sch. op. cit.* p. 455.

? *Br. obtusifolium* *Turn. Musc. hib.* p. 116, t. II, f. I (1804); *Brid. Bryol. univ. I*, p. 639 ? *ejusd. hb ! (fid. C. M.)*.

*Br. tortifolium* *Funch mss.; Brid. op. cit.* p. 844 (1826).

*Mnium cyclophyllum* *Schwaegr. Suppl. II. P. II*, p. 160, t. 194 (1827).

2. *Ad terr. palud. stagn. exsicc. et rup. madid. — Eur. : Suec., Dania merid., Boruss. occid., Siles., Belg. sept., Voges., Silva nigr., Jurass., Francon., Bavar. As. : Jenissei infer. Am. sept. : Fret. Smith, Groenland., Niagara, M. Rupestr., Pensilv.*

? *var. laxifolium* { *Sanio. — Webera Ludwigi.*  
*var. lyceense* }

*cygnicollum* *C. M. in Bot. Zeit.* 1858, p. 156.

*Syn. — Am. merid. : Summ. And. Chilens.*

*cygnicollum* *C. M.* (1848). — *Anomobryum.*

*cygnopelma* *C. M. in Bull. hb. Boissier*, 1897, p. 550.

2. — *Am. sept. : Jamaic.*

*cylindricum* { *Dicks. — Webera elongata (fid. C. M.)*  
— *Webera longicolla (fid. Limpr.)*  
*Mont. — Webera cylindrica.*  
*C. M. — Anomobryum soquense.*  
*cymbifolium* { *Lindb. — Anomobryum.*



- cymbuliforme* Card. — *Webera cucullata* var. *carinata*.
- dalachanicum* Jur. et Milde Beitr. ad fl. bryol. Or. p. 596 (1870).  
2. — As. : Kurdistan, Persia.
- Daviesii* Dicks. — *Glyphomitrium*.
- dealbatum* Dicks., Drum. — *Amblyodon*.
- Decaisnei* Dz. et Mk. M. Archip. ind. p. 19, t. 9 (1844); C. M. Syn. I, p. 251 (1849).  
Br. conicum Dz. et Mk. in Ann. sc. nat. 1844, p. 300.  
*Rhodobryum Decaisnei* Par. Ind. p. 1116 (1897).  
2. Lignic. — As. : M. Java.
- decipiens* Kaur. ap. Joergens. in Christiania Vid. Selsh. Foerh. 1894, p. 78 (descr. *paupera*, fid. Hagen).  
— Eur. : Norveg.
- decipiens* Brid. — *Webera annotina*.
- decolorifolium* C. M. in Nuov. Giorn. bot. ital. 1897, p. 249.  
— As. : China centr. (prov. Schen-Si).
- decurrens* C. M. in Bot. Zeit. 1855, p. 751.  
Ster. — Afr. : Prom. B. Sp.
- decurrentinervium* C. M. in Hedwig. 1897, p. 95.  
Ster. — Am. merid. : Argentin. patag.
- decursivum* C. M. in Hedwig. 1898, p. 224.  
Ster. — Am. sept. : M. Porto-Ricens.
- decursivum* C. M. (Afr. austr.) — *Br. pseudo-decursivum*.
- delicatulum* { Hedw. — *Mniobryum carneum*.  
                  { Wils. — *Webera*.
- delphinense* Corbière in Rev. bryol. 1899, p. 83.  
Br. pallescens ? (fid. Hagen).  
Heteroic. Ad rup. silic. juxta mol. glac. — Eur. : Alp. Delphin.
- Demawendii* C. M. — *Webera*.
- demissum* Hook. — *Plagiobryum*.
- dendroides* { L. f. — *Mniodendron comosum*.  
                  { *polycephalum*, *Phyllitidis folio undulato pellucido*, *thecis ovatis pendulis* Dill. — *Mnium undulatum*.
- densifolium* Mitt. — *Br. schistoneuron* var. *densifolium*.
- denticulatum* Kindb. in Bull. Torr. bot. Club, XVII, p. 273 (1890).  
2. — Am. sept. : Alberta, M. Selkirk.
- dentiferum* Hpe. Symb. 1877, p. 726.  
Syn. — Am. merid. : Brasil. austr. or.
- depressum* C. M. in Dusen M. Camer. n. 663; Broth. in Engler's Bot. Jahrb. 1897, p. 245.  
2. fr. ign. Ad rup. inund. — Afr. : Cameroon.
- diaphanum* C. M. in Hedwig. 1897, p. 94.  
— Am. merid. : La Plata.
- diaphanum* { Gmel. — *Orthotrichum*.  
                  { Web. — *Tetraphis pellucida*.

*dichotomum* Hedw. *Sp. M.* p. 163, t. 42 (1801); *Brid. Bryol. univ. I.*, p. 646 (1826); *C. M. Syn. I.*, p. 304 (1849); *Mitt. M. austr. am.* p. 301 (1869).

Br. bulbillosum Mont. (fid. Mitt.).

Mnium dichotomum P. B. *Prodr.* p. 74 (1805).

2. Ad terr. et saxa. — *Am. sept.* : Ins. Bermud. *Am. merid.* : Columb., And. Quitens., Monte-Video.

*dichotomum* H. f. et W. — Br. atro-purpureum.

*dicranoides* { Hsch. — Brachymenium.

{ W. P. Sch. — Brachymenium commutatum.

*Didymodon* { Gmel. — Barbula rigida.

{ Hoffm. — Ditrichum tortile var. pusillum.

*didymodontium* Mitt. — Brachymenium.

*dilatato-marginatum* C. M. in Hedwig. 1898, p. 102.

Rhodobryum dilatato-marginatum Par. *Suppl. Ind.* p. 299 (1900).

2. — Pac. : Austral. or. (Nov. Vales).

*dilatatum* Joerg. — Br. eubryoides.

*dimorphum* (C. M.) Broth. *Natür. Pflanzenfam.* p. 613.

Mnium dimorphum C. M. in Linn. 1878-79, p. 467.

2. — Am. merid. : Venezuela.

*dimorphum* C. M. — Br. afro-dimorphum.

*dissolutinerve* C. M. in Linn. 1880-83, p. 376.

Ster. Terr. — Am. merid. : Uruguay.

*distans* Wils. — Br. turbinatum.

*Dobsonianum* C. M. in Hedwig. 1898, p. 108.

Rhodobryum Dobsonianum Par. *op. et l. cc.*

Ster. — Pac. : Tasman.

*doliolum* Duby in Moritz Syst. *Verz.* p. 133; *C. M. Syn. I.*, p. 304 (1849).

Br. coronatum var. doliolum Jaeg. *Ad. I.*, p. 617 (1873-74).

1. Terr. — As. : Java, ins. Andaman, Aroë, Malacca, Celebes.

*dolomiticum* Kaurin in Hagen M. *Norveg. bor.* p. 223 (1898-99).

Heteroic. Ad rup. dolomit. littor. — Eur. : Norveg. bor.

*domingense* { C. M. — Br. capillare var. domingense.

{ Steud. — Rhodobryum Antillarum.

*Donianum* Grev. in *Trans. of the Linn. Soc.* 1828, p. 345, t. 3, f. 6; W.

P. Sch. *Coroll.* p. 75 (1856) et *op. cit.* p. 454.

Br. capillare  $\beta$  conicum *Brid. Bryol. univ. I.*, p. 849 (1826).

Br. corsicum Walk. *Arn. mss.*

Br. insulare de Not. *Spic.* p. 7 (1837) et *Syll.* p. 128 (1838).

Br. Mülleri Spr. M. *Pyren. n.* 138 et in *Ann. and Mag. of nat. hist.* 1849,

Br. platyloma (non Schwaegr.) *Bryol. eur. fasc.* 6-9 (Vol. IV), *Mon.* p. 58,

t. 366 (1839); *C. M. Syn. I.*, p. 282 (1849) et II, p. 571 (1851).

2. Ad rup. et terr. glareos. — Eur. : Angl., Belg., Galloprov., Corsica, Sardin., Itel., Dalmat., ins. Ionien., Melit., Balear.; Algarb.

Afr. : Alger., Canar., Madeir. *Am. sept.* : Ins. Vancouver.

♂ *subsp. humile* Kindb. in *Rev. bryol.* 1898, p. 91.

— Eur. : Lusit.

*Donianum* Kindb. — Br. heteroneuron.

dovrense W. P. Sch. *Syn. II ed. p.* 413 (1876).

Syn. Ad terr. humid. et turfosa. — *Eur.* : Alp. Norveg. (Dovre).

*Drummondii* C. M. — Webera.

dubiosum *Par. Suppl. Ind. p.* 62 (1900).

Br. zonatiforme *Kindb. Eur. and N. Am. Bryin. p.* 371 (1897).

Fl. et fr. ign. Rupic. — *Eur.* : Alp. Norveg. (Dovre).

duplicatum *Broth. in Oefv. af K. Vet.-Akad. Handl.* 1900.

*Rhodobryum duplicatum Par. op. et l. cc.*

2. fr. ign. Lignic. et ad terr. silvat. — *Am. merid.* : Brasil. centr. (Matto-Grosso).

Duriei W. P. Sch. *mss.* ; *Besch. Cat. M. Alger. p.* 24 (1882).

2. Terr. — *Afr.* : Alger.

duriusculum W. P. Sch. in *Besch. Fl. bryol. Nouv. Caled. p.* 214 (1873).

2. — *Pac.* : Nov. Caled.

*duriusculum* H. f. et W. — Br. chrysoneuron.

*Dusenii* C. M. — Brachymenium.

Duvallii *Voit in Sturm Deutsch. Fl. II, Heft 12 c. ic.* (1812); *Brid. Bryol. univ. I, p.* 679 (1826); *Bryol. eur. fasc.* 6-9 (*Vol. IV*), *Mon. p.* 64, *t.* 371 (1839); *C. M. Syn. I, p.* 260 (1849); *W. P. Sch. op. cit. p.* 458.

Br. Sphagni *Brid. op. cit. p.* 852.

Br. ventricosum var. *Schwaegr. Sp. M. p.* 54 (1830).

Br. Weigelii *Spreng. Mant.-prim. Add. p.* 55, *n.* 87 (1807).

*Mnium Duvallii Schwaegr. Suppl. I, P. II, p.* 126, *t.* 79 (1816).

2. Ad scatur. et in prat. uligin. — *Eur.* : Norveg., Scotia, Fennia, German., Bohem., Tatra, Stir., Bavar., Voges., Westphal., Belg., Gall. centr. (Arvern.) *As.* : Jenissei super., Obi, Kashmir, Caucas. *Am. sept.* : Groenland., fret. Smith, Labrador, Terra nov., ins. Miquel., Nov. Angl., Canada merid., Alaska, M.-Rupestr. et Selkirk, Washington, Oregon, Calif., Nov. Mexic.

« var. gaspeanum *Kindb. in Mac. Cat. p.* 272 (1892).

Br. gaspeanum *ej. Eur. and N. Amer. Bryin. p.* 352 (1897).

Ad rup. humid. — *Am. sept.* : Gaspes.

« var. lato-decurens *C. M. et Kindb. in Mac. Cat. p.* 132.

Br. lato-decurrens *Mac. Canad. M. n.* 487.

Ad terr. humid. — *Am. sept.* : M. Selkirk, Columb. brit.

*duvalioides* Itz. — Br. ps. triquetrum var. *duvalioides*.

*Eatoni Mitt. in Journ. of the Linn. Soc. 1877 et in Phil. Trans. of the roy. Soc. 1877, p.* 35, *t.* III, *f.* 4.

Br. Orthotheciella *C. M. in Engler's Bot. Jahrb. 1883, p.* 78 et in *Forschungsreise..... Bot. p.* 15, (1889); *Par. Ind. p.* 200 (1894).

Syn. — *Afr.* : Ins. Kerguelen.

*Ecklonianum* C. M. — Webera.

*Edwardsianum* C. M. et *Kindb.* — Br. Riddlecomiae.

*Ehrenbergianum* *C. M. Syn. I, p.* 255 (1849).

*Rhodobryum Ehrenbergianum Par. Ind. p.* 1116 (1897).

2. — *Am. sept.* : Mexic.

*elegans* Nees v. Esenb. mss. in *Brid. Bryol. univ.* I, p. 846 (1826); W. P. Sch. op. cit. p. 452.

Br. capillare { *var. cochlearifolium* Brid. op. cit. p. 666; *Bryol. eur.* fasc. 6-9 (Vol. IV), *Mon.* p. 61, t. 367 f. n.; C. M. Syn. I, p. 281 p.p. (1849).  
*var. elegans* Boul. M. Fr. I, p. 264 (1884).  
*var. laete-virens* Hüb. *Musc. germ.* p. 442 (1833).

2. In rup. fiss. humid. — *Eur.* : Spitzberg., Westphal., Riesengeb., Tatra, Austr. inf., Salisb., Stir., Carinth., alp. Algov., Bavar., Tirol., Rhaet., Helvet., Sabaud., Jurass. *Am. sept.* : Fret. Scoresby, ins. Miquelon, Gaspes.

*var. carinthiacum* Breidl. — Br. capillare *var. carinthiacum*.

*var. Ferchelii* ej. — Br. capillare *var. Ferchelii*.

*var. sanguineum* Arn. — Br. capillare *var. sanguineum*.

*var. subelimbata* Joerg. — Br. capillare *var. subelimbata*.

*elegans* { Grev. — Br. pallens *var. speciosum*.  
 { Vent. et Bott. p.p. — Br. Venturii.

*elegantulum* Ltz. *Moosst.* p. 157 (1864).

2. — *Am. merid.* : Chile austr.

*ellipsifolium* C. M. in *Fl.* 1890, p. 447.

Ster. — *Afr.* : Kilima N'Djaro.

*ellipticum* { Brid. — Mnium punctatum.  
 { Mitt. — Brachymenium.

*elongatum* { Dicks. — Webera.  
 { Drumm. — Webera cruda.  
 { Dz. et Mk. — Webera leptocarpa.  
 { Hoffm. — Rhacomitrium canescens *var. ericoides*.  
 { Mitt. (1876). — Webera austro-elongata.  
 { Stroem. — Trematodon ambiguus.

*var. alpinum* Bryol. eur. — Webera longicolla.

*var. longicollum* H. et T. — Webera longicolla.

*var. macrocarpum* Bryol. eur. — Webera elongata *var. macrocarpa*.

*elwendicum* Fehln. in *Verh. d. K. zool. bot. Ges. in Wien*, 1883, p. 435 et in *Hedwig*, 1887, p. 116.

Syn. Rupic. — *As.* : M. Elwend.

*emergens* C. M. — Miehllichhoferia.

*encalyptaceum* C. M. in *Linn.* 1880-83, p. 366.

Ster. — *Am. merid.* : Uruguay.

*erectis* { *angustifolium, caule reclinato* Dill. — Dicranum scoparium.  
 { *brevibus, foliis reflexis* Dill. — Dicranella squarrosa.  
*capitulis* { *calyptra laxa conica, foliis Serpilli pellucidis, angustioribus* Dill. — Encalypta vulgaris.  
 { *trichodes, foliis latiusculis congestis in pilum canescentem desinentibus* ej. — Tortula muralis.  
 { *falcatis capitulis, trichodes, foliis latiusculis extantibus, in pilum canescentem desinentibus* ej. — Tortula ruralis.  
 { *et obtusis capitulis, calyptra laxa conica, foliis Serpilli pellucidis, latioribus* Dill. — Encalypta ciliata.  
*longis* { *acutis et falcatis capitulis, calyptra subfusca, foliis Serpilli pellucidis* ej. — Tortula subulata.



- Mnium erythrocaulon* *Schwaegr. Suppl. I, P. II, p. 127, t. 80* (1816).  
*Rhodobryum erythrocaulon* *Par. Ind. l.c.*  
 Ster. — *Afr.* : Usambara, Madagasc., ins. Franciae, Rodriguez.  
*erythroneuron* *Mitt. M. austr. am. p. 312* (1869).  
 2. — *Am. merid.* : And. Nov. Granat. et Quitens.  
*erythrophylloides* *Kindb. in Mac. Cat. p. 272* (1892).  
 Ster. — *Am. sept.* : Alberta.  
*erythrophyllum* *Kindb. in op. cit. p. 131*.  
 Ster. Paludic. — *Am. sept.* : Nov. Scot., Ontario, Columb. brit., fret. Behring.  
*erytostegum* *C. M. in Fl. 1886, p. 279*.  
 2. Cortic. — *Afr.* : Ins. Sao Thoma.  
*erythrotropis* *C. M. in Linn. 1875, p. 383*.  
 2. fr. ign. Cortic. — *Afr.* : Niam-Niam.  
*eubryoides* *Limpr. mss.*  
*Br. dilatatum* *Jorgens. in Christiania Vid. Selsk. Foerh. 1894, p. 102*.  
 — *Eur.* : Norveg.  
*euryloma* *Card. et Ther. in Bot. Gaz. 1900, p. 20, t. V*.  
 2. — *Am. sept.* : Puget Sound.  
*eurystomum* *Ren. et Card. in Bull. Soc. roy. bot. Belg. 1891, II, p. 188*;  
*Ren. M. Masc. Mad. n. 70*.  
 2. Muric. — *Afr.* : Ins. Borbon.  
*evadiinerve* *Broth. in Act. Soc. sc. Fenn. 1898, n. 2, p. 24*.  
 2. fr. ign. Rupic. — *As.* : Kasbmir.  
*exaltatum* *Spr.* — *Acidodontium*.  
*excurrens* *Lindb. in Bot. Not. 1886, p. 93*.  
*Br. versicolor* *Kaur. in op. cit. 1885, p. 161*.  
 2. In aren. humid. — *Eur.* : Norveg.  
*exiguum* { *creberrimis capsulis* { var. *minor* f. A-E. — *Pottia truncatula*.  
                                   *rufis* Dill. { var. *major* f. F-K. — *Pottia intermedia*.  
                                   *erectis parvis subrotundis creberrimis capitulis rufis, foliolis*  
                                   *Serpilli angustis pellucidis* ej. — *Pottia truncatula*.  
*exile* *Dz. et Mk.* — *Brachymenium*.  
*eximium* *Mitt. in Handb. of the N. Zeal. Fl. p. 440* (1867).  
 Ster. — *Pac.* : Nov. Seeland. sept. et med.  
*exstans* *Mitt. M. Ind. or. p. 72* (1859).  
 Syn. — *As.* : Tibet. occid.  
*extinctorium* { L. p.p. — *Encalypta vulgaris*.  
                           L.  $\beta$  -- *Encalypta ciliata*.  
*extenuatum* *Ren. et Card. in Bot. Gaz. 1890, p. 57, t. 8 A*.  
 2. — *Am. sept.* : Oregon.  
*extenuatum* *Mitt.* — *Brachymenium*.  
*Fabronia* *C. M. in Linn. 1880-83, p. 387*.  
 Ster. Terr. — *Am. merid.* : Argentin. temper.  
*fabronioides* *C. M. Brachymenium*.  
*fabroniopsis* *C. M.* — *Anomobryum*.  
*Falconeri* *C. M.* — *Epipterygium*.

*fallax* Milde in Rab. *Bryoth. eur. n.* 242 c. *diagn.* (1859); W. P. Sch. *op. cit.* p. 426.

Br. lacustre var. alpinum *Bryol. eur. fasc.* 6-9 (Vol. IV), *Mon.* p. 16, t. 332  $\beta$ .

Br. pallens  $\beta$  fallax Jur. *Laubmfl. Oesterr. Ung.* p. 287 (1882).

2. Loc. arenos. et ad rivul. — *Eur.* : Norveg., ins. Aland, Dania, Britann., Westphal., Siles., Salisb., Carinth., alp. Bavar., Helvet., Gall. centr. et merid., Pyren., Astur. *Am. sept.* : Fret. Behring.

$\alpha$  var. baldense (Vent.) *Limpr. Laubm. Deutschl.*.... II, p. 331 (1892).

Br. baldense Vent. *mss.*; de Not. *Epil.* p. 381 (1869).

Br. pallens var. baldense Vent. et *Bot. Enum. critt.* p. 34.

In sabul. humid. — *Eur.* : Tirol., Helvet.

*fallax* Dicks. — Barbula.

*fasciculare* } Dicks. — Funaria.

                  } Hoffm. — Rhacomitrium.

*fasciculatum* Dicks. } — Eucladium verticillatum.

                          } — Hymenostylium curvirostre.

Fendleri C. M. in Linn. 1878-79, p. 475.

2. — *Am. merid.* : Venezuela.

Ferchellii Funck. — Br. capillare var. Ferchellii.

ferriviae C. M. in Ule *Bryoth. brasil. n.* 139.

Terr. — *Am. merid.* : Brasil. austr. or.

ferrugineum Jungh. — Mniodendron divaricatum.

                  } Dicks., Par. — Anomobryum.

*filiforme* } Mitt. — Anomobryum cymbifolium.

                  } Vill. — Blindia acuta.

*filinerve* Mitt. — Anomobryum.

*filum* W. P. Sch. — Webera.

finmarchicum Kaur. ap. Joergens. in *Christiania Vid. Selsk. Foerh.* 1894, p. 88.

Br. lutescens Bomanss. in *Rev. bryol.* 1897, p. 1; *Par. Suppl. ind.* p. 67 (1900).

Br. pallens var. brevisetum Lindb. et Arn. *M. As. bor.* II, p. 36 (1890).

Br. Winkelmanni Ruthe in Hedwig. 1897, p. 283.

— *Eur.* : Norveg. bor. *As.* : Jenissei infer.

fixum Ruthe in Hedwig. 1897, p. 386.

— *Eur.* : Pomer., ins. Aland.

flaccidisetum C. M. — Brachymenium.

*flaccidum* } Brid. — Br. capillare var. domingense.

                  } H. f. et W. — Br. caespiticium.

flaccum Wils. in *Kew Journ. bot.* IX, p. 332 (1857); *Mitt. M. Ind. or.* p. 72 (1859).

2? — *As.* : Sikkim.

flagellaceum Warnst. in *Dorpat. Naturf. Ges.* 1894, p. 426.

Syn. In calcar. — *Eur.* : Livonia.

*flagellare* } Brid. — Br. warneum.

                  } Hoffm. — Dicranum.

                  } Kaur. — Br. lapponicum.

flagellicoma C. M. in Linn. 1880-83, p. 369.

Ster. — *Am. merid.* : Argentin. subtropic.

flagellosum Kindb. *Eur. and N. Am. Bryin.* p. 356 (1897).

— *Am. sept.* : M. Rupestr.

*flavescens* Kindb. Enum.; Bryhn in Christiania Vid. Sellsk. Foerh. 1888, n. 6.

B. arctico  $\times$  pallens Philib. in Rev. bryol. 1889, p. 62.

2. Rupic. — Eur. : Norveg.

*flavescens* Dicks. { — *Barbula fallax*,  
(fid. Wils. in W. P. Sch. Syn.). — *Dichodontium*  
pellucidum var. serratum.

*flavifolium* C. M. in Hedwig. 1896, p. 101.

*Rhodobryum flavifolium* Par. Suppl. Ind. p. 299 (1900).

2. — Pac. : Austral. or. (Victoria).

*Fleischeri* Warnst. in Fl. u. W. Bryoth. Eur. merid. n. 56.

Ster. Ad rup. calcar. cataract. — Eur. : Neapol. (ins. Lizi).

*flexicaule* C. M. in Nuov. Giorn. bot. ital. 1896, p. 96.

Ster. Terr. — As. : China centr. (prov. Schen-Si).

*flexifolium* { C. M. — *Brachymenium*,  
Dicks. — *Didymodon*.

*flexisetum* Mitt. M. austr. am. p. 295 (1869).

1 vel syn. Paludic. — Am. merid. : And. Quitens.

*flexisetum* Arn. — Br. Arnellii.

*flexuosum* Aust. in Coult. Bot. Gaz. IV, p. 152; Lesq. et Jam. Man. p. 227 (1884).

2? Arenic. — Am. sept. : Columb. brit.

*flexuosum* { C. M. — *Webera*,  
L. p.p. { — *Campylopus*,  
— *Campylopus torfaceus*.

*floresianum* C. M. — *Acidodontium*.

*floridanum* Ren. et Card. in Rev. bryol. 1893, p. 4.

— Am. sept. : Florida.

*fluminale* C. M. in Dusen M. Camer. n. 666; Broth. in Engler's Bot. Jahrb. 1897, p. 248.

*Rhodobryum fluminale* Par. Ind. p. 1116 (1897).

2. Ad saxa rivul. — Afr. : Cameroon.

*foliis* { *capillaribus fasciculatis, seta bilineari, capsulis sphaeroideis,*  
*operculo conico arcuato* Stroem. — *Cynodontium schisti* (p.p. ?)  
*lanceolatis falcatis, setis alaribus, capsula ovata inclinata* Hall.  
— *Trimmia megapolitana*,  
*latiusculis congestis, capsulis longis nutantibus* Dill. — Br. capil-  
lare.  
*mollibus subulatis, setis brevissimis alaribus, capsulis ovatis*  
Hall. — *Bartramia Halleriana*,  
*setaceis curvatis, capsulis erectis, obtuse ovatis, capitello oblique*  
*rostrato, apophysi capitulo subjecta* Oeder. — *Dicranum Blyttii*,  
*striatis acuminatis reflexis capsulis sphaericis subnutantibus,*  
*maturitate striatis* Oeder. — *Bartramia Oederi*.

*fontanum* Sw. — *Philonotis*.

*formosum* Mitt. mss.; E. S. Salmon in Journ. of bot. 1900, p. 329, t. 413.

2? — As. : M. Neilgherr.

*Forsteri* Dicks. — *Zygodon*.



foveolatum *Hugen in Saertryk af Meddel. om Groenland, 1898, p. 402 (nom.) et in Nyt. Mag. for Naturvid. 1900, p. 333.*

Syn. — *Eur.* : Norveg. bor. *Am. sept.* : Fret. Scoresby.

*fragile* } C. M. — Brachymenium.  
Dicks. — Campylopus.

Friederici Mülleri *Ruthe in Hedwig. 1897, Beibl. p. 119.*

1. In gramin. aren. — *Eur.* : Ins. Horkum.

Fridtzii *Hagen in Tromsø Mus. Aarsheft. 1898-99, p. 217.*

1. — *Eur.* : Norveg. bor.

Froudei *Kindb. in Ott. Nat. V, p. 180.*

Syn. — *Am. sept.* : Fret. Behring.

fulvellum *Wils. in Kew Journ. bot. IX, 330 (1857); Mitt. M. Ind. p. 69 (1859).*

2. — *As.* : Sikkim.

*fulvellum* } Dicks. — Dicranum.  
Wils. p.p. — Br. lacustre.

*funarioides* Brid. — Funaria dentata.

Funckii *Schwaegr. Suppl. I, P. II, p. 89, t. 69 (1816); Brid. Bryol. univ. I, p. 660 (1826); Bryol. eur. fasc. 6-9 (Vol. IV), Mon. p. 77, t. 382 (1839); C. M. Syn. I, p. 305 (1849); W. P. Sch. op. cit. p. 446.*

Br. articulatum *Sendtn. in sched. ; C. M. Syn. I, p. 305 (1849).*

Br. Geheebii *C. M. (fd. Milde in Hedwig. 1870, p. 48; cf. Geheeb in Rev. bryol. 1886, p. 22).*

Br. julaceum *Funck Crypt. exs. fasc. IX, n. 194.*

2. Ad terr. argill. aren. et mur. delaps. — *Eur.* : Suecia, Boruss., Transilv., Hungar., Siles., Bohem., Sudet., Harz, Austr. super., Francon., Bavar., Tirol., Helvet., vall. Rhen., Westphal.; Delphinat., Pyren. *As.* : Jenissei med., Caucas.

var. *Ferchelii* *Schwaegr.* — Br. capillare var. *Ferchelii*.

« var. *tenue* (*Ravaud*) *Boul. M. France, I, p. 257 (1884).*

Br. Payoti (*fd. Limpr.*)

Br. *tenue* *Rav. in Bull. Soc. bot. Fr. 1867, p. 260.*

Ut typ. — *Eur.* : Delphinat.

var.  $\beta$ ? *Bryol. eur.* — Br. Blindii.

*Funckii* *Schwaegr. p.p.* — Br. conicum?

furvum *Hagen in Tromsø Mus. Aarsheft. 1898-99, p. 151.*

Syn. Ad ped. rup. — *Eur.* : Norveg. bor.

fuscens *Rota Cat. M. Bergam.; de Not. Epil. p. 763 (1869).*

Ster. Ad terr. aren. — *Eur.* : Ital. super. (Vall. Tellin.).

*fuscens* Spr. — Br. torquescens f. *monoica*.

fusco-mucronatum *C. M. in Hedwig. 1899, p. 94.*

2. — *Am. merid.* : La Plata.

fuscum *Lindb. in Bot. Not. 1864, p. 67.*

Br. intermedium var. *fuscum* *Bryhn in N. Mag. f. Naturv. p. 126 (1892).*

Syn. In aren. humid. — *Eur.* : Suecia, Fennia.

*fuscum* Ferg. — Br. purpurascens.

*fusiferum* Mitt. — Brachymenium.

- gambirenses* C. M. in Linn. 1871-73, p. 148.  
2. Terr. — Pac. : Austral. or., Tasman.
- Gamblei* Broth. in Act. Soc. sc. Fenn. 1898, n. 2, p. 25.  
2. Terr. — As. : Kashmir.
- ganophyllum* C. M. in Linn. 1880-83, p. 270.  
Ster. — Am. merid. : And. Boliv.
- Gardneri* Mitt. — Br. Pabstianum.
- Garovaglii* de Not. Cron. d. briol. ital. in Comm. II, n. 21 (1866) et Epil. p. 400 (1869); W. P. Sch. op. cit. p. 445.  
2 (de Not.), syn. (Limpr.) — Eur. : Vall. Tellin., m. Stelvio.
- gaspeanum* Kindb. — Br. Duvalii var. *gaspeanum*.  
*Gaudichaudii* Spreng. — Lepthotheca.
- Gayanum* Mont. in C. M. Syn. I, p. 267 (1849) et Syll. p. 33 (1856); Mitt. M. austr. ant. p. 294 (1869).  
Syn. — Afr. : Ins. Kerguelen. Am. merid. : Chilo.
- gedeanum* Bryol. jav. I, p. 147, t. 120 (1855-61).  
Br. *pallens* Dz. et Mk. M. frond. ined. Archip. ind. p. 23 (1854).  
2. Cortic. — As. : Java.
- Geheebii* C. M. { — Br. Combae (fid. de Not.).  
                              { — Br. Funckii (fid. Milde).
- gelidum* Hag. — Webera Schimper.
- gemmascens* Kindb. Eur. and N. Am. Bryin. p. 360 (1897).  
Ster. Rupic. — Am. sept. : Ins. Vancouver.
- gemmaceolum* C. M. in Engler's Bot. Jahrb. 1883, p. 77 et in Forschungsreise..... Bot. p. 13 (1889).  
Syn (nec 2, fid. Broth.). — Afr. : Ins. Kerguelen.
- gemmatum* C. M. in Fl. 1885, p. 401.  
2. Paludic. — Am. merid. : Fuegia.
- gemmiparum* de Not. Cron. briol. ital. in Comm. II, n. 28 (1866) et Epil. p. 406 (1869); W. P. Sch. op. cit. p. 442.  
Br. alpinum { var. *gemmiparum* Husn. Muscol. gall. p. 247 (1889).  
                              { *mediterraneum* de Not. Syll. n. 162 (1838).  
                              { f. *ovata*, *gemmipara* Boul. M. Fr. I, p. 253 (1884).  
Br. *laxifolium* Lindb. in Oefv. af K. Vet.-Akad. Foerh. 1867-68, p. 4.  
Br. *muticum* Lange, Toskanske Mossor, in Bot. Tidskr. Kjobenh. (1868).  
Miehlichhoferia *crassinervia* Jur. in Bot. Zeit. 1866, pp. 20 et 78.  
2. In calcar. et rivul. alv. — Eur. : Anglia, vall. Rhen., Belg., Galloprov., Sardin., Etrur., Graecia. As. : Caucas. (Tiflis). Afr. : Alger.
- gemmuligerum* Kindb. Eur. and N. Amer. Bryin. p. 368 (1897).  
1 et 2. — Am. sept. : Ottawa.
- geniculatum* Brid. Bryol. unit. I, p. 842 (1826); C. M. Syn. I, p. 276 (1849).  
Br. *Webera geniculata* Brid. op. et l. cc.  
— Eur. : Tirol.
- genucaule* C. M. in Nuov. Giorn. bot. ital. 1897, p. 19.  
Br. *terebellum* W. P. Sch. in Mandon Pl. Boliv. n. 1666.  
Ad rivul. — Am. merid. : And. Boliv.

- germinigerum *C. M. in op. cit.* 1896, p. 95.  
 Ster. Rupic. — *As.* : China centr. (prov. Schen-Si).
- Gerwigii (*C. M.*) *Limpr. Laubm. Deutschl.*..... p. 419 (1892).  
 Hypnum Gerwigii *C. M. mss.*; *Jack in Mitt. d. Badish. bot. Ver.* 1891.  
 Ster. Ad rup. calcar. — *Eur.* : Helvet. (Schaffhausen).
- giganteum Sull. — Rhodobryum pseudo-giganteum.
- Gilliesii *Hook. Misc. I, p. 3, t. 2* (1830); *C. M. Syn. I, p. 292* (1849); *Mitt. M. austr. am. p. 288* (1869).  
 Brachymenium Gilliesii *Jaeg. Ad. I, p. 574* (1873-74); *Par. Ind. p. 122* (1894).  
 Br. orthodontioides *ej. op. et l. cc.*  
 Bryum orthodontioides *C. M. op. cit. p. 293 (fid. Mitt.)*.  
 Pohlia Gilliesii *Mont. in d'Orbigny Voy. Am. merid. VII, II, p. 93 et VII, p. 18* (1839).  
 Orthodontium julaceum *Schwaegr. Suppl. II. P. II, II, p. 123, t. 88* (1816).  
 2. Limic. — *Am. merid.* : Amazon., And. Chilens., Argentin. Cordob.; Brasil. ?
- gilvum *Hagen in Tromsø Mus. Aarsheft.* 1898-99, p. 161.  
 Polyoic. Ad aren. delaps. — *Eur.* : Norveg. bor.
- Giraldii *C. M.* — Plagiobryum.
- glaciale Schleich. — Mniobryum albicans var. glacialis.
- glaucescens Dicks. — Ditrichum.
- glaucoviride *C. M. in Linn.* 1880-83, p. 374.  
 Ster. — *Am. merid.* : Argentin. subtropic.
- glaucum { *C. M.* — Brachymenium.  
 { *L.* — Leucobryum.
- globigoma *C. M. in Nuov. Giorn. bot. ital.* 1897, p. 246.  
 Rhodobryum globigoma *Par. Ind. p. 1116* (1897).  
 Ster. — *As.* : China centr. (prov. Schen-Si).
- globirameum *C. M.* — Brachymenium.
- globosum *Lindb. in Oefv. af K. Sv. Vet.-Akad. Foerh.* 18 6, p. 545.  
 Br. mammosum var. globosum *Berggr. in op. cit.* 1875, n. 7, p. 63.  
 Br. rhodocladium *ej. in op. cit. n. 8, p. 27*.  
 2. (Lindb.), syn. (Limpr.). Ad ripas aren. rivul. glacial. — *Eur.* : Spitzberg., Norveg. occid. *As.* : Jenissei infer. *Am. sept.* : Terra nov.
- globosum Mitt. — Brachymenium columbicum.
- gracile Schleich. — Webera commutata var. gracilis.
- gracilentum Tayl. — Br. atro-purpureum.
- gracilescens *C. M.* — Br. schistoneuron.  
 var. decumbens *C. M.* {  
 var. densifolium *C. M.* { — Br. schistoneuron varr. ejusd. nom.  
 var. duplicatum *R. C.* }
- gracilifolium *C. M. in Linn.* 1875, p. 390.  
 Br. coronatum *Hpe. et Ltz. in Krause M. Ecuador, Bot. Zeit.* 1868, p. 812.  
 2. Terr. — *Am. merid.* : And. Quitens.
- gracilisetum *Hsch. Fl. brasil. I, p. 44* (1840); *C. M. Syn. I, p. 339* (1849); *Mitt. M. austr. am. p. 299* (1869).  
 Terr. ? — *Am. merid.* : Brasil. austr. or.
- Graefianum *Schlieph. in Fl.* 1885, n. 19, t. VI.  
 Syn. Rupic. — *Eur.* : Alp. Norveg. (Dovre), Suecia, Helvet. (Via mala).

- « *var. dichroa* Joergens. in *Christiania Vid. Selsk. Foerh.* 1894, p. 85.  
— *Eur.* : Norveg.
- grammocarpaceum* C. M. in *Bull. hb. Boissier*, 1898, p. 32 et in *Ule Coll. M. brasil.* n. 1171.  
2. Terr. — *Am. merid.* : Brasil. or. (S. do Itatiaia)
- grammophyllum* C. M. — Webera.  
*grande* Kindb. — Webera Ludwigii.
- grandiflorum* Arn. in *Rev. bryol.* 1899, p. 36.  
Polyg. Ad terr. marit. — *Eur.* : Suecia.
- grandirete* Kindb. *Eur. and N. Amer. Bryin.* p. 352 (1897).  
Ster. — *Am. sept.* : Columb. brit.
- Griffithianum* Dicks. — *Oedipodium*.
- griseum* Dusen M. *Am. austr. n.* 741.  
Saxic. — *Am. merid.* : Chile austr.
- guatemalense* Hpe. in *sched.*; C. M. in *Bull. hb. Boissier*, 1897, p. 182.  
2. — *Am. sept.* : Guatemala.
- Haamii* Rehm. M. *Afr. austr. n.*  
— *Afr.* : Orange.
- haematocarpaceum* C. M. et Kindb. in *Bidrag.* p. 453 (nom., 1890) et in *Mac. Cat.* p. 125 (1892).  
2. Ad rup. humid. — *Am. sept.* : Columb. brit.
- haematoneuron* C. M. in *Nuov. Giorn. bot. ital.* 1872, p. 17.  
2. — *Afr.* : Abyss.
- haematophyllum* Kindb. in *Mac. Cat.* p. 118 (1892).  
Br. finmarckicum ? (*fid. Hagen*).  
2. Ad rup. humid. — *Am. sept.* : Anticosti, M. Rupestr.
- haematostomum* Joerg. in *Christiania Vid. Selsk. Foerh.* 1894, n. 8.  
Br. salinum Hag. ?  
— *Eur.* : Norveg.
- Hagenii* Limpr. *Laubm. Deutschl.* .... p. 335 (1892).  
Syn. Rupic. — *Eur.* : Norveg. occid.
- « *subsp. curvatum* Hag. et Arn. mss.; Kindb. *Eur. and N. Amer. Bryin.* p. 383 (1897).  
— *Eur.* : Norveg., ins. Aland.
- Haistii* W. P. Sch. *Syn. II ed.* p. 449.  
2. Ad mur. vinear. — *Eur.* : Helvet. (Neufchâtel).
- Hallerianum* Neck. — *Diphyscium foliosum*.
- halophytum* Hagen in *Tromsø Mus. Aarsheft.* 1898-99, p. 142.  
lir. Arnelli J. O. Bomanss. in *Rev. bryol.* 1901, p. 52.  
1. Ad rip. marit. argill. — *Eur.* : Suecia.
- hamipilum* C. M. in *Linn.* 1880-83, p. 381.  
Ster. — *Am. sept.* : Argentin. Tucuman.
- hammatocaulon* C. M. in *Dusen M. Camer. n.* 159.  
— *Afr.* : Cameroon.
- Harveyanum* C. M. — *Brachymenium acuminatum*.
- Hauthalii* C. M. in *Hedwig.* 1897, p. 93.  
Syn. ? — *Am. merid.* : Argentin. Buenos-Ayr.
- Heimii* Gmel. — *Pottia*.

*helveticum* Philib. — *Br. arcticum* var. *helveticum*.

*hemisphaericum* C. M. *Syn.* II, p. 576 (1851).

2. — *As.* : Himalaya, Nepal.

*Hendersoni* Ren. et Card. et Bot. Gaz. 1890, p. 44, t. 7 c.

2. ♂ ign. — *Am. sept.* : Oregon, Calif.

*heteromallum* Dill. — *Dicranella*.

*heteroneuron* C. M. et Kindb. in *Bidrag*, p. 453 (nom., 1890) et in *Mac. Cat.* p. 130 ut subsp. *Br. capillaris* (1892).

*Br. Donianum* Kindb. in *Bull. Torr. bot. Club*, XVII, p. 273 (1890).

*Br. obconicum* ej. in op. cit. p. 274; *Mac. Canad. M. n.* 181.

*Br. provinciale* Mac. in *Coll. cit. n.* 171.

2. Rupic. et terr. — *Am. sept.* : Ontario, Columb. brit., ins. Vancouver.

• var. *brevicuspdatum* Kindb. in *Mac. Cat.* p. 131.

Lignic. — *Am. sept.* : Ins. Vancouver.

*heteroneuron* Spreng. — *Acidodontium*.

*heterophyllum* Hook. — *Mnium*.

*heteropterum pellucidum* Dill. — *Aulacomnium heterostichum*.

*heterostichum* } Arn. — *Aulacomnium*.

                              } Hoffm. — *Rhacomitrium*.

*Hogbergii* C. M. in *Linn.* 1875, p. 391.

*Br. coronatum* Besch. *Prodr. bryol. mexic.* p. 54 (1871).

2. — *Am. sept.* : Mexic.

*Holmgrenii* Lindb. *M. scand.* p. 17 (1879).

*Br. archangelicum* (fid. Kindb. *Eur. and N. Am. Bryin.* p. 379).

*Syn.* — *Eur.* : Lappon.

*homalobolax* C. M. mss.; Ren. *Prodr. fl. bryol. Madag. etc.* p. 167.

— *Afr.* : Madagascar.

*Hookeri* Spreng. — *Brachymenium nepalense*.

*horizontale* (Hpe.) Broth. *Natür. Pflanzenfam.* p. 596.

*Rhodobryum horizontale* Hpe. *Symb.* 1874, p. 498; *Par. Ind.* p. 1116 (1897).

2. Terr. — *Am. merid.* : Brasil, austr. or.

*Hornschuchianum* C. M. — *Brachymenium*.

*Hornschuchii* Spreng. — *Webera longicolla*.

*hornum* Sw. — *Mnium*.

*horridulum* C. M. in *Hedwig.* 1899, p. 67.

2. — *Afr.* : Zambesia merid.

*huillense* W. et Dy. — *Brachymenium*.

*humidulum* Sull. in *Proced. of the Amer. Acad.*.... 1859 p. 278.

2. Ad terr. humid. — *Afr.* : Prom. B. Sp.

*humile* } Mont. — *Orthodontium*.

                  } *pilis carens, viride et pellucidum* Dill. — *Tortula cuneifolia*.

*humillimum* C. M. in *Nuov. Giorn. bot. ital.* 1898, p. 164.

*Br. schensianum* Par. *Suppl. Ind.* p. 72 (1900).

2. — *As.* : China centr. (prov. Schen-Si).

*humillimum* C. M. (Boliv.). — *Anomobryum*.

*hybridum* Bruch. — *Br. cuspidatum*.

*hydrophilum* Kindb. in *Bull. Torr. bot. Club*, p. 95.  
 Ster. Ad scatur. — *Am. sept.* : Ins. Vancouver.

*hygrometricum* Scop. — *Funaris*.

*hymenophylloides* Hartm. — *Mnium*.

*hyperboreum* { Gunn. — *Dicranum*.  
 { Stroem. — *Cynodontium schisti* (p.p. ?)

L. — *Rhacomitrium lanuginosum*.

Schreb. — *Rhacomitrium canescens*.

{ *setis et capsulis exiguis* Dill. — *Rhacomitrium*  
 { *microcarpum*.

*alpinum* { *operculis obtusis* Dill. — { *canescens* var. obtu-  
 { *Rhacomitrium* { sum (fid. *Limpr.*).  
 { sudeticum (fid. *C. M.*).

*aquaticum, calyptris nigris acutis* { *aciculare* (fid. *Limpr.*).  
 Dill. — *Rhacomitrium* { protensum (fid. *C. M.*).

*aquaticum, calyptris nigris acutis* var. *strigosior* Dill. —  
*Rhacomitrium fasciculare*.

*hypnoides* { *ericæ facie, capsulis barbatis, alpinum* Dill. — *Rhacomitrium*  
 { *canescens* var. *ericoides*.

{ *virescens, fasciculare, alpinum* Dill. — *Rhacomitrium*  
 { *fasciculare*.

*hirsutie* { *canescens vulgare* { f. 27 A-E Dill. — *Rhacomitrium*  
 { *canescens*.  
 { f. 27 F-G Dill. — *Rhacomitrium heterostichum*.

*pendulum, sericeum, coma insigni atro-rubente* Dill. —  
*Bryum alpinum*.

*polycephalum, lanuginosum, montanum* Dill. — *Rhacomitrium lanuginosum*.

var. *barbatum* Retz. — *Rhacomitrium canescens* var. *ericoides*.

var. *medium* Leers. — *Rhacomitrium heterostichum*.

var. *microcarpum* Retz. — *Rhacomitrium microcarpum*.

var. *obtusum* Retz. — *Rhacomitrium sudeticum*.

var. *pilosum* Retz. — *Rhacomitrium canescens*.

var. *virescens* Retz. — *Rhacomitrium fasciculare*.

var.  $\alpha$  Neck. — *Rhacomitrium canescens*.

var.  $\beta$  L. — *Rhacomitrium canescens*.

var.  $\gamma$  L. — *Rhacomitrium canescens* var. *ericoides*.

var.  $\gamma$  Huds. — *Rhacomitrium microcarpum*.

var.  $\delta$  Huds. — *Rhacomitrium sudeticum*.

var.  $\eta$  Huds. — *Rhacomitrium protensum*.

*imberbe* L. — *Barbula fallax*.

*imbricatifolium* C. M. — *Brachymerium imbricatum*.

*imbricatissimum* C. M. — *Br. argenteum* var. *costa-ricense*.

*imbricatulum* C. M. *Syn. II*, p. 575 (1851).

2. — *As.* : Himalaya.

*imbricatum* (*Schwaegr.*) *Bryol. eur. fasc.* 6-9 (*Vol. IV*), *Mon. p.* 29, *t.* 240 (1839); *C. M. Syn. I*, p. 303 (1849).

*Br. marginatum* Brid. *Bryol. univ. I*, p. 849 (1826).

*Pohlia imbricata* *Schwaegr. Suppl. I, P. II*, p. 71, *t.* 64 (1816).

2. — *Eur.* : *Rariss.* : *Helvet.*

*imbricatum* de Not., W. P. Sch. (*Syn. II ed.*) — *Br. comense*.

*immarginatum* Bröth. in *Bail. Contrib. Queensl. Fl. Bull.* 13, p. 21 (*nom.*, 1891) et in *Oefv. af K. Sv. Vet.-Akad. Foerh.* 1893, p. 50.

2. ♂ ign. Ad terr. aren. — *Pac.* : Austral. or. (Queensland).

*immersum* Dicks. — *Campylopus flexuosus*.

*imperfectum* Card. in *Rev. Bryol.* 1900, p. 44.

Syn. — *Am. merid.* : Fret. Gerlache.

*inaequale* Tayl. in *Lond. Journ. of bot.* 1846, p. 342; *C. M. Syn. I*, p. 302 (1849).

— *Pac.* : Austral. or.

*inclinans* Dicks. — *Cynodontium polycarpum* var. *strumiferum*.

*inclinatum* (Sw.) *Bryol. eur. fasc.* 6-9 (*Vol. IV*), *Mon.* p. 17, t. 334 (1839); *C. M. Syn. I*, p. 269 p.p. (1849); *W. P. Sch. op. cit.* p. 419.

*Cladodium inclinatum* Brid. *Bryol. univ. I*, p. 621 (1826).

*Hypnum intermedium* W. et M. *Bot. Taschb.* p. 482 p.p. (*fid. C. M.*).

*Leskia inclinata* eor. *op. cit.* pp. 244 et 482 (1807).

*Mnium intermedium* var. *Roehl.* *Deutschl. Fl. III*, p. 93 (1813).

*Pohlia inclinata* Sw. *Musc. Suec.* pp. 45 et 96, t. V, f. II (1799).

Syn. Ad sax., mur., lign., terr. turfos. — *Eur.* : E reg. campestr. usq. ad subalp. in zon temper. *As.* : Japon., penins. Tschutschica, Jenissei infer., Himalaya (var.), Persia, Caucas. *Afr.* : Alger. *Am. sept.* : Fret. Hudson et Kotzebue, Groenland., ins. Miquel., Nov. Scot., Gaspes.; Alaska, fret. Behring, M. Rupestr., reg. occid. et or.

« var. *gracile* Lindb. in *Oefv. af K. Sv. Vet.-Akad. Foerh.* p. 547.

— *Eur.* : Spitzberg. *Am. sept.* : Groenland.

« var. *hydrophilum* Warnst. in *Verh. d. bot. Ver. d. Prov. Brandenb.* 1899, p. 56.

— *Eur.* : Lusat. infer.

« var. *hyperborea* Joergens. in *Christiania Vid. Selsk. Foerh.* 1894, p. 83.

— *Eur.* : Norveg.

var. *longisetum* Fiedl. — *Br. longisetum*.

« var. *magellanicum* Card. in *Rev. bryol.* 1900, p. 41.

— *Am. merid.* : Patagon. austr.

« var. *rimosa* Hagen in *Tromsø Mus. Aarsheft.* 1898-99, 130.

In pratis siccis. — *Eur.* : Norveg. bor.

« subsp. *subinclinatum* Kindb. *Eur. and N. Am. Bryin.* p. 382 (1897).

— *Am. sept.* : Labrador.

*inclinatum* { *Bryol. eur. p.p.* — *Br. longisetum*.  
Dicks. — *Distichium*.  
*plano-operculatum* Breidl. — *Br. archangelicum*.

*inclusum* C. M. in *Fl.* 1890, p. 477.

In gramin. edit. — *Afr.* : Ins. Mayotte.

*inconnexum* Card. in *Rev. bryol.* 1900, p. 44.

Ster. — *Am. merid.* : Fret. Gerlache.

*incurvifolium* C. M. in *Bot. Zeit.* 1851, p. 549.

2. Rupic. — *Pac.* : Nov. Seeland.

*incurvum* Huds. — *Oligotrichum hercynicum*.

*indicum* Dz. et Mk. — *Brachymenium*.

*inflatum* Philib. in *Rev. bryol.* 1886, p. 81.

Syn. — *Eur.* : Norveg.

*inflexum* C. M. — *Philonotis*.

*inordinatum* Brid. — *Aulacomnium palustre*.

*insulare* de Not. — Br. Donianum.

*insularum* J. O. Bomanss. in *Act. Soc. pr. Faun. et Fl. fenn.* 1899, n. 4.  
Syn. Ad litt. mar. — Eur. : Ins. Aland.

*integridens* C. M. — Webera.

*integrifolium* Rehm. — Rhodobryum leucothrix.

*integrisculum* C. M. — Webera.

*intermedium* (Ludw.) Brid. *Mant. M.* p. 120 (1819); *Bryol. eur. fasc.* 6-9 (Vol. IV), *Mon.* p. 47, t. 356 (1839); C. M. *Syn. I.* p. 278 (1849); W. P. Sch. *op. cit.* p. 428.

Br. occidentale Mac. *Canad. M. n.* 390.

Br. pallescens  $\beta$  *intermedium* Schwaegr. *Sp. M. I.* p. 61 (1830).

Br. turbinatum var. *H. et T. Muscol. brit.* p. 122 (1818).

Br. Webera intermedia Brid. *Bryol. univ. I.* p. 632 (1826).

Hypnum *intermedium* W. et M. *Bot. Taschb.* p. 283, t. X, f. 6, *fid. Linpr.* (1807).

Mnium *intermedium* Ludw. Moos. *Cent. I.* n. 81 (1802).

Pohlia intermedia Brid. *Musc. rec. II, P. III.* p. 144, t. 2, f. II (1803).

Webera intermedia Schwaegr. *Suppl. I, P. II.* p. 67, t. 75 (1816).

Syn. Ad sax. et terr. aren. — Eur. : Britann., Suecia, Dania, Boruss., Siles., Saxon., Austr. infer., Voges., Ligur. As. : Jenissei, Himalaya (var.) Am. sept. : Groenland., fret. Hudson, Terra nov., Nov. Scot., Canada merid.; M. Rupestr., Columb. brit., ins. Vancouver, reg. sept., or. et centr. Pac. : Nov. Seeland. ? Tasman.

var. *aristatum* Fiedl.

var. *cirratum* Bryol. eur. } — Br. cirratum.

var. *fuscum* Bryhn. — Br. fuscum.

• var. *Limprichtii* Warnst. *Moossfl. Prov. Brandenb.* p. 52 (1885).  
In arenos. — Eur. : Brandeb.

• var. *microcarpum* Warnst. in *Verh. d. Prov. Brandenb.* 1897, p. 32.  
Ad scaturig. in silvulis. — Eur. : Boruss.

• var. *subcylindricum* Limpr. *Laubm. Deutschl. .... II*, p. 351 (1892).  
Ut et ubi praeced.

*interruptum* Dicks. — *Philonotis fontana*.

*intricatum* Brid. — Br. calophyllum.

*ischyrorhodon* C. M. in *Hedwig.* 1898, p. 103.

Rhodobryum *ischyrorhodon* Par. *Suppl. Ind.* p. 299 (1900).

2. — Pac. : Austral. merid.

*Isleanum* Besch. *Note M. Ins. S. Paul et d'Amsterdam*, p. 3 (1875).

2. — Afr. : Ins. S. Paul et Amsterdam.

*iteratum* Par. — Br. curtum.

*Jackii* C. M. in *Bot. Zeit.* 1864, p. 348.

2. Terr. — Eur. : Alp. Norveg. (Dovre) et Carinth.

*Jamesoni* Tayl. — *Brachymenium*.

*japonense* (Besch.) Broth. *Natür. Pflanzenfam.* p.

*Brachymenium cernuum* W. P. Sch. in *Savatier M. Japon.* n. 385; Par. *Ind.* p. 121 (1894).

Br. *japonense* Besch. in *Ann. sc. nat.* 1893, p. 340.

— As. : Japon.



Joergensenii *Kaur. ap. Joergens. in Christiania Vid. Selsk. Foerh.* 1894, p. 82.

Br. malangense *Kaur. et Arn. mss.*; *Arn. in Rev. bryol.* 1896, p. 39 et in *Bot. Not.* 1899, p. 79 p.p.; *Par. Suppl. Ind.* p. 67.

1. In fiss. rup. schistos. — *Eur.*: Norveg.

julaceo-riparium *C. M. in Ule Bryoth. brasil. n.* 42.

— *Am. merid.*: Brasil. austr. or.

*julaceum* { *Funck.* — Br. *Funckii*.  
Kunth. — Br. *semi-ovatum*.  
Schröd. — Br. *argenteum* var. *majus*.  
Sm. — *Anomobryum filiforme*.

var. *concinnum* *Wils.* — *Anomobryum*.

*juliforme* *W. P. Sch.* — *Anomobryum*.

Junghunianum *Hpe. in Bryol. jav. I*, p. 146, t. 119 (1855-61).

Br. *pachytheca Dz. et Mk. in Pl. Jungh. I*, p. 323.

2. — *As.*: M. Java.

*juniperi foliis, capsulis rectioribus* *Dill.* — *Atrichum angustatum*.

*juniperifolium* *Hoffm.* { *Atrichum angustatum*?  
*Atrichum undulatum* var. *abbreviatum*.

*juniperinum* *C. M.* — *Webera*.

*Juratzkæ Milde Beitr. zur Moosfl. d. Orient.* p. 596 (1870).

2. — *As.*: Persia.

*Kaernbachii C. M. in Act. hort. Petropol.* 1887, fasc. 1.

2. — *As.*: Caucas. (Batoum).

*kashmirens* *Broth.* — *Anomobryum*.

*Kaurini Philib. in Rev. bryol* 1886, p. 21.

Br. *arcticum* (fd. *Limpr. Laubm. Deutschl.*... II, p. 296).

Br. *confluens* *Joerg. Om floraen i Nordreisen*, p. 161 (1894).

1. (Philib.), syn. (Limpr.). Terr. et in fiss. rup. — *Eur.*: Norveg.

*kaurnianum Warnst. in Hedwig.* 1882, n. 2.

Br. *Graesslanum Schlieph. (fd. Hag. in Det. K. Norske vid. selsk. Skript.* 1897, n. 2).

Syn. *Rupic.* — *Eur.*: Norveg. (Dovre), ins. Aland.

*Keniae C. M.* — *Rhodobryum*.

*kerguelense Mitt. in Journ. of the Linn. Soc.* 1877, p. 67 et in *Phil.*

*Trans. of the roy. Soc.* 1877, p. 36, t. III, f. 3.

1. — *Afr.*: Ins. Kerguelen.

*Kiaeri Lindb.* — Br. *Blindii*.

*Killiasii Amann in Ber. d. schweiz. bot. Ges. III*, p. 66 (1893).

Br. *rhaeticum* *ej. in Rev. bryol.* 1892, p. 56.

Syn. Terr. — *Eur.*: Helvet. (summ. Albula).

*Kindbergii Philib. in Rev. bryol.* 1885, p. 83.

Syn. — *Eur.*: Norveg., Helvet. (Vales.).

*Kirkii Broth. in Oefv. af Finska Vet. Soc. Foerh.* 1893, p. 179.

2. Terr. — *Pac.*: Nov. Seeland.

*Klingraeffii W. P. Sch. Syn. II ed.* p. 436 (1876).

Br. *constrictum* *Br. mss.*; *Husn. Musc. gall.* p. 245 (1889).

Br. *erythrocarpum* var. *turfaceum* *W. P. Sch. Syn. I ed.* p. 363 (1860).

Br. *macrostomum* *Jur. in Verh. d. zool. bot. Ges. Wien*, 1867, p. 542.

- Br. pumilum *Saut. in Fl. Herzogt. Salzbg. III, p. 45* (1870).  
 Br. turbinatum forma *Jur. in op. cit.* 1866, p. 486.
2. Ad terr. aren. vel turf. humid. — *Eur.* : Scandin., Boruss., Siles., Gallie., Morav., Hungar., Austr. inf., Salsb., Stir., Carinth., Bavar., Tirol., Helvet., Palatin.
- Klotzchyi* Mitt. — *Brachymenium*.  
*Knowltoni* *Barnes in Bot. Gaz.* 1889, p. 44.  
 Polyg. Rupic. — *Am. sept.* : Terra nov., Nov. Scot.
- Koratranum* C. M. — *Brachymenium julaceum*.  
*Korbianum* C. M. — *Webera*.
- Kunzii* H. et H. — *Br. caespitium var. imbricatum*.  
*labradorensis Philib. in Rev. bryol.* 1887, p. 55 et 1889, p. 61.  
 Polyg. Ad terr. turf. — *Eur.* : Penins. Kola, Dovre, N. Fennia.  
*Am. sept.* : Labrador.
- laceratum* Besch. — *Anomobryum*.  
 var. *comorense* ej. — *A. laceratum var. comorense*.  
*lacerum* Till. — *Bartramia Oederi*.
- lacustre Bland. M. exs. fasc. III, n. 132* (1804); *Bryol. eur. fasc. 6-9* (Vol. IV), *Mon. p. 16, t. 332* (1839); *C. M. Syn. I, p. 268* (1849); *W. P. Sch. op. cit. p. 422*.  
 Br. *Barnesi* *Wood in W. P. Sch. op. cit. p. 471*.  
 Br. *caespitium var. H. et T. Musc. brit. p. 121* (1818).  
 Br. (*Pohlia*) *cernuum Br. in Brid. Bryol. univ. I, p. 851* (1826).  
 Br. *fulvellum Wils. in Kew. Journ. bot. I, p. 330* (1849).  
 Br. *Webera lacustris Brid. op. cit. p. 637*.  
 Hypnum *lacustre W. et M. Bot. Taschb. p. 285* (1807).  
 Mnium *lacustre Schwaegr. Suppl. I, P. II, p. 124, t. 77* (1816).  
 Pohlia *lacustris Hüb. Musc. germ. p. 483* (1833).
- Syn.* In gramin. aren. udis. — *Eur.* : Spitzberg., ins. Ursor., penins. Kola, ins. Faroër, Britann., Suec., Fennia, Dania, Boruss. sept., Siles., Ducat. Megapol., vall. Rhen.; Anglia. *As.* : Amour, Sikkim. *Am. sept.* : Fret. Kotzebue, Alaska, Groenland.
- var. *alpinum* *Bryol. eur.* — *Br. fallax*.  
 var. *angustifolium* *Ruthe.* — *Br. ammophilum*.
- « var. *norvegicum* *Limpr. in LXI Jahrb. d. schlesisch. Ges. f. Vaterl. Kult.* 1884.  
 — *Eur.* : Alp. Norveg. (Dovre).
- « var. *phaeodon* *Hagen in Tromsø Mus. Aarsheft.* 1898-99 p. 128.  
 — *Eur.* : Norveg. bor.
- laete-nitens* C. M. in *Besch. Fl. bryol. Réunion, etc. p. 94* (1881-82).  
 Ster. — *Afr.* : Ins. Franciae.
- laetum* *Lindb.* — *Br. oblongum*.
- laevigatulum* *Broth. in Oefv. af Finska Vet.-Soc. Foerh.* 1898, p. 176.  
 2. — *Pac.* : Nov. Seeland. (Auckland).
- laevigatum* *Hook. f. et Wils. in Journ. of bot.* 1844, p. 546 et *Fl. antarct. I, p. 134, t. 60* (1844); *C. M. Syn. I, p. 319* (1849) et *II, p. 577* (1851); *Handb. of the N. Zeal. Fl. p. 440* (1867).  
 2. — *Afr.* : Ins. Kerguelen, Marion. *Am. merid.* : Ins. Eremit., Maclov., Fuegia. *Pac.* : Austral. or., Nov. Seeland. med. et merid., Tasman.

*Lagerheimii* Joergens. in *Christiania Vid. Selsk. Foerh.* 1894, p. 77.  
Syn. Ad litt. marit. — *Eur.* : Norveg.

*lagunicolum* C. M. in *Bull. hb. Boissier*, 1897, p. 183.  
2. — *Am. sept.* : Guatemala.

*lamprocarpum* C. M. — *Plagiobryum*.

*lamprocomum* C. M. in *Linn.* 1878-79, p. 292.  
— *Am. merid.* : Argentin. Tucuman.

*lamprostegum* C. M. in *Bot. Zeit.* 1853, p. 22; *Mitt. M. Ind. or.* p. 71  
(1859).  
2. — *As.* : N. W. Himalaya, M. Neilgherr.

*lanatum* Brid. — *Br. argenteum* var. *lanatum*.

*lanceolatum* { Dicks. — Pottia.  
                  { *bimum, setis et capsulis* { f. 73 F. — *Br. bimum*.  
                  { *longis pendulis* Dill. } f. A-B. — *Br. pseudo-triquetrum*.  
                  { *pellucidum, capsulis subrotundis, pedicellis carneis* Dill.  
                  { — *Mniobryum carneum*.

*Lansbergii* Dz. et Mk. — *Br. Crügeri*.

*lanuginosum* Hoffm. — *Rhacomitrium*.

*lapponicum* Kaur. et E. Joergens. mss.; *Kindb. Eur. and N. Amer.*  
*Bryin.* p. 382 (1897).

*Br. flagellare* Kaur. ap. Joergens. in *Christiania Vid. Selsk. Foerh.* 1894,  
p. 100 (*fid. Joergens. op. cit.* p. 104).

Syn. — *Eur.* : Lappon., Suecia, Dovre.

*lapponicum* Dicks. — *Amphoridium*.

*laterale* Lightf. — *Bartramia Halleriana*.

*laticeps* C. M. in *Linn.* 1878-79, p. 295.  
Ster. Limic. — *Am. merid.* : Argentin. Cordob.

*latifolium* { Brid. — *Br. Schleicheri* var. *latifolium*.  
                  { Bryol. eur. — *Br. calophyllum*.  
                  { Dz. et Mk. — *Br. Marattii*.

*lato-decurrens* Mac. — *Br. Duvalii* var. *lato-decurrens*.

*lato-marginatum* C. M. in *Sikora Pl. Mad. n.* 14293.  
— *Afr.* : Madagasc.

*Lawersianum* Philib. in *Rev. bryol.* 1899, p. 99.  
Syn. — *Eur.* : Scotia (m. Ben Lawers).

*laxifolium* Besch. *Fl. bryol. Nouv. Caled.* p. 216 (1873).  
2. Paludic. — *Pac.* : Nov. Caledon.

*laxifolium* { Lindb. — *Br. gemmiparum*.  
                  { Warnst. — *Webera Ludwigii*.

*laxo-gemmaceum* C. M. in *Hedwig.* 1899, p. 75.  
2. — *Afr.* : Prem. B. Sp.

*laxum* Reichdt. — *Webera nutans*.

*Lechleri* C. M. in *Bot. Zeit.* 1856, p. 416.

*Br. andicola* (*fid. Mitt.*).

*Br. truncorum* W. P. Sch. in *Lechler Pl. Chil. n.* 626 b.  
*Rhodobryum Lechleri* Par. *Ind.* p. 1117 (1897).

2. Cortic. — *Am. merid.* : Chile austr.

- Leloutrei* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1899, *II*, p. 221.  
Ster. — Afr. : Madagascar (Diego-Suarez).
- lepidopiloides* C. M. — Epipterygium.
- lepidum* Hagen in *Tromsoë Mus. Aarsheft.* 1898-99, p. 138.  
Syn. Ad litt. mar. — Eur. : Norveg. bor.
- lepto-areoblastum* C. M. in *Dusen M. Camer.* n. 71.  
— Afr. : Cameroon.
- leptoblepharon* C. M. — Webera.
- leptocercis* Philib. in *Rev. bryol.* 1894, p. 86.  
Br. Hagenii Limpr. (fid. Hagen in *Det. K. Norske Sells. Skript.* 1897, n. 2).  
Polyg. — Eur. : Ins. Aland.
- leptocladon* Sull. — Anomobryum.
- Leptodontium* Mitt. — Webera.
- lepto-flagellans* C. M. in *Nuov. Giorn. bot. ital.* 1896, p. 96.  
Ster. Terr. — As. : China centr. (prov. Schen-Si).
- leptoglyphodon* Philib. in *Rev. bryol.* 1900, p. 24.  
Syn. In aren. glac. et ad fontes. — As. : M. Alatau.
- leptoloma* C. M. in *Linn.* 1878-79, p. 478.  
2. — Am. merid. : Venezuela.
- leptoloma* Broth. — Br. Uleanum.
- leptopelma* C. M. in *Hedwig.* 1898, p. 88.  
2. ? — Pac. : Austral. or. (Nov. Vales.).
- leptopelmatum* C. M. — Webera.
- leptophyllum* Br. et Sch. — Brachymenium.
- leptospeiron* C. M. in *Besch. Fl. bryol. Réunion, etc.* p. 95 (1881-82).  
2. — Afr. : Madagasc., Comor. magn., ins. Franciæ.
- leptostomoides* C. M. — Brachymenium.
- leptostomum* W. P. Sch. — Anomobryum sericeum.
- leptothecium* Tayl. in *Phyt.* 1844, n. 41, p. 1094; C. M. *Syn. I*, p. 254 (1849).  
Br. truncorum H. f. et W. *Fl. of the N. Zeal. II*, p. 87 (1855), *Fl. Tasman. II*, p. 192 (1860) et *Handb. of the N. Zeal. Fl.* p. 439 (1867).  
Rhodobryum leptothecium Par. Ind. p. 1117 (1897).  
— Pac. : Ins. Societ., Taïti, Austral. or. (Victoria), Nov. Seeland., Tasman., ins. Norfolk et Campbell.
- leptothrix* C. M. in *Hedwig.* 1898, p. 94.  
Ster. — Pac. : Austral. (Trinity-Bay).
- leptothrix* C. M. (Afr.) — Rhodobryum.
- leptotorquescens* C. M. in *Dusen M. Camer.* n. 208; Broth. in *Engler's Bot. Jahrb.* 1897, p. 246.  
2. fr. ign. Rupic. in rivul. — Afr. : Cameroon.
- leptotrichaceum* Rehm. *M. Afr. austr.* n. 256.  
— Afr. : Orange.
- leptotrichum* C. M. in *Hedwig.* 1897, p. 95.  
Muric. — Am. merid. : Argentin. temper.

*Lescurianum* Sull. — Webera.

*leucanthemum* Hpe. mss.; *Mitt. in Trans. and Proceed. of the roy. Soc. of Victoria*, 1883, p. 73.

— *Pac.* : Ins. Lord Howe.

*leucoglyphodon* Philib. in *Rev. bryol.* 1900, p. 73.

1. — *As.* : M. Terskei Alatau.

*leucolomatum* C. M. et Kindb. in *Bidr. p.* 454 (nom., 1890) et in *Mac. Cat.* p. 123 (1892).

Ster. Terr. — *Am. sept.* : Columb. brit.

*leucophyllum* Dz. et Mk. *M. frond. ined. Archip. ind. p.* 15, t. 6 (1854); *C. M. Syn. I.* p. 315 (1849); *Bryol. jav. I.* p. 148 (1855-61).

2. Terr. — *As.* : Java.

*leucothecium* C. M. in *Hedwig.* 1898, p. 106.

*Rhodobryum leucothecium* Par. *Suppl. Ind. p.* 300 (1900).

2. — *Pac.* : Austral. or. (Victoria).

*leucothrix* C. M. (Afr. austr.) — *Rhodobryum*.

*leucurum* C. M. in *Linn.* 1878-79, p. 479.

2. — *Am. merid.* : Venezuela.

*Leveilleum* Dz. et Mk. — *Br. plumosum*.

*Levieri* C. M. in *Hedwig.* 1898, p. 92.

2. Paludic. — *Pac.* : Nov. Seeland. sept.

*Liebmanni* W. P. Sch. in *Besch. Prodr. bryol. mexic. p.* 56 (1871).

*Br. Schimperii* Sb. in *Jaeg. Ad. II.* p. 712 (1877-78).

*Rhodobryum Liebmanni* Par. *Ind. p.* 1117 (1897).

2 ? — *Am. sept.* : Mexic.

*Liebmannianum* C. M. — *Br. brevicaule*.

*ligulatum* Schreb. — *Mnium undulatum*.

*liliputanum* C. M. — *Brachymenium stenopyxis*.

*limbo-marginatum* C. M. in *Fl.* 1896, p. 440.

*Rhodobryum limbo-marginatum* Par. *op. cit. p.* 1118.

Ster. — *Pac.* : Ins. Hawai.

*limbatum* C. M. — Webera.

*limosum* Hagen in *Tromsoë Mus. Aarsheft.* 1898-99, p. 175.

Heteroic. — *Eur.* : Norveg. bor. et merid.

*Limprichtii* Kaur. in *Bot. Not.* 1885, p. 87.

Polyg. Ad rup. madid. — *Eur.* : Alp. Norveg. (Dovre), Finmark, alp. Gray. *Am. sept.* : Groenland.

*Lindbergii* Kaur. in *op. cit.* 1886, p. 129.

1. — *Eur.* : Norveg. occid.

*Lindgrenii* W. P. Sch. *Syn. II ed. p.* 412.

*Br. archangelicum* (fd. Kaur.).

*Br. arcticum* Lindgr. *M. Suec. n.* 60.

*Syn. Terr.* — *Eur.* : Suec. merid.

*Lindmannianum* Broth. in *Bihang till K. Sv. Vet.-Akad. Foerh.* 1900.

2. Terr. — *Am. merid.* : Brasil. austr. (Rio Grande do Sul).

- linearifolium C. M. in *Linn.* 1880-83, p. 365.  
Ster. — *Am. merid.* : Argentin. subtropic.
- lingulatum J. O. Bomanss. in *Act. Soc. pr. Faun. et Fl. fenn.* 1899, n. 4.  
Heteroic. Ad litt. glareos. marit. — *Eur.* : Ins. Aland.
- linoides Dicks. — *Barbula unguiculata*.
- Lisae de Not. *Epil.* p. 483 (1869); W. P. Sch. *op. cit.* p. 457.  
2 (de Not.), syn. ! (Limpr.) — *Eur.* : M. Pedemont.
- liriense Warnst. et Fleisch. *Bryoth. Eur. merid. et in Bot. Centralbl.*  
*XVII Jahrg. Bd.* 65, n. 9, p. 300.  
— *Eur.* : Neapol. (ins. de Liri).
- litorale Ryan et Hag. in *D. K. N. Vid. Skr.* 1896, n. 1.  
Br. pycnoderium *Limpr. forma?*  
2. Polyg. — *Eur.* : Norveg. bor.
- litorum J. O. Bommanss. in *Rev. bryol.* 1898, p. 10 et in *Act. Soc. pr. Faun. et Fl. fenn.* 1899, n. 4.  
1. Ad litt. mar. — *Eur.* : Ins. Aland.
- lonchocaulon C. M. in *Fl.* 1875, p. 93.  
Syn. — *Am. sept.* : Colorado.
- lonchoneuron C. M. in *Hedwig.* 1898, p. 91.  
2. — *Pac.* : Austral. or. (Nov. Vales.).
- lonchophyllum Broth. — Br. zonatiforme.
- lonchopyxis C. M. in *Hedwig.* 1899, p. 72.  
2. — *Afr.* : Prom. B. Sp.
- lonchotrachylon C. M. — *Brachymenium*.
- longescens C. M. — *Webera*.
- longibracteatum Kindb. — *Mniobryum*.
- longicollum Sw. { — *Webera cruda* var. minor?  
— *Webera*.
- longifolium { C. M. — *Brachymenium*.  
C. M. et Hpe. — *Webera*.  
Dicks. — *Dicranum falcatum*.  
Hoffm. — *Dicranum*.
- longipedicellatum C. M. — *Brachymenium*.
- longirostrum { Brid. — *Mnium rostratum*.  
Griff. — *Mnium rostratum* var. *rhynchophorum*.
- longisetum Bland. *M. exs. fasc. V*, n. 222 (1808); *Brid. Bryol. univ. I*,  
p. 673 (1826); W. P. Sch. *op. cit.* p. 420.  
Br. inclinatum var. longisetum Fiedl. *Syn. Laubm. Mecklenb.* p. 66 (1844).  
Br. inclinatum forma *Bryol. eur. fasc. 6-9 (Vol. IV)*, Mon. p. 17, t. 334  
p.p. (1839); C. M. *Syn. I*, p. 269 p.p. (1849).  
Br. turbinatum var. *H. et T. Musc. brit. II ed.* p. 292 (1827).  
Br. Willdenowii Jur. in *Rab. Bryoth. eur. n.* 973 (1867).  
Meesea nutans Willd. *Prodr. fl. berol. n.* 983 (1787).  
Pohlia longiseta Hüb. *Musc. germ.* p. 480 (1833).  
Syn. In pratis palud. — *Eur.* : German. sept., Wurtemb., Bavar.
- Lorentzii W. P. Sch. *op. cit.* p. 413.  
Syn. Ad terr. turf. et in sphagn. — *Eur.* : Norveg. (semel lectum).
- lucidum { D<sup>na</sup> E. G. Britt. — *Roellia*.  
James. — *Webera cruda* var. *lucida*.

- Ludwigii* } Brid. — *Webera commutata*.  
               } Spreng. — *Webera*.  
               var. *gracile* Bryol. eur. — *Webera commutata* var. *gracilis*.  
*luridum* Ruthe in *Verh. d. bot. Ver. d. Prov. Brandenb.* 1867; W.  
               *Sch. op. cit.* p. 410.  
               Syn. In cav. arenos. — *Eur.* : German. sept. (Neomark).  
*luteolum* } C. M. — *Brachymenium*.  
               } Hoffm. — *Weisia viridula*.  
               Dicks. — *Rhacomitrium fasciculare*.  
*lutescens* } Kindb. — *Webera*.  
               J. O. Bomanss. — *Br. finmarkicum*.  
*lycopodiiforme* Schleich. — *Philonotis fontana*.  
*lycopodioides* } Hook. — *Mnium*.  
                       } Sw. — *Syrrophodon*.  
*maceratum* C. M. in *Linn.* 1880-83, p. 385.  
               2. — *Am. merid.* : Argentin. subtropic.  
*Mac-Leanum* C. M. in *Hedwig.* 1899, p. 74.  
               — *Afr.* : Transvaal.  
*Macounii* Aust. — *Br. Atwateriae*.  
*macrantherum* C. M. in *Engler's Bot. Jahrb.* 1883, p. 77 et in *Forschungs-*  
               *reise..... Bot.* p. 14 (1889).  
               *Br. pendulum* Mitt. in *Journ. of the Linn. Soc.* 1876.  
               Syn. — *Afr.* : Ins. Kerguelen, Marion.  
*macrocarpum* Hedw. — *Leptostomum*.  
*macrodemissum* C. M. — *Plagiobryum*.  
*macro-erythrocarpum* C. M. in *Hedwig.* 1898, p. 92.  
               2. Ad rup. humid. — *Pac.* : Nov. Zeland. austr.  
*macro-gracilescens* C. M. in *Hedwig.* 1898, p. 224.  
               Ster. — *Am. sept.* : Haiti.  
*macropelma* C. M. *Syn. I*, p. 275 (1849).  
               2. — *Am. merid.* : Chile austr.  
*macropelma* C. M. (Austral.). — *Br. curtum*.  
*macropoma* C. M. — *Brachymenium*.  
*macrostomum* Jur. — *Br. Klinggræffii*.  
*magellanicum* C. M. — *Brachymenium*.  
*malachiticum* C. M. — *Brachymenium*.  
*malacodictyon* C. M. in *Hedwig.* 1898, p. 92.  
               2. — *Pac.* : Nov. Seeland occid.  
*malangense* Kaur. et Arn. p.p. } — *Br. Joergensenii*.  
   } — *Br. salinum*.  
*mammillatum* Lindb. in *Not. ur Saellsh. pr. Faun. et Fl. fenn. Foerh.*  
               IX (1868); W. P. *Sch. op. cit.* p. 417.  
               1. Ad rip. aren. fluv. et in aren. marit. — *Eur.* : Ins. Ursor., Aland,  
               Gothland, Pomeran.  
               var. *globosum* Berggr. — *Br. globosum*.  
*mammilligerum* Kindb. in *Mac. Cat.* p. 122 ut subsp. *Br. inclinati* (1892).  
               Syn. Ad rup. humid. — *Am. sept.* : M. Rupestr.  
*mammillosum* C. M. — *Brachymenium*.

*mancum* C. M. — Webera?

*Mandoni* C. M. — Epipterygium.

*Mangiferae* C. M. — Webera.

*Marattii* Wils. *Bryol. brit. App.* p. 9, t. 32, f. b. (1855); *Bryol. eur. fasc.* 62-64 (Vol. IV), *Mon. Suppl.* p. 60, t. 640 (1855); *W. P. Sch. op. cit.* p. 423.

*Br. latifolium* *Ds. et Mh. Prodr. fl. batav.* (1851).

1. In aren. humid. et ad rip. turf. — *Eur.* : Ins. Faroër, Anglia, Neerland., Suecia, Dania, ins. Aland et Gothland, Borussiae marit.

*marchicum* Roth. — Philonotis.

*marginatum* *Bryol. eur. fasc.* 6-9 (Vol. IV), *Mon.* p. 74, t. 379 sub *Br. inclinatum* (1839); *C. M. Syn. I.* p. 297 (1849); *W. P. Sch. op. cit.* p. 437.

2. Ad rup. arenac. — *Eur.* : Prope Bipontem Voges. semel lectum (*W. P. Sch.*); pr. « Bar-sur-Seine » et Andegav. Galliae (C. M.).

*marginatum* { *Brid.* } — *Br. imbricatum.*  
                   { *Dicks.* } — *Epipterygium Tozeri.*  
                   { *Dicks.* } — *Mnium serratum.*

var. *orthorhynchum* Hartm. — *Mnium orthorhynchum.*

var. *Drumm.* — *Mnium inclinatum.*

*Mariei Besch. in Fl. bryol. Réunion, etc.* p. 94 (1880-81).

2. Ad cement. calcar. — *Afr.* : Nossi Be.

*maritimum* *J. O. Bomanss. in Rev. bryol.* 1897, p. 1 et in *Act. Soc. pr. Faun. et Fl. fenn.* 1899, n. 4.

1. In aren. marit. — *Eur.* : Finmark, Suecia, ins. Aland.

*Massabiae* C. M. in *Hedwig.* 1898, p. 224.

2. — *Am. merid.* : Aequator litt.

*matto-grossense* *Broth. in Bihang till K. Sv. Vet.-Akad. Foerh.* 1900.

2. Ad saxa hinc inde inund. — *Am. merid.* : Brasil. centr. (*Matto Grosso*).

*maynense* *Spr. in Mitt. M. austr. am.* p. 312 (1869).

2. Ad saxa ligna. inund. — *Am. merid.* : Ins. Trinitat., And. Quintens. et Peruv.

*medianum* *Mitt. M. Ind. or.* p. 74 (1858).

*Rhodobryum medianum* *Par. Ind.* p. 1118 (1897).

2. — *As.* : Khasia, m. Neilgherr.

*Mayrii* *Broth. in Hedwig.* 1899, p. 218.

*Syn.* — *As.* : Ins. Kuril.

*medium* Vill. — *Cynodontium polycarpum.*

*meesoides* *Kindb. in Bull. Torr. bot. Club,* 1889, p. 95.

2. Ad rup. humid. — *Eur.* : Norveg. *Am. sept.* : Anticosti, Terra nov., Gaspes., Minnesota, M. Rupestr. et Selkirk, ins. Vancouver.

*megalacron* Schwaegr. — *Brachymenium.*

*megalocarpum* Hook. — *Acidodontium.*

*megalodictyon* Sull. in *Proced. of the Amer. Acad.*.... 1859, p. 279.

*Br. cellulare* Hook.?

2. Muric. — *As.* : Ins. Loo-Choo.

*megamorphum* C. M. — Webera.



*melanodon* Brid. — *Mniobryum carneum*.

*melanopyxis* C. M. — *Brachymenium*.

*melanothecium* C. M. — *Brachymenium*.

*Melvilleanum* C. M. *Syn. I*, p. 267.

*Ptychostomum pulchellum* R. Br. *App. ad iter Parray*. p. 197 et in *Brid. Bryol. univ. I*, p. 601 (1826).

— *Am. sept.* : Ins. Melville.

*membranaceum* C. M. in *Bal. Pl. Parag. n.* 3051 a.

— *Am. merid.* : Paraguay.

*membranaceum, foliis membranaceis obtusis* Dill. — *Aulacomnium heterostichum* var. *humile*.

*Menziesii* Hook. — *Rhizogonium*.

*meruense* C. M. in *Fl.* 1890, p. 475.

2. — *Afr.* : Kilima N'Djaro.

*mexicanum* C. M. — *Brachymenium*.

*Meyenianum* C. M. — *Brachymenium*.

*micans* *Limpr. Einig. neue Art.* 1883, p. 236 et *Laubm. Deutschl.... II*, p. 303.

*Polyg. Ad terr. turfos.* — *Eur.* : Ins. Faroër, Norveg. *Am. sept.* : Fret. Scoresby.

*micans* Mitt. — *Brachymenium*.

*micro-alpinum* C. M. in *hb. Saharampour*, n. 7627.

— *As.* : N. W. Himalaya.

*microblastum* C. M. in *Bot. Centralbl.* 1883, n. 2-4.

*Syn. (fd. Broth.)*. — *As.* : Penins. Tschutschica.

*micro-calophyllum* *Philib. in Rev. bryol.* 1899, p. 13.

1. — *As.* : M. Terskei Alatau.

*micro-capillare* C. M. in *Hedwig*, 1898, p. 264.

*Br. caespiticium* Sull. in *Wright M. cubens. n.* 61.

2 ? *Terr.* — *Am. sept.* : Cuba.

*microcarpon* (*Hsch.*) C. M. *Syn. I*, p. 294 (1849); *Mitt. M. austr. am.* p. 300 (1869).

*Cladodium microcarpon* *Hpe. Enum. M. Brasil.* p. 30 (1879).

*Pohlia microcarpa* *Hsch. Fl. brasil. I*, p. 38 (1840).

2. *Terr.* — *Am. merid.* : Brasil. austr. or.

*microcarpum* Gmel. — *Rhacomitrium*.

*micro-cephalum* C. M. et *Kindb. in Bidr.* p. 454 (*nom.*, 1890) et in *Mac. Cat.* p. 134 (1892).

2. *Rupic.* — *Am. sept.* : Columb. brit.

*microchaeton* *Hpe. in Ann. sc. nat. V ser., III*, p. 342 (1866); *Mitt. M. austr. am.* p. 298 (1869).

2 ? — *Am. merid.* : Brasil. austr. or.

*micro-commutatum* *Kindb.* — *Webera norvegica*.

*micro-comosum* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 19.

*Rhodobryum micro-comosum* *Par. Ind.* p. 1118 (1897).

— *Am. merid.* : And. Boliv.

*microdontium* *Kindb.* — *Webera micro-denticulata*.

- micro-erythrocarpum *C. M. et Kindb. op. et ll. cc. pp. 454 et 124.*  
*Br. murale Mac. Canad. M. n. 186.*  
 2. In aren. humid. — *Am. sept.* : Ins. Vancouver.
- microglobum *C. M. et Kindb. op. et ll. cc. pp. 454 et 129.*  
 2. Terr. — *Am. sept.* : Ontario.
- microglossum *C. M. in Hedwig. 1897, p. 94.*  
 2. — *Am.* : Argentin, Patagon.
- micro-laevigatum *C. M. in Engler's Bot. Jahrb. 1883, p. 77 et in Forschungsreise.... Bot. p. 12 (1889).*  
 Ster. — *Afr.* : Ins. Kerguelen.
- micron *C. M. in Linn. 1880-83, p. 372.*  
 2. — *Am. merid.* : Argentin.
- micro-pendulum *C. M. in op. cit. 1878-79, p. 476.*  
 2. — *Am. merid.* : Venezuela.
- micropendulum *C. M. (Argent.) — Br. nano-pendulum.*
- microrhodon *C. M. in Hedwig. 1898, p. 108.*  
*Rhodobryum tasmanicum Par. Suppl. Ind. p. 301 (1900).*  
 2. — *Pac.* : Tasman.
- microsporum *Broth. in Oefv. af Finska Vet.-Akad. Foerh. 1899, p. 100.*  
 2. — *Pac.* : Tasman.
- microsporum *Kindb. — Webera pycnodecurrans.*
- microstegioides *Kindb. in Ott. Nat. 1900, V, p. 88.*  
 — *Am. sept.* : Ins. Pribylof.
- microstegium *Bryol. eur. fasc. 32, Mon. Suppl. p. 7, t. 362 (1846); C. M. Syn. I, p. 279 (1849); W. P. Sch. op. cit. p. 433.*  
 Syn. Terr. — *Eur.* : Alp. Norveg., Tirol., Helvet. *Am. sept.* : Groenland., Labrador.
- microstomum { *C. M. — Brachymenium.*  
 { *Dicks. — Hymenostomum.*
- microtheca *C. M. Syn. I, p. 314.*  
 2. — *As.* : Manill.
- microthecium *C. M. in Hedwig. 1898, p. 95.*  
 Syn. — *Pac.* : Austral. or. (Nov. Vales.).
- Miehlichhoferia C. M. — Webera.*  
*miehlichhoferiaceum C. M. — Brachymenium.*
- Mildeanum *Jur. in Verh. d. zool. bot. Ges. in Wien, 1862, p. 967; W. P. Sch. op. cit. p. 446.*  
*Br. rubrum Milde mss.; Ltz. Bryol. Notizb. p. 46 (1865).*  
 2. Ad terr. humid. et mur. — *Eur.* : Siles., Hungar., Sudet., Bohem., Austr. inf., Salisb., Stir., Carinth., Carniol., Bavar., Tirol., Helvet., Delphin., Rhoengeb., Francon., Westphal., Belg. *As.* : Caucas.
- miniaturum *Lesq. in Mem. Acad. Calif. I, p. 23; Lesq. et Jam. Man. p. 233 (1884).*  
 2. Ad rup. humid. — *Am. sept.* : Columb. brit., Calif.
- miniaturum *Mac. — Br. capitellatum.*  
*minus (Dicranum?) f. pumila Mich. — Leucobryum minus f. pumila.*

- minuscolum* C. M. in *Fl.* 1885, p. 400.  
 Syn. ! (*fid. Broth.*), nec 2. Paludic. — *Am. merid.* : *Fuegia austr.*
- minutirete* C. M. in *Fl.* 1888, p. 409.  
 Ster. — *Afr.* : Kilima N'Djaro.
- minuti-rosatum* C. M. in *Fl.* 1828, p. 408.  
 Br. roseum Mitt. in *Journ. of the Linn. Soc.* 1886, p. 307 ?  
*Rhodobryum minuti-rosatum* Par. Ind. l. c.  
 Ster. In graminos. — *Afr.* : Kilima N'Djaro.
- minutissimum* C. M. in *Hedwig.* 1898, p. 88.  
 2 ? — *Pac.* : Austral. or. (Queensland).
- minutulum* W. P. Sch. in *Besch. Prodr. bryol. mexic.* p. 53 (1871).  
 2. — *Am. sept.* : Mexic.
- minutum* { Dicks. — *Pottia Starkeana*.  
 { Gmel. — *Seligeria pusilla*.
- mirabile* C. M. in *Bot. Centralbl.* 1883, n. 2-4.  
 Syn. — *As.* : Penins. Tschutschica.
- misandrum* Hagen in *Tromsø Mus. Aarsheft.* 1898-99, p. 178.  
 Syn. In fiss. rup. alp. — *Eur.* : Norveg. bor.
- mnioides* (W. P. Sch.) Broth. *Natür. Pflanzenfam.* p. 552.  
*Webera mnioides* W. P. Sch. in *Besch. Fl. bryol. Antill. fr.* p. 30 (1876).  
 2. — *Am. sept.* : Guadalup.
- mnioides* { Gmel. — *Leptobryum piriforme*.  
 { Wils. — *Mnium subglobosum*.
- Moei* W. P. Sch. *Syn. II ed.* p. 417 (1876); *Philib. in Rev. bryol.* 1886, p. 20.  
 Syn. — *Eur.* : Alp. Norveg.
- Mohrii* Lesq. in *Bull. Torr. bot. Club*, 1874, p. 50 (*in Man. non memor.*).  
 — *Am. sept.* : Ludovic.
- Molleri* C. M. — *Brachymenium*.
- molliculum* *Philib. in Rev. bryol.* 1899, p. 81.  
 1. — *As.* : M. Alatau.
- moniliforme* C. M. in *Dusen M. Camer.* n. 240 a.  
 — *Afr.* : M. Cameroon.
- Montagneanum* C. M. — *Brachymenium pendulum*.  
 C. M. — *Webera*.  
 { Lam. — *Distichium capillaceum*.
- montanum* { Sm. — *Bartramia pomiformis* var. *crispa*.  
 { *hemiheterophyllum, operculis acutis* Dill. — *Rhacomitrium aciculare*.
- Morisi* Br. — *Br. erythrocarpum*.
- mucronatum* Mill. in *Handb. of the N. Zeal. Fl.* p. 442 (1867).  
 Hermaphr. Terr. — *Pac.* : Nov. Seeland.
- mucronifolium* *Philib. in Rev. bryol.* 1899, p. 27.  
 1. — *As.* : M. Alatau.
- mucronigerum* *Philib. in op. cit.* 1900, p. 91.  
 2 vel syn. — *Am. sept.* : Prince William's Sound, prom. Fox.
- mucronulatum* Dicks. — *Barbula unguiculata*.

Mühlenbeckii *Bryol. eur. fasc. 32 (Vol. IV), Mon. Suppl. p. II, t. 381* (1846); *C. M. Syn. I, p. 286* (1849); *W. P. Sch. op. cit. p. 441*.

Br. alpinum var. brevifolium *Myr. Coroll. fl. Upsal. p. 68* (1833).

Br. Raii *Aust. in Coult. Bot. Gaz. II, p. 110*.

2. Ad rup. madid.; calcar. fugit. — *Eur.*: Scand., Riesengeb., Gosenke, Tatra, Salisb., Stir, Carinth., Tirol., Helvet. *As.*: Caucas. *Am. sept.*: Groenland., Terra nov., Wyoming, Oregon, Washington, Nov. Mexic.

*Mülleri* Spr. — Br. Donianum.

multicaule *Tayl. in Lond. Journ. of bot. 1846, p. 342; C. M. Syn. I, p. 302* (1849).

— *Pac.*: Austral. or.

multiflorum *C. M. — Webera*.

Mundtii *C. M. in Bot. Zeit. 1859, p. 206*.

Br. canariense *ej. Syn. I, p. 253 pp.* (1849).

2. Cortic. — *Afr.*: Prom. B. Sp.

« var. flaccidum *C. M. in op. et l. cc.*

Ut et ubi typ.

murale *Wils. in Milde Bryol. siles. p. 213* (1869); *W. P. Sch. op. cit. p. 437*.

Br. atro-purpureum *de Not. Epil. p. 399* (1869) *et Erb. critt. ital.*

Br. erythrocarpum γ murorum *W. P. Sch. Syn. I ed. p. 363* (1860); *Mandon Pl. Madeir. n. 21; Rab. Bryoth. eur. n. 674*.

Br. murorum *Wils. (nom. sol.) Ltz. Bryol. Notizb. p. 46* (1865).

2. Ad cement. calcar. mur. — *Eur.*: Britann., vall. Rhen., Gall., Helvet. occid., Ital., Astur. *As.*: Caucas. *Afr.*: Alger., Madeir.

murale } *Mac.* — Br. micro-erythrocarpum.  
          } *L.* — Tortula muralis.

var. β *Huds.* — Tortula cuneifolia.

murmanicum *Broth. in Act. Soc. pr. Faun. et Fl. fenn. VI, n. 4, p. 58* (1890).

*Syn. Arenic.* — *Eur.*: Lappon. Murman.

murorum *Wils.* — Br. murale.

muticum *M. T. Lge.* — Br. gemmiparum.

mutilum *Hagen in Tromsø Mus. Aarsheft. 1898-99, p. 124*.

*Syn. Inter gramin. et musc. al.* — *Eur.*: Norveg. bor.

Myurella *C. M.* } — Brachymenium.

                  } — Brachymenium myurelloides.

myurelloides *Par.* — Brachymenium.

nagasakense *Broth. in Hedwig. 1899, p. 219*.

2. Rupic. ad rivul. — *As.*: Japon

nanocoma *C. M. in Linn. 1880-83, p. 365*.

2. — *Am. merid.*: Argentin. Tucuman.

nano-pendulum *ej. in op. cit. op. 363*.

Br. micro-pendulum *ej. in op. cit. 1878-79, p. 294*.

*Syn.*

— *Am. merid.*: Argentin. Tucuman.

nanophyllum *C. M. in Nuov. Giorn. bot. ital 1897, p. 18*.

2 ?

— *Am. merid.*: And. Boliv.

- nanopyxis* C. M. in *Hildebr. M. Madag.* n. 2050.  
— Afr. : Nossi-Be.
- nanorhodon* C. M. in *Besch. Fl. bryol. Réunion, etc.* p. 90 (1880-81).  
Ster. — Afr. : Ins. Franciae.
- nano-rosula* C. M. — *Rhodobryum roseum*.
- nano-torquescens* C. M. in *Fl.* 1890, p. 476.  
Ster. — Afr. : Kilima N'Djaro.
- nanum larignis foliis albis* Dill. — *Octoblepharum albidum*.
- naviculare* Hpe. *Symb.* 1877, p. 724.  
Ster. — Am. merid. : Brasil, austr. or.
- naviculare* R. C. — *Webera cucullata* var. *carinata*.
- Neesii* C. M. — *Brachymenium*.
- neilgherriense* Mont. in C. M. *Syn. I*, p. 255 (1849) et *Syll.* p. 32 (1856).  
Mnium ramosum (fid. Mitt.).  
*Rhodobryum neilgherriense* Par. Ind. p. 1118 (1897).  
2. — As. : M. Neilgherr.
- neilgherriense affine* Wils. — *Rhodobryum roseum*.
- nemicaulon* C. M. — *Webera*.
- neodamense* Itzigsohn in *Fl.* 1848, I, p. 360 ; C. M. *Syn. I*, p. 268 (1849) ;  
W. P. Sch. op. cit. p. 460.  
Br. cavifolium Wils. mss. ; Braithw. Brit. Moossf. 1894, p. 188.  
Br. pseudo-triquetrum var. *cavifolium* W. P. Sch. Coroll. p. 76 (1856).  
2. In uligin. — Eur. : Spitzberg, Britann., Scandin., Dania, German.  
sept., Neerland., vall. Rhen., Pyren. Am. sept. : Fret. Scoresby.  
var. *ovatum* L. et A. — Br. *ovatum*.  
Dz. et Mk. — *Brachymenium brevicaulis* ?
- nepalense* { Mitt. — *Brachymenium*.  
Spreng. — *Brachymenium Weisia*.
- nervosum* Hoffm. — *Barbula unguiculata* var. *cuspidata*.
- nevadense* C. M. — *Webera*.
- Nietneri* C. M. — *Brachymenium*.
- nigrescens* Vill. { *Rhacomitrium aciculare* (fid. Limpr.).  
Rhacomitrium protensum (fid. C. M.).
- nigricans* Kaurin in *Hagen M. Norveg. bor.* 1898-99, p. 172.  
Syn. Ad terr. aren. — Eur. : Norveg. bor.
- nigricans* { Crome. — Br. *caespitium*.  
Dicks. { — Br. *turbinatum* (fid. C. M.).  
— Mniobryum carneum (fid. Limpr.).
- nigro-purpureum* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 18.  
2. — Am. merid. : And. Boliv.
- nigritum* Dicks. — *Catoscopium*.
- nimbigenum* Tayl. — *Anomobryum concinnatum*.
- nitens* Hook. in *Wall. Cat. n.* 7592 et *Is. pl. rar. t.* 20, f. 6 (1841) ;  
Harv. in *Journ. of bot.* 1840, II, p. 9 ; C. M. *Syn. I*, p. 339 (1849) ;  
Mitt. M. Ind. or. p. 70 (1859).  
Br. *coronatum* var. Wils. in *Kew Journ. bot.* IX, p. 333 (1857).  
Br. *pachypoma* Mont. (fid. Mitt.).  
2. Lignic. — As. : Nepal., Ceylan, Java. Afr. : Ins. Rodriguez.
- nitens* }  
*nitescens* } Kindb. — *Webera cruda* var. *minor*.

nitido-nigrum C. M. in *Sikora Pl. Madag.* n. 14294-95.

— *Afr.* : Madagascar

nitidulum Lindb. in *Oefv. af K. Sv. Vet.-Akad. Foerh.* 1866, p. 545.

Syn. — *Eur.* : Spitzberg., Scand. *Am. sept.* : Groenland.

nitidum	foliis	Serpilli	Mitt. — Anomobryum.	
			inferioribus Serpilli, superioribus magis.... Dill. —	
			Mnium affine.	
			angustioribus	majus ej. — Bryum pseudo- triquetrum. medium ej. — Webera cruda. minus ej. — Bryum pal- lens.
			pellucidis	capitulis reflexis.... ej. — Bryum capillare. oblongis, capitulis.... ej. — Mnium cuspidatum. rotundis, capitulis.... ej. — Mnium punctatum. subrotundis, elatius ej. — Mnium affine.
			oblongis undatis.... ej. — Mnium undulatum.	

nivale C. M. *Syn. I*, p. 262 (1849).

Syn. — *Am. merid.* : Summ. And. Chilens.

norvegicum } Günn. — Bartramia Halleriana.

Kindb. — Webera micro-commutata.

Notarisii Mitt. — Haplodontium.

nudicaule Lesq. — Webera.

nudum Arn. in *Rev. bryol.* 1897, p. 7.

Syn. Ad arbor. period. submers. — *As.* : Jenissei infer.

nudum Dicks. — Discelium.

H. f. et W. — Webera clavaeformis ?

nutans } Mitt. } — Webera sphagnadelphus ?

(1876). — Webera pseudo-nutans.

Schreb. — Webera.

var. minus Kth. — Mielichoferia longiseta.

var. pinetorum H. M. — Webera nutans var. strangulata (1).

var. Drumm. — Br. arcticum.

nutanti-polymorphum C. M. — Webera.

obconicum Hsch. in *litt.* ; *Bryol. eur. fasc.* 6-9 (Vol. IV), *Mon.* p. 59, t. 367 (1839) ; *C. Syn. I*, p. 282 (1849) et *II*, p. 571 (1851) ; *W. P. Sch. op. cit.* p. 451.

Br. capillare var. obconicum Hüb. *Musc. germ.* p. 412 (1833).

2. Ad mur. et rup. aren. — *Eur.* : Angl., vall. Rhen., Hercyn., Pyren.

*Am. sept.* : Minnesota, Colorado, Calif., Florid. *Pac.* : Austral., Nov. Seeland., Tasman.

« f. longipes Besch. in *Ann. sc. nat.* 1892, I, p. 66.

Terr. — *As.* : Yunnan.

obconicum } Kindb., Mac. — Br. heteroneuron.

Sull. — Br. sanguilentum.

(1) Omnes var. *Bryol. eur.* et subspec. *Kindbergii* ut var. et subspec. ejusd. nom. ad *Weberam nutantem* relatae sunt.

- obliquatulum *Dus. M. Am. austr. n.* 365.  
Terr. — *Am. merid.* : Chile austr.
- obliquum *C. M. Bryol. austr. Georg. p.* 17 (1890).  
2. — *Am. merid.* : Austr. Georg.
- oblongum *Lindb. in Act. Soc. pr. Faun. et Fl. fenn. 1883, p.* 127 et in *Rev. bryol.* 1886, p. 33.  
Br. laetum *ej. in Bot. Not. 1883, p.* 64 et in *op. cit.* 1886, p. 252.  
2. Terr. — *Eur.* : Norveg., Fennia. *As.* : Jenissei med.
- obovatum *Mitt. in Journ. of the Linn. Soc.* 1863, p. 4.  
2. Terr. — *Afr.* : Madeir.
- obtusatissimum *C. M. in Nuov. Giorn. bot. ital.* 1897, p. 26.  
— *Am. merid.* : And. Boliv.
- obtusatissimum *C. M.* (Costa Rica). — *Anomobryum.*
- obtusifolium *Lindb. in Oefv. af Finska Vet.-Akad. Foerh.* 1886, p. 544.  
2. Ster. — *Eur.* : Spitzberg., Septem Ins., ins. Parry, Urior., Norveg.  
*As.* : Penins. Samojed., Jenissei infer. *Am. sept.* : Groenland, fret. Behring.
- obtusifolium } Turn. — Br. cyclophyllum.  
Vill. — *Webera cucullata.*
- obtusissimum *Ren. et Card. in Bull. Soc. roy. bot. Belg.* 1899, II, p. 222.  
Ster. In aquos. — *Am. sept.* : Costa-Rica.
- obtusum *Dicks.* — *Entosthodon ericetorum.*
- occidentale *Sull. in Bot. Pac. R. R. Survey (Whipple), p.* 188, t. 7;  
*Lesq. et Jam. Man. p.* 236 (1884).  
2. Rupic. et terr. — *Am. sept.* : Ottawa, Montana, Oregon, Calif.
- occidentale *Mac.* — Br. intermedium.  
*var.* *Sull. et Lesq.* — Br. sanguilentum.
- occultum *C. M. in Hildebr. M. Madag. n.* 2135.  
— *Afr.* : Madagasc. centr.
- ochropyxis *C. M. in Rehm. M. Afr. austr. n.* 283.  
— *Afr.* : Prom. B. Sp.
- octoblepharis *Gmel.* — *Octoblepharum albidum.*  
*Ederi* *Gunn.* — *Bartramia.*
- oediloma *C. M. in Bihang till. K. Sv. Vet.-Akad. Foerh.* 1895, p. 3.  
2. ♂ ign. Ad saxa rivul. — *Am. merid.* : Brasil. or.
- oedoneuron *C. M.* — *Webera.*
- oelandicum *Philib. in Rev. bryol.* 1887, p. 9.  
Br. warneum (*fd. Linpr.*).  
1. In aren. marit. — *Eur.* : Ins. Oeland.
- oeneum *Blytt in Bryol. eur. fasc. 32, Mon. Suppl. p.* 4, t. 338 (1846);  
*W. P. Sch. op. cit. p.* 427.  
Br. rutilans *Brid. Bryol. univ. I, p.* 624 (1826); *C. M. Syn. I, p.* 274 (1849).  
2. In fiss. rup. humid. — *Eur.* : Spitzberg, ins. Parry, Urior, Finmark, alp. Norveg. *Am. sept.* : Fret. Scoresby, Smith, ins. Sabine.  
\* *var.* *gracilescens W. P. Sch. op. et l. cc.*  
*Ut typ.* — *Eur.* : Finmark, alp. Norveg.

*Ærstedtianum* C. M. — Webera.

*oligochloron* C. M. et Kindb. in *Bidr.* p. 454 (nom., 1890) et in *Mac. Cat.* p. 129 (1892).

Ster. Rupic. — *Am. sept.* : Columb. brit.

*oncophorum* Hpe. *Symb.* 1877, p. 726.

2? — *Am. merid.* : Brasil. austr. or.

*eophyllum* C. M. in *Hedwig.* 1898, p. 97.

2. Ad rivul. — *Pac.* : Nov. Seeland. sept.

*opdalense* *Limpr. Emige neue Art...* 1883, p. 238 et *Laubm. Deutschl.* II, p. 310 (1892).

Syn. In aren. humid. — *Eur.* : Norveg.

*oranicum* C. M. in *Hedwig.* 1899, p. 68.

Br. argenteum var. australe ej. in *Rehm. M. Afr. austr. n.* 260.

2. — *Afr.* : Orange.

*orbiculare pulvinatum, hirsutie* } *immersis* Dill. — *Grimmia pulvinata.*  
*canescens, capsulis* } *erectis praedita* ej. — *Grimmia leucophoea.*

*orbiculatum* Mitt. — *Anomobryum.*

*oreganum* Sull. in *M. Wilkes Exped.* p. 10, t. 7; *Lesq. et Jam. Man.* p. 230 (1884).

Syn. vel polyg. — *Am. sept.* : Oregon; Yellowstone Park?

*organum* Bosw. { — Br. pallens (*fid. Limpr.*).  
 { — Br. Weesleyi (1).

*orarium* J. O. Bomanss. in *Rev. bryol.* 1899, p. 9 et in *Act. Soc. pr. Faun. et Fl. fenn.* 1899, n. 4.

Syn. Ad litt. aren. marit. — *Eur.* : Ins. Aland.

*orthocarpulum* C. M. — Webera.

*orthocladon* Br. et W. P. Sch. in *W. Sch. M. Abyss. n.* 589; C. M. *Syn.* I, p. 318 (1849).

2. — *Afr.* : Abyss.

*orthodontioides* C. M. — Br. Gilliesii.

*orthodontum* P. B. — *Dissodon borbonicus.*

*orthopelma* C. M. — *Brachymenium.*

*orthophyllum* Besch. in *Ann. sc. nat. VII ser. II*, p. 90.

2. — *Afr.* : Ins. Mayotte.

*orthorhynchum* Brid. — *Mnium.*

*Orthotheciellae* C. M. — Br. Eatoni.

*Osculationum de Not. in Mem. Acad. Torin...* II ser., XVIII, p. 442.

— *Am. merid.* : Columb.

*ovale* Hffm. — } *Grimmia commutata* (*fid. C. M.*).  
 } *Grimmia ovata* (*fid. Limpr.*).

*ovalifolium* Sull. — Br. Crügeri.

*ovatum* Jur. *Laubmfl. v. Oesterr. Ung. p.* 291 (1882).

Br. neodamense var. ovatum *Lindb. et Arn. M. As. bor.* 1890, II, p. 34.

(1) Ob nimian similitudinem cum praeced. nom. mutatum.



- Br. pseudo-triquetrum*  $\delta$  *cavifolium* Saut. *Fl. d. Herzogth. Salz. III*, p. 47.
2. *Paludic.* — *Eur.* : *Salisb.*, *Stir. As.* : *Jenissei infer. Am. sept.* : *Fret. Scoresby.*
- ovatum* Dicks. — *Pterygoneuron cavifolium.*
- ovicarpum* Broth. in *Oefv. af Finska Vet.-Akad. Foerh.* 1899, p. 101.  
2. fr. ign. — *Pac.* : *Austral. or.* (Nov. Vales.).
- oxystegium* Hagen in *Tromsoë Mus. Aarsheft.* 1898-99, p. 211.  
*Syn.* In turf. — *Eur.* : *Norveg. bor.*
- Pabstianum* C. M. in *Bot. Zeit.* 1852, p. 751.  
*Br. Gardneri* Mitt. in *Journ. of bot.* 1856, p. 233, t. XI B et M. austr. am. p. 293 (1869).  
*Syn. Terr.* — *Am. merid.* : *Brasil. austr. or.*, *And. Peruv.*
- pachyloma* Card. in *Ann. Rev. of the Missouri bot. Gard.* 1897, p. 62.  
2? — *Afr.* : *Ins. Azor.*
- pachyneuron* Dusen M. *Am. austr. n.* 135.  
*Terr.* — *Am. merid.* : *Chile austr.*
- pachypoma* { *Mont.* { — *Br. plumosum.*  
                  { (*fid. Mitt.*). — *Br. nitens.*  
                  { *Wils. p.p.* { — *Br. alpinum.*  
                              { — *Br. caespitium.*  
                              { — *Br. erythrinum.*
- pachypomatoides* Hpe. mss.; Mitt. in *Trans. and Proced. of the roy. Soc. of Victoria*, 1882, p. 71.  
— *Pac.* : *Absq. loco (Austral. vel Tasman.).*
- pachypomatulum* Broth. in *Oefv. af Finska Vet.-Akad. Foerh.* 1899, p. 103.  
2. fr. ign. — *Pac.* : *Austral. or.* (Nov. Vales.).
- pachythea* C. M. *Syn. I*, p. 307 p.p. (1849).  
*Br. balanoides* Tayl. in *Drumm. M. Swan River (fid. Fl. of the N. Zeal. II*, p. 191).  
*Br. dichotomum* (*fid. Mitt.*).  
*Br. coronatum*  $\beta$  minus Hpe. in *Preiss Pl. N. Holl. n.* 117.  
2. — *Pac.* : *Austral. occid.*, *Nov. Seeland.*, *Tasman.*  
 $\alpha$  *var. inflatum* Hook. f. et Wils. *Fl. Tasman. II*, p. 191 (1860).  
— *Pac.* : *Austral.*, *Nov. Seeland.*
- pachythea* { C. M. pp. — *Br. pseudo-pachythea.*  
                  { Dz. et Mk. — *Br. Junghunianum.*
- pallens* Sw. M. *Suec. pp.* 47 et 98, t. 4, f. 12 (1799); *Brid. Bryol. univ. I*, p. 642 (1826); *Bryol. eur. fasc. 6-9 (Vol. IV)*, Mon. p. 68, t. 373 (1839); C. M. *Syn. I*, p. 256 (1849); W. P. Sch. *op. cit.* p. 456.  
*Br. annotinum lanceolatum pellucidum capsulis oblongis pendulis* Dill. *Hist. M.* p. 399, t. 50, f. 68 (1741) et *hb. fol.* 125, n. 68; t. 51, fig. 73 C-E *fid. Lindb.*  
*Br. nitidum*, foliis Serpilli angustioribus, minus Dill. *Cat. Giss.* p. 228 (1718).  
*Br. turbinatum H. et T. Musc. brit. p.* 122 p.p. (1818).  
*Br. turbinatum var. pallens* Drumm. M. *bor. am. I ed.* p. 261.  
*Hypnum pallens* W. et M. *Bot. Taschb.* p. 286 (1807).  
*Mnium gracile* Schleich. *Cent. II Pl. helv. n.* 46.

- Mn. pallens* P. B. *Prodr.* p. 47 (1805).  
*Pohlia pallens* Brid. *Musc. rec.* II, P. III, p. 145 (1803-4).
2. Ad scatur., terr. saxaq. humid. — *Eur.* : E reg. campestr. montos. usq. in alp. *As.* : Japon., Jenissei, Caucas. *Am. sept.* : Groenland., Gaspes., New Hampshire, M. Adirondack, Ontario, M. Rupest. et Selkirk, Wyoming, Oregon, Columb. brit., ins. Vancouver. *Am. merid.* : And. Quitens.
- α *var. abbreviatum* Bryol. eur. et W. P. Sch. II, cc.  
*Br. attenuatum* Dicks. *Pl. crypt. fasc.* IV, t. 10, f. 8 (1802); *Brid. Bryol. univ.* I, p. 638 (1826).  
*Orthopyxis attenuata* P. B. *op. cit.* p. 45.  
*Ut typ.* — *Eur.* : Reg. alp.
- α *var. angustifolium* Breidl. *Laubm. v. Steierm.* p. 134 (1891).  
*Br. triste* Jur. *Laubmfl. v. Oesterr. Ung.* p. 287 (1882).  
*Ad terr. et rap. calcar.* — *Eur.* : Austr. infer., Stir.
- α *var. arcuatum* W. P. Sch. *Syn.* I ed. p. 377 (1860).  
*Cynodontium cernuum* Schleich. *Cent. III Pl. helvet. n.* 9.  
*Ptychostomum cernuum* Hüb. *Muscol. germ.* 435 (1833).  
*In depressis.* — *Eur.* : Neerland., Helvet.  
*var. baldense* Vent. et Bott. — *Br. fallax var. baldense.*  
*var. brevisetum* Lindb. et Arn. — *Br. finmarkicum.*  
*var. fallax* Jur. — *Br. fallax.*  
*var. microstoma* Bryol. eur. — *Br. pallens var. speciosum.*
- α *var. œnodes* Limpr. in *LXI Jahresb. d. schles. Ges. f. Vaterl. Kult.* 1884.  
— *Eur.* : Alp. Norveg. (Dovre).
- α *subspec. rufescens* Kindb. *Eur. und N. Am. Bryin.* p. 352 (1897).  
*Ster.* — *Eur.* : Alp. Norveg.
- α *var. speciosum* (Voit) W. P. Sch. *Syn.* I ed. p. 377 (1860) et II ed. I, c.  
*Br. elegans* Grev. in *Trans. of the Linn. Soc.* XV, p. 344, t. 3, f. 5?  
*Br. pallescens var. microstoma* Bryol. eur. I, c. f. 8  
*Br. speciosum* Voit in *Sturm Deutsch. Fl. II, Heft.* 14 c. ic. (1813); *Brid. op. cit.* p. 666.  
*Ut typ.* — *Eur.* : Reg. alp.
- α *var. vulturienne* de Not. in *Piccone Enum. M. lig. n.* 101 (1863).  
— *Eur.* : Ligur.  
*var. Wils.* — *Br. calophyllum.*
- pallenticoma* C. M. in *Hedwig.* 1898, p. 100.  
*Br. Billardieri* Hpe. in *M. Preissianis.*  
*Rhodobryum pallenticoma* Par. *Suppl. Ind.* p. 300 (1900).  
2. — *Pac.* : W. Austral. (Swan River).
- pallescens* Schleich. *Cent. Pl. helvet. n.* 28; *Schwaegr. Suppl. I, P. II,* p. 107, t. 75 (1816); *Brid. Bryol. univ.* I, p. 645 (1826); *Bryol. eur. fasc.* 6-9 (Vol. IV), *Mon.* p. 51, t. 359 (1839); *C. M. Syn. I,* p. 279 (1849) et II, p. 570 (1851); *W. P. Sch. op. cit.* p. 433.  
*Br. rupicola* Schleich. *Cat.* 1821.  
*Br. turbinatum* Drumm. *M. bor. am. I ed. n.* 266 pp.

1. Ad mur. et rup. arenac. — *Eur.* : F. reg. campestr. usq. in alp.; Spitzberg. *As.* : Japon., Kamschatka, Amour, Jenissei, Himalaya (*var.*), Kashmir, Gilgit, Caucas. *Afr.* : Ins. Kerguelen. *Am. sept.* : Groenland., Labrador, Terra nov. ?, ins. Miquelon, New Brunswick, Ontario, M. Rupestr., Washington, Columb. brit., Utah, Colorado, Calif.; reg. sept.

var. *Baenitzii* Hagen. — Br. Baenitzii.

« var. boreale (*Schwaegr.*) *Bryol. eur. l. c. t.* 360; *W. P. Sch. op. cit.* p. 434.

Br. boreale *Schwaegr. Suppl. I, P. II, p. 96, t. 69* (1816); *Brid. op. cit.* p. 643.

Hypnum boreale *W. et M. Bot. Taschb. p. 287 p.p.* (1807) (*fid. Limpr.*).

Ut typ. — *Eur.* : Reg. alp.

« var. contextum (*H. et H.*) *Bryol. eur. et W. P. Sch. ll. cc.*

Br. contextum *H. et H. in Flora* 1819, *I, p. 91*; *Brid. l. c. p. 646.*

Ut typ. — *Eur.* : Spitzberg., Scandin., Alp. Helvet., Sabaud. *As.* : Jenissei. *Am. sept.* : Groenland.

« var. cylindrica *Hagen in Tromsø Mus. Aarsheft. 1898-99, p. 189.*  
Heteroic. — *Eur.* : Norveg. bor.

« var. flexisetum *Limpr. in LXI Jaresber. d. schles. Ges. f. Vaterl. Kult. 1884.*

— *Eur.* : Norveg.

var. *intermedium* *Schwaegr.* — Br. *intermedium.*

« var. laxifolium *Kindb. in Bull. Torr. bot. Club, XVII, p. 274* (1890).  
In silv. humid. — *Am. sept.* : M. Rupestr.

« var. longifolium *Kindb. in op. et l. cc.*  
In aren. humid. — *Am. sept.* : M. Rupestr.

var. *subrotundum* *Bryol. eur.* — Br. *subrotundum.*

« var. synoicum *Schiffn. Bryol. Durchforsch. d. südl. Theil v. Boehmen in Sitz. ber. natur. med. Ver. Lotos, 1898, n. 5, p. 33.*  
Muric. — *Eur.* : Bohem.

*pallescens* Dz. et Mk. — Br. *gedeanum.*

*pallidipes* C. M. in *Hedwig. 1897, p. 93.*  
— *Am. merid.* : La Plata.

*pallidisetum* *Brid.* — *Mniobryum carneum.*

*pallido-julaceum* C. M. — *Brachymenium.*

? *palmifolium* *Brid.* — *Conomitrium.*

*pallidum* *Schreb.* — *Ditrichum.*

*paludicola* *W. P. Sch. Syn. II, p. 415* (1876).

Br. pendulum var. *paludicola* N. Bryhn *De Bryin. i. Norvegia. p. 13* (1891)  
*Syn.* (1 in adn.). In uligin. — *Eur.* : Alp. Norveg.

*paludosum* L. — *Seligeria pusilla.*

β *viridulum* *Web.* — *Weisia viridula.*

- Brid. — Didymodon rubellus.  
Dicks. — Dicranella squarrosa.  
L. — Hymenostylium curvirostre.  
Sw. — Aulacomnium.  
*aestivum* Conservae facie Dill. — Eucladium verticillatum.
- palustre* { *brevifolium capsulis nigrican-* } — Barbula fallax var. brevi-  
              *tibus* Dill.                     *folia* (fd. C. M.).  
  — Didymodon tophaceus (fd.  
  *Limpr.*).  
    *complicatum, rubens, capsulis turbinatis pendulis* Dill. —  
        Br. turbinatum.  
    *pellucidum, capsulis et foliis brevibus recurvis* Dill. — Dichod-  
        ontium pellucidum.  
    *pilosum verticillatum* Dill. — Eucladium verticillatum.  
    *scopis teretibus stellatis, vertice multifidis* Dill. — Philonotis  
        fontana.  
var. *ramosum* Turn. — Aulacomnium palustre var. polycephalum.
- Pancheri Jaeg. Ad. I, p. 639 (1873-74).  
Br. crassinervium Besch. Fl. bryol. Nouv. Calédon. p. 215 (1873).
2. Ad terr. ferrug. — Pac. : Nov. Calédon.
- papillosum* { C. M. — Brachymenium.  
              } Dicks. — Ceratodon purpurens.
- Pappeanum C. M. in Bot. Zeit. 1855, p. 752 et 1859, p. 206.  
1. — Afr. : Prom. B. Sp.
- « var. humile C. M. in op. cit. p. 206.  
Ubi typ.
- paradoxum Schwaegr. Suppl. III, P. I, I, t. 224 (1823); C. M. Syn. I,  
p. 290 (1849); Mitt. M. Ind. or. p. 74 (1859).  
1. — As. : Nepal.
- paradoxum* { Jur. — Br. cuspidatum.  
              } Philib. — Br. terskeiense.  
var. *cirratum* Jur. — Br. cuspidatum var. cirratum.
- paraguense Besch. Not. M. Parag. p. 264 (1877).  
2. — Am. merid. : Paraguay.
- parasiticum Sw. — Calymperes.
- parvirete Dusen M. Am. austr. n. 684.  
Terr. — Am. merid. : Chile austr.
- parvulum Hoffm. — Dicranella cerviculata var. pusilla.  
    *erectis subrotundis majusculis capitulis subfuscis, foliis*  
    *Serpilli pellucidis* Dill. — Pottia intermedia.
- parvum* { *surculis et setis geminatis* ej.  
          *trichodes ramosum, erectis capitulis,*  
          *subfuscis, in pediculis obscure* } — Ceratodon purpu-  
          *rubris* ej. reus.
- patens* { Dicks. — Rhacomitrium.  
          } Wils. — Br. rigidum.
- patulum C. M. — Brachymenium.
- paucifolium Hpe. in Ann. sc. nat. V ser. IV, p. 344 (1866).  
— Am. merid. : Nov. Granat.

*paucifolium* { Dicks. } — *Seligeria subcernua* (fd. Lindb.).  
                   { Mitt. — *Gyroweisia tenuis* (fd. W. P. Sch.).  
*pauperum* C. M. — Webera.

*Payoti* W. P. Sch. *Syn. II* ed. p. 470 (1876).

Br. *Blindii* var. *Payoti* Husn. *Musc. gall.* p. 243 (1889).

Ster. Secus rivul. — *Eur.* : Alp. Sabaud. (m. Blanc.)

*pellucens* Hook. — *Orthodontium*.

*pellucidum* { Abb. — *Tetraphis*.  
                   { L. — *Dichodontium*.  
       β L. — *Dicranella squarrosa*.

*pendulinum* Hpe. in Linn. 1874, p. 214.

— *Afr.* : Madagascar.

*pendulum* (Hsch.) W. P. Sch. *Coroll.* p. 70 (1856) et *Syn. II* ed. p. 414 (1876).

Br. caespitium *Drumm. M. bor. am. I* ed. n. 265 p.p.

Br. cernuum *Bryol. eur. fasc.* 6-9 (*Vol. IV*), *Mon.* p. 14, t. 331 (1839);  
*C. M. Syn. I*, p. 266 (1849).

Br. pendulum ovatum, caespitium et pilosum, seta bicolori var. *H. Dill. Hist. M. p.* 397, t. 50, f. 66 H (1741).

*Cynodon cernuus* *Brid. Mant. M. p.* 99 (1819) p.p.

*Cynodontium cernuum* *Schwaegr. Suppl. I, P. I*, p. 110 (1811).

*Cynodontium cernuum* *Hedw. Sp. M. p.* 58, t. 9 (1801).

*Didymodon cernuus* *Sw. M. Suec. p.* 29, t. I, f. 9 (1798) fd. *Brid.*

*Ptychostomum cernuum* *Hsch. in Fl.* 1882, *Syll. I*, p. 64; *Brid. Bryol. univ. I*, p. 600 (1826).

*Pt. pendulum* *Hsch. in op. et l. cc.*; *Brid. op. cit.* p. 599.

*Pt. radiculosum* *Brid. op. cit.* p. 838.

*Pt. radiculosum* var. *spadiceum* *ej. op. cit.* p. 837?

*Swartzia cernua* *Poir.*

*Sw. radiculosa* *Hedw. in hb. DC.*

*Syn.* Ad mur. saxa et terr. — *Eur.* : E reg. campestr. usq. in alp.

*As.* : Japon., Saghalien, Amour, Pamir, Himalaya, Panjab, Caucas.

*Afr.* : Alger. ? *Am. sept.* : Fret. Smith, Groenland, Labrador,

Anticosti, Terra nov., Canad. merid., Wyoming, M. Rupestr.,

Columb. brit., fret. Behring, temperata tota.

« var. *angustatum* *Ren. in Husn. Musc. gall.* p. 235 (1889) et *M. Gall.* n. 667.

Ad mur. et terr. — *Eur.* : Boruss. or., Pyren.

« var. *compactum* (Hsch.) *Hartm. Skand. Fl. IV* ed. p. 407 (1843);  
*W. P. Sch. Coroll. l. c.*

Br. algovicum *Sendtn. mss.*; *C. M. Syn. II*, p. 569 (1851).

Br. caespitium *Brid. Bryol. univ. I*, p. 837 (1826).

Br. compactum *Kindb. Eur. a. N. Amer. Bryin. p.* 380 (1897).

*Ptychostomum compactum* *Hsch. in Fl. Syll. I*, p. 62 (1824).

Rupic. — *Eur.* : Summ. Alp. *Am. sept.* : Fret. Scoresby.

« var. *microcarpum* *Warnst. in Abh. d. bot. Ver. d. Prov. Brandenb.* 1900, p. 203.

— *Eur.* : Boruss. or.

var. *paludicola* *Bryhn.* — Br. *paludicola*.

- Brid. — *Breutelia*.  
 Ehrh. — *Leptobryum piriforme*.  
 Fl. dan. — *Bartramia Halleriana*.  
 Mitt. (1876). — *Br. macrautherum*.  
*foliis variis pellucidis, thecis ovatis* Dill. — *Mnium affine et cuspidatum*.  
*hornum molle, foliis et lanceolatis et gramineis* Dill. — *Webera cruda*.  
*julaceum argenteum et sericeum* Dill. — *Br. argenteum*.  
*pendulum* { *ovatum caespitium et pilosum, seta bicolori* Dill. } — *Br. atro-purpureum* (fid. Limpr.).  
 { — *Br. caespitium* (fid. C. M.).  
*ovatum caespitium et pilosum, seta bicolori* var.  $\epsilon$  ej. — *Br. erythrocarpum*.  
*Serpilli folio* { *longiore pellucido, thecis oblongis cuspidatis* Dill. — *Mnium cuspidatum*.  
 { *rotundiore pellucido, thecis ovatis* Dill. — *Mnium punctatum*.  
*surculis teretibus viridibus* Dill. — *Br. atro-purpureum*.  
*penicillatum* Hpe. in Linn. 1874, p. 215.  
 — Afr. : Madagascar.  
*pennatum* With. — *Schistostega osmundacea*.  
*pensilvanicum* Brid. — *Mnium serratum* var. *apiculatum*?  
*perangustis foliis et cauliculis, foliis* { *crebrioribus et circa summitates magis congestis capitulis erectis et* } *ad summitates magis egredientibus* Dill. — *Didymodon rigidulus*.  
 { *rarioribus cinctis, capitulis erectis et surculis annotinis egredientibus ej.* — *Ceratodon purpureus*.  
*perappressum* C. M. — *Anomobryum*.  
*peraristatum* C. M. in Hedwig. 1898, p. 106.  
*Rhodobryum peraristatum* Par. Suppl. Ind. p. 300 (1900).  
 2. — Pac. : Austral. or. (Victoria).  
*peraristatum* C. M. (Venez.) — *Brachymenium*.  
*percurrentinerve* Kindb. in Bull. Torr. bot. Club, XVII, p. 274 (1890)  
 Ster. Ad rup. irror. — Am. sept. : Columb. brit.  
*perdecurrens* C. M. — *Webera*.  
*pergracilescens* C. M. in Bull. hb. Boissier, 1897, p. 184.  
 Ster. — Am. sept. : Guatemala.  
*perimbricatum* C. M. — *Anomobryum*.  
*perminutum* C. M. in Bull. hb. Boissier, 1897, p. 182.  
 Ster. — Am. sept. : Guatemala.  
*Peromnium* C. M. — *Brachymenium radiculosum*.  
*perspinidens* Broth. in Engler's Bot. Jahrb. 1897, p. 246.  
 2. — Afr. : Reg. lacustr.

*pertenne* Besch. in *Journ. de bot.* 1894, p. 60.

2. Terr. — *Am. sept.* : Guadalup.

*pertomentosum* Dusen *M. Am. austr. n.* 601.

Rupic. — *Am. merid.* : Patagon. occid.

*pertriste* C. M. in *Linn.* 1878-79, p. 297 (*sub nom.* *Br. tristis*, *post. correct.*).

Calcic. ? — *Am. merid.* : Argentin. subtropic.

*pervirens* C. M. in *Rab. Bryoth. eur. n.* 1309.

— *Afr.* : Prom. B. Sp.

*Pfefferi* de Not. *Epil.* p. 409 (1869).

Ad rup. granit. — *Eur.* : Ital. super. (Vall. Tellin.).

*Phallus* C. M. in *Linn.* 1875, p. 392.

*Br. coronatum* ej. *Syn. I*, p. 307 p.p. (1849); *Mitt. M. austr. am.* p. 321 (1869).

2. — *Am. merid.* : Chile.

*phascoides* Jaeg. — *Diphyscium foliosum*.

*Philiberti* Amann. — *Br. comense*.

*Philippianum* C. M. in *Linn.* 1844, p. 701 et *Syn. I*, p. 772 (1849).

*Syn.* — *Am. merid.* : Chile.

*philonotum* C. M. — *Mniodendron*.

*Philonotula* } C. M. — *Webera*.

} Hpe. — *Brachymenium*.

*phyllitidifolium* Neck. — *Atrichum undulatum*.

*Phyllitidis folio rugoso acuto, capsulis incurvis* Dill. — *Atrichum undulatum*.

*piliferum* Br. et W. P. Sch. in W. Sch. *M. Abyss.*

— *Afr.* : Abyss.

*piliferum* Dicks. } — *Desmatodon latifolius* var. *brevicaulis* (fid. C. M.).

} — *Pottia latifolia* var. *pilifera* (fid. W. P. Sch.).

*pilosum* } *molle setis intortis* Dill. } — *Campylopus flexuosus* (fid. C. M.).

W. P. Sch.).

} *Sphagni subulati facie* Dill. — *Blindia acuta*.

*verticillatum* Dill. — *Eucladium*.

*Pimpanae* C. M. in *Hedwig.* 1898, p. 94.

2. *Pac.* : Austral. or. (Queensland).

*piriforme* } Wegg. — *Leptobryum*.

} L. — *Physcomitrium*.

var. Weis. — *Webera nutans*.

*plagiopodium* W. P. Sch. in C. M. *Syn. II*, p. 572 (1851).

2. — *Am. sept.* : Mexic.

*planifolium* Kindb. *Art. d. Laubm. Schwed. u. Norweg.* p. 75 (1883).

*Syn.* — *Eur.* : Norveg. (Dovre).

*planiusculum* Lindb. et Arn. *M. As. bor.* 1890, II, p. 38.

2. Terr. — *As.* : *Jenissei* med.

*platyloma* } Bryol. eur. — *Br. Donianum*.

} Schwaegr. — *Br. capillare* var. *platyloma*.

*platyphylloides* C. M. in *Linn.* 1878-79, p. 298.

2. Limic. — *Am. merid.* : Argentin. Cordob.

*platyphyllum* (Schwaegr.) C. M. *Syn. I*, p. 291 (1849).

*Pohlia platyphylla* Schwaegr. *Suppl. IV*, p. 324 (1842).

2. *Rupic.* — *Am. merid.* : And. Chil. austr.

*plebejum* C. M. in *Hedwig*. 1898, p. 94.

2. — *Pac.* : Austral. or. (Victoria).

*plumaefolium* C. M. — Webera.

*Plumella* C. M. — *Brachymenium*.

*plumosum* Dz. et Mk. *M. frond. ined. Archip. ind.* p. 16, t. 7 (1854);  
*Bryol. jav. I*, p. 143, t. 117 (1855-61).

*Br. ambiguum* Duby in *Moritz Syst. Verz.* p. 132; C. M. *Syn. I*, p. 294 (1849).

*Br. apiculatum* Dz. et Mk. *op. cit.* p. 24, t. 8.

*Br. Leveilleum eor. op. cit.* p. 16.

*Br. pachypoma* Mont. in *Ann. sc. nat.* 1845, *IV*, p. 104; C. M. *op. cit.* p. 299.

*Br. plumosum* γ *Leveilleum Bryol. jav. l. c.*

2. *Terr.* — *As.* : Tonkin., Java, Sumatra.

♣ *var. elongatum* *Bryol. jav. l. c. t.* 144.

Ut typ. — *As.* : Java.

*var. Leveilleum* *Bryol. jav.* — *Br. plumosum*.

♣ *var. majus* Hpe. in *Bryol. jav. l. c.*

— *As.* : Java.

*pluricolor* Dusen *M. Am. austr. n.* 181.

*Muric.* — *Am. merid.* : Chile austr.

*pohliaeforme* W. P. Sch. in *Besch. Prodr. bryol. mexic.* p. 54 (1871).

2. — *Am. sept.* : Mexic.

*pohliaeforme* Brid. — *Br. subrotundum*.

*pohliaeopsis* C. M. in *Hedwig*. 1898, p. 107.

*Rhodobryum pohliaeopsis* Par. *Suppl. Ind.* p. 300 (1900).

2. — *Pac.* : Austral. or. (Victoria).

*polare* Hagen in *Saertryk of Meddel. om Groenland*, 1898, p. 98.

*Syn. Ad terr. humos. arenac.* — *Eur.* : Finmark. *Am. sept.* : Fret. Scoresby.

*polycarpon* Hfm. — *Cynodontium*.

*polycarpum* Mitt. — Webera.

*polychaete* C. M. — *Brachymenium*.

*polymorphoides* Kindb. — Webera.

*Bryol. eur.* — Webera.

*polymorphum* { C. M. { — Webera *acuminata* var. minor.

{ — Webera *acuminata* var. *polyseta*.

Neck. — *Mnium affine*.

Varr. omn. *Bryol. eur.* — *Ad Weberam polymorphum spectant.*

*polyphyllum* Dicks. — *Ptychomitrium*.

*polysetulum* C. M. — Webera.

*polytrichoides palustre, setis longioribus rubris sericeis* Dill. — *Ceratodon purpureus*.

*polytrichoides* C. M. in *Bot. Zeit.* 1859, p. 206.

2. *Rupic.* — *Afr.* : Prom. B. Sp.

*pomiforme* L. — *Bartramia*.



- var. *Halleri* Vill. — *Bartramia Halleriana*.  
 var. *majus* Web. — *Bartramia ithyphylla*.  
 var. *Æderi* Vill. — *Bartramia Æderi*.  
 var. Scop. — *Bartramia Halleriana*.
- Pomoniae* C. M. in *Linn.* 1876, p. 244.  
 2. Terr. — *Afr.* : Usambara, ins. Anjouan.
- pootenianum* Par. *Ind.* p. 206 (1894).  
 Br. *arenarium* C. M. in *Bot. Centralbl.* 1883, n. 2-4.  
 2. vel syn. — *As.* : Penins. Tchutschica.
- porphyroloma* C. M. in *Rehm. M. Afr. austr.* n. 250.  
 — *Afr.* : Prom. B. Sp.
- porphyroneuron* C. M. in *Bot. Zeit.* 1853, p. 22; *Mitt. M. Ind. or.* p. 70 (1859).  
 2. — *As.* : M. Neilgherr., Java, Sumatra.
- porphyreothrix* C. M. in *Rehm. M. Afr. austr.* n. 103 et in *Hedwig.* 1899, p. 70.  
 1. — *Afr.* : Prom. B. Sp.
- posthumum* C. M. in *Hedwig.* 1897, p. 94.  
 2. — *Am. merid.* : Uruguay.
- pottiaefolium* C. M. in *Linn.* 1875, p. 382.  
 Syn. Terr. — *As.* : M. orient. mar. Erythraei.
- praecox* Warnst. in *Verh. d. bot. Ver. d. Prov. Brandenb.* 1899, p. 58.  
 — *Eur.* : Lusat. infor.
- praemorso-obtusum* C. M. in *Dusen M. Cam.* n. 321.  
 — *Afr.* : Cameroon.
- Preissianum* Hpe. — *Brachymenium*.
- Preussii* Broth. in *Engler's Bot. Jahrb.* 1894, p. 187.  
*Rhodobryum Preussii* Par. *Ind.* p. 1119 (1897).  
 2. — *Afr.* : Cameroon.
- procerrimum* Par. — *Anomobryum*.
- procerum* W. P. Sch. in *Besch. Prodr. bryol. mexic.* p. 55 (1871).  
*Rhodobryum procerum* Par. *op. et l. co.*  
 — *Am. sept.* : Mexic.
- producticolle* Kindb. *Eur. and N. Amer. Bryin.* p. 364 (1897).  
 Rupic. — *Am. sept.* : Ins. Vancouver.
- proliferum* Mitt. *M. austr. am.* p. 311 (1869).  
 2. — *Am. merid.* : And. Nov. Granat.
- proligerum* Kindb. — *Webera*.
- Promontorii* C. M. in *Hedwig.* 1899, p. 69.  
*Mielichhoferia procumbens* *Rehm. M. Afr. austr.* n. 214.  
 2. — *Afr.* : Transvaal.
- proprium* Hagen in *Tromsoë Mus. Aarsheft.* 1898-99, p. 135.  
 Syn. Ad litt. marit. — *Eur.* : Norveg. bor.
- prostratum* C. M. — *Anomobryum*.  
 var. *minus* Par. — *Anomobryum prostratum* var. *minus*.

provinciale *Philib. in W. P. Sch. Syn. II ed. p. 432* (1876).

Br. Billardieri *Bryol. eur. fasc. 6-9 (Vol. IV). Mon. p. 58, t. 336* (1839).

Br. canariense *W. P. Sch. Syn. I ed. p. 574* (1860).

Br. canariense var. provinciale *Husn. Musc. gall. p. 239* (1889).

Syn. In pinet., suberet. et ad tumul. formic. — *Eur.* : Angl., Galloprov., Istr., Dalmat., Etrur., Roma. *Afr.* : Alger. *Am. sept.* : Florid., Calif.

*provinciale* Mac. — Br. heteroneuron.

pseudo-Acidodontium *C. M. in Linn. 1880-83, p. 364.*

2. — *Am. merid.* : Argentin. subtropic.

pseudo-acuminatum *Kindb. — Webera polymorpha var. pseudo-acuminata.*

pseudo-alpinum *Ren. et Card. in Bull. hb. Boissier, 1895, I, p. 62.*

Br. emodi-alpinum *C. M. mss. (fd. Card.).*

2. — *As.* : N. W. Himalaya, Sikkim.

pseudo-argenteum *Warnst. in Verh. d. bot. Ver. d. Prov. Brandenb. 1899, p. 62.*

— *Eur.* : Lusat. infer.

pseudo-capillare *Besch. Fl. bryol. Antill. fr. p. 31* (1876).

2. — *Am. sept.* : Martinic.

pseudo-decursivum *Par. Suppl. Ind. p. 70* (1900).

Br. decursivum *C. M. in Hedwig. 1899, p. 70.*

Ster. — *Afr.* : Prom. B. Sp.

pseudo-Funckii *Anzi Enum. M. Longob. p. 20.*

— *Eur.* : Ital. super. (Vall. Tellin.).

pseudo-Kunzei *Limpr. Laubm. Deutschl. ... p. 359* (1892).

Syn. — *Eur.* : Helvet. (Vales.).

pseudo-marginatum *Geh. et Bpe. in Fl. 1881.*

Rhodobryum pseudo-marginatum *Par. Ind. p. 1119* (1897).

2. — *Am. merid.* : Brasil. austr. or.

pseudo-micron *C. M. in Linn. 1880-83, p. 373.*

Ster. — *Am. merid.* : Argentin. temperat.

pseudo-pachytheca *C. M. in Bot. Zeit. 1859, p. 207.*

Br. pachytheca *ej. Syn. I, p. 307 p.p.*

2. — *As.* : India.

pseudo-pallescens *Hpe. et C. M. mss.; Mitt. in Trans. and Proc. of the roy. Soc. of Victoria, 1882, p. 72.*

— *Pac.* : Absq. loc. (Austral. vel Tasman.).

pseudo-Philonotula *C. M. — Webera.*

pseudo-selerodictyon *C. M. in Bal. M. Paraguay. n. 3647.*

— *Am. merid.* : Paraguay.

pseudo-torquescens *C. M. in Bot. Zeit. 1858, p. 155.*

Br. torquescens *W. P. Sch. in W. Sch. M. Abyss. II, n. 30.*

Syn. — *Afr.* : Abyss.

pseudo-triquetrum (*Hedw. p.p.*) *Schwaegr. Suppl. I, P. II, p. 110 excl. syn.* Br. bimum (1816); *Brid. Bryol. univ. I, p. 676 p.p.* (1826); *Bryol. eur. fasc. 6-9 (Vol. IV), Mon. p. 54, t. 64* (1839); *C. M. Syn. I, p. 258* (1849); *W. P. Sch. op. cit. p. 459.*

Br. annotinum palustre, capsulis ventricosus pendulis *Dill. Hist. M. p. 404, t. 51, f. 72* (1741) *et hb.*

Br. bimo affine *Wils. in Kew Journ. bot. IX, p. 366* (1857).

Br. cubitale *Dicks. Pl. crypt. fasc. II, p. 9, t. 5, f. 2* (1790).

Br. lanceolatum bimum, setis et capsulis longis pendulis *Dill. op. cit. p. 405, t. 51, f. 73 A et B et hb.*

Br. nitidum, foliis Serpilli angustioribus, majus *Dill. Cat. Giss. p. 228* (1718).

Br. subobliquum *Lindgr. in Bot. Not. 1842, p. 18.*

Br. ventricosum *Dicks. op. cit. fasc. I, p. 4* (1785); *Lindb. in Oefv. af K. Sv. Vet.-Akad. Foerh. p. 544* (1866).

Hypnum palustre erectum breve, foliis brevibus, tenuibus, angustis, minus confertis *Dill. Cat. Giss. p. 220* (1718).

H. pseudo-triquetrum *W. et M. Bot. Taschb. p. 288* (1807).

Mnium pseudo-triquetrum *Hedw. Descr. III, p. 19, t. 7* (1792) *p.p.*

? Mn. rubiginosum *Lam. Fl. fr. I, p. 41* (1778).

2. Ad terr. humid., in uligin. et ad rup. madid. — *Eur.* : Spitzberg., ins. Ursor.; e reg. campestr. usq. in alp. *As.* : Japon., China or., Jenissei, Obi, Tibet. occid., Kashmir, Caucas. *Afr.* : Alger., Madeir., *Am. sept.* : Groenland., Anticosti, Terra nov., ins. Miquel., Canada merid. or., Minnesota, Montana, Wyoming, M. Rupestr., Washington, Columb. brit., ins. Vancouver, tota temper.

♣ *var. arctica* *Ryan in Hagen M. Norveg. bor. 1898-99, p. 205.*  
In alv. exsicc. fluv. — *Eur.* : Norveg. bor.

♣ *var. anguste limbatum* *Warnst. in Verh. d. bot. Ver. d. Prov. Brandenb. 1899, p. 62.*

— *Eur.* : Lusat. infer.

♣ *var. angustifolium* *Lindb. in Krit. gransk. af Mossorna, p. 34* (1883); *Dill. Hist. M. p. 404, t. 51, fig. 71 A-C.*

— *Eur.* : ?

♣ *f. aulacodon* *Hagen in Tromsø Mus. Aarsheft 1898-99, p. 205.*

— *Eur.* : Finmark.

*var. bimum* *Brid.* — Br. bimum.

♣ *var. Bornmülleri* *Schiffn. in Oesterr. Zeitschr. 1897, p. 130.*

— *As.* : Persia or.

*var. cavifolium* { *Berggr.* — Br. crispulum.  
                          *Saut.* — Br. ovatum.  
                          *W. P. Sch.* — Br. neodamense.

♣ *var. compactum* *Bryol. eur. et W. P. Sch. II, cc.*

Ad rup. marit. — *Eur.* : Anglia, Norveg. *As.* : Jenissei.

*var. compactum* *Berggr.* — Br. crispulum.

♣ *var. duvalioides* *Itzigs. in litt.; Rab. Deutsch. Kryptfl. I ed., II, p. 214* (1848).

Br. duvalioides *Itzigs. in sched.*

Br. pseudo-triquetrum *var. flaccidum* *W. P. Sch. Coroll. p. 76* (1856) *et op. cit.*

In uligin. aquos. — *Eur.* : Neomark.

*var. flaccidum* *W. P. Sch.* — Br. pseudo-triquetrum *var. duvalioides.*

- « *var. gracilescens* W. P. Sch. *Coroll. et Syn. II ed. II. cc.*; *Dill. Hist. M. p. 404, t. 51, f. 73 A-B.*  
 Ut praeced. — *Eur.*: Suecia, Rhaet., etc. *As.*: Jenissei. *Am. sept.*: Wisconsin, Washington.
- « *var. hyalodontium* C. M. et Kindb. in *Mac. Cat. p. 133* (1892).  
 Ad rup. humid. — *Am. sept.*: M. Rupestr.
- « *var. latifolium* Lindb. in *op. et l. cc.*; *Dill. Hist. M. p. 406 var. colore rubro et p. 404, t. 51, f. 72 D-F.*  
 — *Eur.*: ?
- « *var. neo-inarchicum* Warnst. in *Verh. d. bot. Ver. d. Prov. Brandenb. 1900, p. 206.*  
 Inter al. musc. semel lectum. — *Eur.*: Brandeb.
- « *var. polytrichoides* Corbière in *Husn. Musc. gall. p. 250* (1889).  
 Marit. — *Eur.*: Normannia.
- var. rotundifolium* Zett. — Br. crispulum?
- « *var. synoicum* Arn. in *Oefv. af K. Sv. Vet.-Akad. Foerh. 1900, p. 116.*  
 Syn. Ad terr. aren. — *Eur.*: Spitzberg.
- ptychothecioides* C. M. — *Rhodobryum roseum.*
- ptychothecium* Besch. in *Ann. sc. nat. 1892, I, p. 66.*  
*Rhodobryum ptychothecium* Par. *op. et l. cc.*
2. — *As.*: Yunnan.
- Puiggarii* Geh. et Hpe. in *Fl. 1881, p. 376.*  
 1. — *Am. merid.*: Brasil. austr. or.
- pulchellum* { *Bryol. eur.*, W. P. Sch. *Coroll. et Syn. I ed. p.p.* — Webera.  
 W. P. Sch. *Syn. II ed. p.p.* — *Mniobryum vexans.*  
 Hedw. — Webera.  
 Schultz. — Br. erythrocarpum (*fid. Fiedl.*).  
 Sull. — Webera Lescuriana.
- var. atro-purpureum* Wahlenb. — *Mniobryum.*
- pulchrum* C. M. — *Brachymerium.*
- pulvinatum* { C. M. — Webera.  
 L. — *Grimmia.*
- var. β* Poll. — *Pottia lanceolata.*
- pumilum* Ryan in *Hagen M. Norveg. bor. 1898-99, p. 191.*  
 Heteroic. In fiss. rup. dolomit. — *Eur.*: Norveg. bor.
- pumilum* { Saut. — Br. Klinggraeffii.  
 Wils. p.p. — Br. subrotundum.
- punctatum* { Drumm. — *Mnium subglobosum.*  
 Schreb. — *Mnium.*
- pungens* Tayl. in *Lond. Journ. of bot. 1846, p. 342*; C. M. *Syn. I, p. 302* (1849); Mitt. *M. austr. am. p. 304* (1869).  
 — *Am. merid.*: M. Pichincha.
- pungentifolium* C. M. in *Hedwig. 1898, p. 97.*  
 2. — *Pac.*: Nov. Seeland. sept. occid.
- purpurascens* (R. Br.) *Bryol. eur. fasc. 32 (Vol. IV), Mon. Suppl. p. 2, t. 336* (1846); C. M. *Syn. I, p. 270* (1849) et *II, p. 569* (1851); W. P. Sch. *op. cit. p. 410.*  
 Br. fuscum Ferg. in *Natural. N. S. V. p. 82* (1880).

- Hemisynapium arcticum*  $\beta$  *purpurascens* Brid. *Bryol. univ.* I, p. 606 (1826).  
*Pohlia arctica*  $\beta$  *purpurascens* Schwaegr. *Sp. M.* p. 74 (1830).  
*P. purpurascens* R. Br. *Suppl. App. ad iter Parray.* p. 197 (1824).  
 Syn. In humid. et spongios. — *Eur.* : Lapon., alp. Norveg. *As.* :  
 Jenissei, Alatau. *Am. sept.* : Ins. Melville, Groenland., Labrador,  
 fret. Smith., M. Rupestr., Columb. brit.
- purpuratum* C. M. in Linn. 1880-83, p. 370.  
 2? — *Am. merid.* : Argentina. subtropic.
- purpureo-nigrum* Duby in *Mém...* Genève, 1875, P. II, t. I, f. 3.  
 2. — *Afr.* : Madagascar.
- purpureum Philib.* in *Rev. bryol.* 1886, p. 24.  
 2. — *Eur.* : Norveg.
- purpureum* Huds. — *Ceratodon*.
- pusillum* Broth. in *Oefv. af Finska Vet.-Soc. Föerh.* 1890, p. 99.  
*Rhodobryum pusillum* Par. *op. et l. cc.*  
 2. Cortic. — *Pac.* : Austral. or. (Queensland).
- pusillum* { Gmel. — *Ditrichum tortile* var. *pusillum*.  
 Hedw. — *Pterygoneuron cavifolium*.  
 Hoffm. — *Seligeria*.
- pycnobasium* C. M. — *Anomobryum*.
- pyncoderium* Limpr. in *LXI Jahresb. d. schles. Ges. für Vaterl. Kult.*  
 1884, p. 217.  
 1. In arena. — *Eur.* : Norveg. (Dovre).
- pyncoloma* C. M. in *Rehm. M. Afr. austr. n.* 239.  
 — *Afr.* : Austr.
- pygmaeo-alpinum* C. M. et Kindb. in *Bidr.* p. 454 (nom., 1890) et in *Mac.*  
*Cat. p.* 126 (1892).  
 Ster. Ad rup. humid. — *Am. sept.* : M. ins. Vancouver.
- pygmaeo-mucronatum* Philib. in *Rev. bryol.* 1899, p. 34.  
 Syn. — *As.* : M. Alatau.
- pygmaeum* C. M. in *Engler's Bot. Jahrb.* 1883, p. 77 et in *Forschungs-*  
*reise...* Bot. p. 12 (1889).  
 Ster. Ad rup. sicc. — *Afr.* : Ins. Kerguelen.
- pyrothrix* C. M. in Linn. 1880-83, p. 367.  
 Ster. — *Am. merid.* : Uruguay.
- quitense* Par. — *Anomobryum*.
- radiale* C. M. — *Brachymenium*.  
 var. *leptoradiale* C. M. — *Brachymenium radiale* var. *leptoradiale*.
- radians* P. B. — *Entosthodon*.
- radicale* Rehm. *M. Afr. austr. n.*  
 — *Afr.* : Orange.
- radiculosum* (Schwaegr.) Mitt. *M. austr. am.* p. 284 (1869).  
 Br. *Peromnium C. M. Syn.* I, p. 321 (1849).  
*Peromnium radiculosum* Schwaegr. *Suppl. III, P. I, t.* 250 (1828).  
 2. Cortic. — *Am. merid.* : Brasil. austr. or.
- radiculosum* Brid. — Br. *erythocarpum* var. *ramosum*.
- ramicola* Spr. — *Acidodontium*.

*ramosum* (Hook.) Mitt. *M. Ind. or.* p. 75 (1859).

Br. neelgheriense Mont. (fide Mitt.).

Br. truncorum Wils. in *Kew Journ. bot.* IX, p. 366 (1857).

*Mnium ramosum* Hook. in *Lond. Journ. of bot.* 1840, II, p. II et Io. pl. var. t. 20, f. 2 (1841).

Mn. rostratum var. *ramosum* C. M. *Syn.* I, p. 558 (1849); *Par. Ind.* p. 834 (1896).

*Syn.* — As. : Nepal; m. Neilgherr.?, Ceylan, Java.

*Rau* Aust. — Br. *Mühlenbeckii*.

*reclinatum, foliis fulcatis scoparium* { — *Dicranum majus*,  
*effigie* Dill. { — *Dicranum scoparium*.

*recurvatum* Hoffm. — *Seligeria*.

*recurvifolium* Tayl. in *Trans. of the Edinb. Linn. Soc.* 1845, II, 1; C.

*M. Syn.* I, p. 340 (1849).

*Ster.* — Eur. : Hibern.

*recurvirostrum* { Dicks. — *Didymodon rubellus*.  
                              { Gmel. — *Hymenostylium curvirostre*.

*recurvulum* Mitt. *M. Ind. or.* p. 74 (1859).

2. — As. : N. W. Himalaya, Nepal.

*recurvum* Wulf. — *Bartramia Halleriana*.

*Regnellii* C. M. — *Brachymenium*.

*Rehmanni* C. M. in *Rehm. M. Afr. austr. n.* 234.

— Afr. : Prom. B. Sp.

*Remelii* C. M. in *Fl.* 1874, n. 31.

*Ster.* Muric. — Afr. : Ægypt. super.

*remotifolium* Hpe. in *Linn.* 1863, p. 131.

2. Terr. — Am. merid. : And. Nov. Granat.

*Renauldi* Roell in sched.; *Ren. et Card.* in *Bull. Soc. roy. bot. Belg.* 1899, II, p. 221.

*Ster.* — Am. sept. : Costa Rica.

*repens* Gmel. — *Amblystegium serpens*.

*restitutum* de Not. *Epil.* p. 408 (1809); *W. P. Sch. op. cit.* p. 469.

Br. capillare var. *restitutum* Kindb. in *Boll. d. Soc. bot. ital.* 1896, p. 17.

*Desmatodon* ? de Not. *Syll. n.* 275 (1838).

*Ster.* In pascuis. — Eur. : Ital. super. (agro Comens.).

*reticulatum* Dicks. — *Dissodon Froehlichianus*.

*retusum* Hagen in *Det. K. Norske Vid. sells. Skript.* 1897, n. 2.

*Syn.* — Eur. : Norveg. bor.

*revelatokense* Kindb. in *Rev. bryol.* 1896, p. 22.

2. Ad rup. madid. — Am. sept. : Columb. brit.

*revolutum* C. M. in *Linn.* 1878-79, p. 291.

2. Ad rup. gneiss. — Am. merid. : Argentin. Tucuman.

*Reyeri* Breidl. in *Rev. bryol.* 1888, p. 35 et in *Verh. d. K. K. zool. bot. Ges. in Wien*, 1888, p. 799.

2. ♂ et fr. ign. — Eur. : Pusterthal, Stiria.

*rhaeticum* { Rota. — Webera.

                              { Amann. — Br. Killiasii.

*rhamphostegium* (Hpe). — *Acidodontium*.

*rhodocladium* Herggr. — Br. globosum.

- rhypariocaulon* C. M. in *Fl.* 1886, p. 507.  
 Ster. Cortic. — *Afr.* : Ins. Fernando-Po.
- rigescens* Wils. — Webera.  
*rigidulum* Wils. — Didymodon.
- rigidum* (Hsch.) C. M. *Syn.* I, p. 263 (1849).  
 Br. patens Wils. in *Lond. Journ. of bot.* 1844, p. 155 ; C. M. *op. cit.* p. 250 (*fid. Mitt.*).  
*Mnium rigidum* Hsch. *Fl. brasil.* I, p. 46 (1840).  
*Rhodobryum rigidum* Par. *op. et l. cc.*  
 2. *Rupic* — *Am. merid.* : Brasil. austr. or., Monte Video.
- rigidum* Huds. — *Aloina aloides*.
- ripense* C. M. in *Bull. hb. Boissier*, 1897, p. 551.  
 Paludic. — *Am. sept.* : Jamaic.
- rivale* C. M. in *Linn.* 1878-79, p. 300.  
 2. Ad aren. rivul. — *Am. merid.* : Argentin. subtropic. Tucuman et Jujiens.
- riolare* Arn. in *Rev. bryol.* 1898, p. 6.  
 Ster. Ad rip. rivul. — *Eur.* : Suecia merid. occid.
- riolare* Hoffm. — *Rhacomitrium protensum*.
- robustum* C. M. in *Engler's Bot. Jahrb.* 1883, p. 77 et in *Forschungsreise.... Bot.* p. 14 (1889).  
*Rhodobryum robustum* Par. *op. cit.* p. 1120.  
*Syn.* — *Afr.* : Ins. Kerguelen.
- robustum* Hpe. in *Linn.* 1856, p. 205 et 1859-60, p. 627.  
*Rhodobryum robustum* Par. *op. et l. cc.*  
 — *Pac.* : Austral. or. (Victoria).
- Rodriguezii* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1893, I, p. 129 ;  
*Ren. M. Masc. Mad.* n. 117.  
 Br. Bescherellei eor. in *op. cit.* 1891, II, p. 188.  
 2. — *Afr.* : Ins. Borbon.
- Roellii* Philib. in *Rev. bryol.* 1890, p. 56.  
 Polyg. — *Am. sept.* : Washington.
- Roscheri* Ltz. *Moosst.* p. 157 (1864).  
 2. In sabul. — *Afr.* : Zanzibar.
- Rosenbergiae* Hagen in *Det. K. Norske Vid. sells. Skript.* n. 2.  
 1. Ad rnp. granit. humidiusc. — *Eur.* : Norveg.
- roseolum* C. M. in *Linn.* 1878-79, p. 287.  
*Rhodobryum roseolum* Par. *op. et l. cc.*  
 Fl. et fr. ign. — *Am. merid.* : Argentin. subtropic.
- roseum* { Mitt. — Br. minuti-rosatum.  
 { W. Arn. — *Rhodobryum* Beyrichianum.
- rostratum* { C. M. — *Brachymenium*.  
 { Schrad. — *Mnium*.
- rosulans* C. M. in *Hedwig.* 1897, p. 96.  
 — *Am. merid.* : La Plata.

*rosulatum* C. M. in *Fl.* 1880, p. 474.

*Rhodobryum rosulatum* Par. op. et l. cc.

Ster. In graminos. — *Afr.* : Kilima N'Djaro.

*rosulatum* C. M. in *Bot. Zeit.* 1856, p. 416.

*Brachymenium rosulatum* Besch. *Prodr. bryol. mexic.* p. 55 (1871).

*Rhodobryum rosulatum* Par. op. et l. cc.

2. — *Am. sept.* : Mexic.

*rosulicoma* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1892, I, p. 166.

*Rhodobryum rosulicoma* Par. op. et l. cc.

2. fr. ign. Lignic. — *Am. sept.* : Costa Rica.

*rubellum* { Brid. — *Mniobryum carneum*.  
Hoffm. — *Didymodon*.

*rubens* Mitt. in *Journ. of bot.* 1856, p. 232 et *M. Ind. or.* p. 69 (1859).

Br. *erythrocarpum* var. *limbatum* Berth. in *Rev. bryol.* 1883, p. 67.

Br. *erythrocarpum* var. *silvaticum* Hpe. *exs. n.* 201.

2. — *Eur.* : Norveg., Angl., Gall. centr. et merid.

As. : Sikkim, Assam. *Am. sept.* : Temper.

*rubicundulum* C. M. et Kindb. in *Bidr.* p. 454 (*nom.*, 1890) et in *Mac. Cal.* p. 129 (1892).

Br. *Muhlenbeckii* (fid. Kindb. *Eur. and N. Am. Bryin.* p. 356).

2. Rupic. ad niv. aetern. — *Am. sept.* : M. Selkirk.

*rubicundum* C. M. in *Bot. Zeit.* 1859, p. 207.

— *Afr.* : Prom. Virid.

*rubiginum* C. M. in *Nuov. Giorn. bot. ital.* 1898, p. 165.

2. — *As.* : China centr. (prov. Schen-Si).

*rubiginosum* Hook. f. et Wils. *Fl. Tasman. II*, p. 190, t. 173, f. 9 (1860).

2. fr. ign. — *Pac.* : Tasman.

*rubrifolium* W. P. Sch. in *Besch. Fl. bryol. Antill. fr.* p. 32 (1876).

2. — *Am. sept.* : Guadalup.

*rubro-costatum* C. M. in *Engler's Bot. Jahrb.* 1883, p. 83 et in *Forschungsreise.... Bot.* p. 40 (1889).

Ster. — *Afr.* : Ins. Ascension.

*rubrum* Huds. — *Dicranella varia*.

*rufescens* Hook. f. et Wils. *Fl. Tasman. II*, p. 192, t. 174, f. 1 (1860) et *Handb. of the N. Zeal. Fl.* p. 439 (1867).

Br. *Billardieri* (fid. Mitt).

2. — *Pac.* : Austral. or. (Victoria), Nov. Seeland., Tasman.

« var. *brevifolium* H. f. et W. op. et l. cc.

— *Pac.* : Tasman.

« var. *mammillatum* eor. op. et l. cc.

Ubi praeced.

*rufescens* Sm. — *Dicranella*.

*rufinerve* C. M. in *Linn.* 1874, p. 549.

2. — *As.* : Ins. Philippin.



- rufo-limbatum *Ren. et Card. in Bull. Soc. roy. bot. Belg.* 1892, I, p. 165.  
2. — *Am. sept.* : And. Costa-Ricens.
- rufo-nitens *Hpe. Symb.* 1877, p. 725.  
Ster. — *Am. merid.* : Brasil. austr. or.
- rufum *Fergus. mss.* ; *Limpr. Laubm. Deutschlands.... II*, p. 296 (1892).  
Syn. — *Eur.* : Anglia.
- rugosum { C. M. { — Brachymenium.  
                  Hoffm. — Dicranum undulatum.
- rupeum Neck. — *Leptobryum piriforme.*
- rupicola Schleich. — *Br. pallescens.*
- rurale { L. — *Barbula.*  
          *unguiculatum, hirsutum elatius et ramosius* Dill. — *Har. ruralis.*
- ruralis facie, foliis minoribus et non pilosis Dill. — *Barbula unguiculata*  
var. patagonica.
- Rusbyanum C. M. — *Webera.*
- rutilans { Brid. — *Webera.*  
          Bryol. eur. — *Webera Schimper.*  
          C. M. — *Br. æneum.*
- Salaminae C. M. — *Brachymenium.*
- salinum *Hagen in sched.* ; *Limpr. Laubm. Deutschl.... II*, p. 334 (1892).  
Br. malangense *Kaur. et Arn. mss.* ; *Arn. in Rev. bryol.* 1898, p. 39 et in  
*Bot. Not.* 1899, p. 79, p.p.  
Syn. In arenos. marit. — *Eur.* : Norveg. occid.
- saltense *Par. Ind.* p. 212 (1894).  
Br. angustifolium *Kaur. in Bot. Not.* 1887, p. 113 c. ic.  
Br. campylocarpum ? (*fid. Kindb.*).  
1. — *Eur.* : Norveg. bor.
- Sandbergii Holzing. — *Roellia.*
- Sandii *Ds. et Mk. M. frond. ined. Archip. ind.* p. 21, t. 10 (1854) ;  
C. M. *Syn. I*, p. 300 (1849) ; *Bryol. jav. I*, p. 145 (1855-61).  
1 et Syn. — *As.* : Java.
- sanguineum Brid. — *Br. erythrocarpum.*
- sanguilentum *Ren. et Card. mss.*  
Br. capillare var. *Lesq. et Jam. Man.* p. 236 (1884).  
Br. obconicum *Sull. M. bor. am. I ed.* n. 187 b.  
Br. occidentale var. obconicum *Sull. et Lesq. M. bor. am. II ed.*  
n. 283 et 284.  
2. Lignic. — *Am. sept.* : Calif.
- saprophilum C. M. in *Dusen M. Camer. n.* 752 ; *Broth. in Engler's Bot. Jahrb.* 1897, p. 248.  
*Rhodobryum saprophilum Par. op. et l. cc.*  
2. Lignic. — *Afr.* : Cameroon.
- Sartorii *W. P. Sch. in Besch. Prodr. bryol. mexic.* p. 56 (1871).  
*Rhodobryum Sartorii Par. op. et l. cc.*  
2. — *Am. sept.* : Mexic.
- Sauteri *Bryol. eur. fasc.* 32 (Vol. IV), *Mon. Suppl.* p. 10, t. 377 (1846) ;  
C. M. *Syn. I*, p. 280 (1849) ; *W. P. Sch. op. cit.* p. 435.  
1. Ad terr. arenac. humid. — *Eur.* : Salisb., Pinzgov., Vall. Tellin.  
super. *As.* : Caucas.

*Sawyerii* Ren. et Card. in Bot. Gaz. 1889, p. 95, t. XIII c.

2. Arenic. et lignic. — Am. sept. : Florid., Ludovic.

*saxatile* Hagen in Tromsøe Mus. Aarsheft. 1898-99, p. 170.

Heteroic. Ad rup. madid. — Eur. : Norveg. bor.

*scabridens* Mitt. — Brachymenium.

*scalariforme* Joerg. in Christiania Vid. Selsk. Foerh. 1894, p. 79.

Br. subtumidum (fid. Hagen).

— Eur. : Norveg.

*Schauinslandi* C. M. in Abh. d. nat. Ver. zu Bremen, 1900, p. 502.

2. — Pac. : Ins. Chatham.

*schensianum* Par. — Br. humillimum.

*Schimperi* { C. M. — Webera.

{ Sb. — Br. Liebmanni.

*Sch<sup>is</sup>ti* { Gunn. — Dicranum Blyttii.

{ CEd. — Cynodontium.

*schisticolum* C. M. — Webera.

*schistoneuron* (Kze.) Par. Ind. p. 212 (1894).

Br. gracilescens C. M. Syn. I, p. 261, (1849).

Mnium schistoneuron Kze. in Poeppig Pl. Chil. n. 234.

2. — Am. merid. : Peruv.

« var. decumbens (C. M.) Par. op. cit. p. 213.

Br. Auberti Hsch. Fl. Brasil. p. 45, t. 2, f. I (1840).

Br. gracilescens var. decumbens C. M. op. et l. cc.

— Am. merid. : And. Quitens., Brasil. austr. or.

« var. duplicatum (Ren. et Card.) Par. Suppl. Ind. p. 73 (1900).

Br. gracilescens var. duplicatum R. C. in Bull. Soc. roy. bot. Belg. 1895, I, p. 62.

Ster. — Am. merid. : Brasil. or.

« var. densifolium (Brid.) Par. Ind. l. cc.

Br. Auberti Schwaegr. Suppl. II, P. II, II, p. 195 (1826) et Sp. M. p. 54.

Br. densifolium Mitt. M. austr. am. p. 313 (1869).

Br. gracilescens var. densifolium C. M. op. et l. cc.

Br. Polla densifolia Brid. Bryol. univ. I, p. (1826).

Rupic. et terr. — Am. merid. : And. Quitens. et Peruv., Brasil. austr. or.

*schizotrichum* C. M. in Rehm. M. Afr. austr. n. 241.

— Afr. : Prom. B. Sp.

*Schleicheri* Schwaegr. Suppl. I, P. II, p. 113, t. 73, (1816); Brid. Bryol. univ. I, p. 683 (1826); W. P. Sch. op. cit. p. 462.

Br. alpinum Schleich. Pl. helvet. exs.; Vill. Pl. Dauph. III, p. 889 (1786).

H. et T. Musc. brit. p. 122 (1818).

Br. turbinatum { var. elongatum Bryol. eur. fasc. 6-9, p. 65 (1839).

{ var. gracilescens W. P. Sch. Coroll. p. 77 (1856).

{ β praelongum Bryol. eur. l. o. t. 372 β.

Mnium glaciale Schleich. Cent. III Pl. helv. n. 41; Catal. 1807.

Mn. Schleicheri Brid. Sp. M. III, p. 60 (1817).

2. Ad scatur. — Eur. : Reg. subalp. et alp. zon. centr.; Pyren., S. Nevada. As. : N. W. Himalaya, Tibet. occ., Baltistan, Kashmir, Tchitral, Panjab, Caucas. occ. et centr. Am. sept. : Groenland., M. Rupestr. et Selkirk, Utah, Calif.



- 17

*speciosum* { Mitt. — *Leptotheca* ?  
 { Voit. } — *Br. pallescens* (fid. *Limpr.*).  
*spectabile* C. M. — *Webera*.  
 — *Br. pallens* var. *speciosum* (fid. W. P. Sch.).

*Spegazzinii* C. M. in *Fl.* 1885, p. 400.

Syn. Ad terr. et rup. humid. — *Am. merid.* : *Fuegia austr.*

*spirophyllum* Kindb. in *Nuov. Giorn. bot. ital.* 1896, p. 17.

*Br. capillare* var. *frigidum* W. P. Sch. ?

— *Eur.* : *Ital. super.* (Ticin.).

*sphaericarpon* Sw. — *Philonotis*.

*sphagnadelphus* C. M. — *Webera*.

*Sphagni* { Brid. — *Br. Duvalii*.

{ C. M. p.p. — *Webera sphagnicola*.

*sphagnicola* Bryol. eur. — *Webera sphagnicola*.

*sphagnoides* Jacq. — *Hedwigia ciliata*.

*spinidens* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1893, I, p. 117;

*Ren. M. Masc. Mad.* n. 29.

2. — *Afr.* : *Madagascar.*

*spinifolium* Philib. in *Rev. bryol.* 1899, p. 29.

1. In uligin. — *As.* : *M. Alatau.*

*spinosum* Voit. — *Mnium*.

*spirifolium* C. M. — *Brachymenium*.

*spitzbergense* Arn. in *Oefv. af K. Sv. Vet.-Akad. Foerh.* 1900, p. 119.

Syn. — *Eur.* : *Spitzberg.*

*splachnobryoides* C. M. in *Linn.* 1880-82, p. 375.

Ster. — *Am. merid.* : *Argentina. Cordob.*

*splachnobryoides* C. M. (1883). — *Br. subsplachnobryoides*.

*splachnoides* { C. M. — *Brachymenium*.

{ Dicks. — *Blindia acuta*.

*splachnoideum* C. M. in *Engler's Bot. Jahrb.* 1883, p. 77 et in *Forschungsreise.... Bot.* p. 13 (1889).

Ster. — *Afr.* : *Ins. Kerguelen.*

*spongiosum* W. et Dy. — *Philonotis*.

*Sprucei* Mitt. — *Acidodontium*.

*spurium* Hoffm. — *Dicranum*.

*squalidum* Vill. — *Br. capillare*.

*squarripilum* C. M. in *Fl.* 1886, p. 280.

Ster. Terr. — *Afr.* : *Ins. Sao Thoma.*

*squarrosus* Kindb. in *Hedwig.* 1896, p. 66.

2. — *Am. sept.* : *Oregon.*

*squarrosus* L., Drumm. — *Paludella*.

*Starkeanum* Hoffm. — *Pottia*.

*Steffeni* Dusen. *M. Am. austr.* n. 400.

Terr. — *Am. merid.* : *Patagon. occid.*

- Roth. — *Mnium*.  
*hornum silvaticum capsulis* } — *Mnium hornum* (*fid. C. M.*).  
*magnis nutantibus* Dill. } — *Webera cruda* (*fid. Limpr.*).  
*lacustre, foliis rubris* } — *Didymodon rubellus* (*fid. C. M.*).  
*capillaceis* Dill. } — *Ceratodon purpureus* (*fid. Limpr.*).  
*nitidum, pallidum,* }  
*capsulis tenuis-* } — *Barbula convoluta* (*fid. C. M.*).  
*simis* Dill. } — *Barbula revoluta* (*fid. Limpr.*).  
*stellare* {  
*majus, capsulis ovatis pendulis* Dill. — *Rhodo-*  
*bryum roseum*.  
*roseum* {  
*pendulum minus,* } — *Mnium stellare* (*fid.*  
*capsulis et setis* } *C. M.*).  
*longioribus* Dill. } — *Br. capillare* (*fid.*  
*Limpr.*).  
*stellatum* { Dicks. — *Barbula agraria*.  
Lam. et DC. } — *Mnium stellare* (*fid. Brid.*).  
} — *Mnium hornum* (*fid. Lindb.*).  
Schreb. — *Aloina rigida*.  
*stelligerum* Dicks. — *Hymenostylium curvirostre*.  
*stellipilum* C. M. in *Hedwig*. 1899, p. 68.  
2. — *Afr.* : Transvaal.  
*stenocarpum* *Limpr.* in *LXI Jahresb. d. schles. Ges. f. Vaterl. Kult.* 1884,  
p. 219 et *Laubm. Deutschl. .... II*, p. 292 (1892).  
*Br. Brownii* var. *minor* Kaur. mss. ; *Bryhn* in *N. Mag. f. Naturv.* 1892,  
p. 125.  
1. — *Eur.* : Penins. Kola, alp. Norveg. (Dovre), ins.  
Aland.  
*stenodon* *Hagen* in *Tromsø Mus. Aarsheft.* 1898-99, p. 145.  
Syn. In fiss. rup. — *Eur.* : Norveg. bor.  
*stenopyxis* C. M. in *Linn.* 1878-79, p. 840.  
2. — *Am. merid.* : Venezuela.  
*stenothea* J. O. Bomanss. in *Rev. bryol.* 1899, p. 11.  
Syn. — *Eur.* : Ins. Aland.  
*stenotheceum* (*Hpe.*) *Broth. Natür. Pflanzenfam.* p. 595.  
*Rhodbryum stenotheceum* *Hpe. Symb.* 1874, p. 497 ; *Par. Ind.* p. 1121  
(1897).  
Terr. in silv. — *Am. merid.* : Brasil. austr. or.  
*stenotrichum* C. M. in *Fl.* 1887, p. 219.  
Syn. — *Am. sept.* : Alaska.  
*Stirtoni* W. P. Sch. *Syn. II* ed. p. 470 (1876) ; *Hagen* in *Det. K. Norske*  
*Vid. sells. Skr.* 1899, n. 2.  
*Br. barbatum* *Wils. mss.* ; *Hunt* in *Mem. Lit. Phil. Soc. Manch.* 1872,  
p. 238 ? (*fid. Limpr.*).  
2. — *Eur.* : M. Scot., Norveg. *Am. bor.* : Fret.  
Scoresby.  
*streptophyllum* *Kindb. Eur. and N. Am. Bryin.* p. 359 (1897).  
*Ster.* — *Am. sept.* : Columb. brit.  
*streptorhodon* C. M. in *Bull. hb. Boissier*, 1897, p. 179.  
*Rhodbryum streptorhodon* *Par. op. et l. cc.*  
— *Am. sept.* : Guatemala.

- striatum* L. — *Orthotrichum leiocarpum*.  
 var. *ulophyllum* Ehrh. — *Ulota crispa*.  
*strictipes* C. M. — Webera.
- strictum* Philib. in *Rev. bryol.* 1886, p. 18.  
 Br. *warneum* (fid. *Limpr.*).  
 1. — Eur. : Norveg.
- strictum* Hoffm. — *Ceratodon purpureus*.
- strigosum* Wils. in *Kew Journ. bot.* IX, p. 369 (1857).  
 Br. *Wrightii* Mitt. *M. Ind. or.* p. 71 (1858).  
*Rhodobryum strigosum* Par. *Suppl. Ind.* p. 300 (1900).  
 — As. : M. Neilgherr., Ceylan.
- strumiferum* Dicks. — *Cynodontium polycarpum* var. *strumiferum*.  
*styrcticum* Ren. — Br. *torquescens* f. *monoica*.
- subalpinum* Warnst. in *Fleisch, et Warnst. Bryoth. Eur. merid. n.* 161.  
 — Eur. : Italia.
- subapiculatum* Hpe. *Symb.* 1872, p. 51.  
 1. — Am. merid. : Brasil. austr. or.
- subappressum* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1894, II, p. 123.  
 2. — Afr. : Madagascar.
- subareoblastum* C. M. in *Dusen M. Camer. n.* 115 a.  
 — Afr. : Cameroon.
- subargenteum* Hpe. in *Linn.* 1874, p. 213.  
 2. — Afr. : Madagascar.
- subatro-purpureum* C. M. in *Linn.* 1871-73, p. 147.  
 2. — Pac. : Austral. or. (Queensland).
- subcalophyllum* Philib. in *Rev. bryol.* 1899, p. 15.  
 1. In uligin. — As. : M. Terskei Alatau.
- subcampylopodioides* Par. *Suppl. Ind.* p. 73 (1900).  
 Br. *campylopodioides* C. M. in *Nuov. Giorn. bot. ital.* 1896, p. 95.  
 Ster. Terr. — As. : China centr. (prov. Schen-Si).
- subcapillifolium* C. M. — *Brachymenium*.
- subcapitellatum* Par. *op. et l. cc.*  
 Br. *capitellatum* C. M. in *op. cit.* 1898, p. 164.  
 2. — As. : China centr. (prov. Schen-Si).
- subcavifolium* Par.  
 Br. *cavifolium* C. M. in *Rehm. M. Afr. austr. n.* 255.  
 — Afr. : Austral.
- subcorrugatum* C. M. in *Bull. hb. Boissier*, 1897, p. 182.  
 2. — Am. sept. : Guatemala.
- subcucullatum* Kindb. — Webera.
- subcurvatum* Mitt. — Webera.
- subdecursivum* C. M. in *Rehm. M. Afr. austr. n.* 277 et in *Hedwig.* 1899, p. 74.  
 Ster. — Afr. : Prom. B. Sp.
- subenerve* Brid. — *Eriopus remotifolius*.

- suberythrocarpon *C. M. in Bot. Zeit.* 1856, p. 407.  
*Br. erythrocarpon var. australis Hpe. in Preiss Pl. Nov. Holl.*  
 2. — *Pac.* : Austral.
- suberythrocaulon *C. M. in Krause M. Ecuador.*  
*Rhodobryum suberythrocaulon Par. Ind. p. 1191 (1897).*  
 — *Am. merid.* : And. Quitens.
- subfasciculatum (*Hpe.*) *Mitt. in Trans. a. Proced. of the roy. Soc. of Victoria*, 1883, p. 73.  
*Rhodobryum subfasciculatum Hpe. in Linn.* 1876, p. 312 ; *Par. op. et l. cc.*  
 Ster. — *Pac.* : Austral. or. subtropic.
- subgemmaferum *Kindb. Eur. and N. Am. Bryin. p. 368 (1897).*  
 2. Ad terr. arenac. — *Eur.* : Norveg.
- subglobosum *Schlieph. in Fl.* 1897, p. 330.  
 2? — *Am. merid.* : And. Venezuel.
- sublencophyllum *C. M. in Fl.* 1897, p. 330.  
 2? — *Am. merid.* : And. Venezuel.
- ? submicrostegium *Kindb. in Bot. Not.* 1896, p. 5.  
 Ster. — *Am. sept.* : Groenland.
- submucronatum *Philib. in Rev. bryol.* 1899, p. 25.  
 1. — *As.* : M. Alatau.
- subnitidulum *Kindb.* — *Br. archangelicum.*  
*subobliquum Lindgr.* — *Br. pseudo-triquetrum.*
- subobtusifolium *C. M. in Bot. Centralbl.* 1883, n. 2-4.  
 2. — *As.* : Penins. Tschutschica.
- suboeneum *Hpe. et C. M. in Linn.* 1853, p. 494.  
 2. — *Pac.* : Austral. or.
- subolivaceum *C. M. in Hedwig.* 1898, p. 103.  
*Rhodobryum subolivaceum Par. Suppl. Ind. p. 300 (1900).*  
 2. — *Pac.* : Austral. or. (Nov. Vales.).
- subpachypoma *Hpe. in Linn.* 1869-70, p. 518.  
 2? — *Pac.* : Austral. or.
- subpilosum *Mitt. M. austr. am. p. 307 (1869).*  
 2. — *Am. merid.* : And. Quitens. ; Peruv.
- subplumosum *Broth. et Par. in Rev. bryol.* 1900, p. 77.  
 Ster. Lignic. et terr. — *As.* : Tonkin.
- subpurpurascens *Kindb. in Mac. Cat. p. 119 (1872).*  
 2. Ad terr. humid. — *Am. sept.* : Columb. brit.
- subroseum *Besch. Prodr. bryol. mexic. p. 56 (1871).*  
*Br. Liebmanni (fid. Broth.).*  
*Rhodobryum subroseum Par. Ind. p. 1191 (1897).*  
 2. — *Am. sept.* : Mexic.
- subrotundifolium *Jaeg. Ad. II, p. 707 (1877-78).*  
*Argyrobryum subrotundum Hpe. in Linn.* 1876, p. 312.  
 Ster. — *Pac.* : Austral. or.



subrotundum *Brid. Bryol. univ. I, p. 673* (1826); *Bryol. eur. fasc. 32. Mon. Suppl. p. 9, t. 361* (1846); *C. M. Syn. I, p. 275* (1849); *W. P. Sch. op. cit. p. 434*.

*Br. brunnescens Schleich. in sched.*

*Br. caespitium Wils. in Kew Journ. Bot. IX, p. 331* (1857).

*Br. cernuum ej. in op. et l. cc.*

*Br. pallescens* ♂ subrotundum *Bryol. eur. fasc. 6-9, Mon. p. 52, t. 360* (1839).

*Br. pohliaeforme Brid. Bryol. univ. I, p. 687* (1826).

*Br. pumilum Wils. in op. et l. cc.*

1. *Ad terr. humos. et in fiss. rup. — Eur. : Alp. Norveg., Sudet., Salisb., Stir., Carinth., Tirol., Helvet. As. : Jenissei infer., Himalaya. Am. sept. : Groenland., Wyoming, Columb. brit., Colorado, Calif.*

« *var. clavata Hagen in Tromsø Mus. Aarsheft. 1898-99, p. 191.*

*Ut typ. — Eur. : Norveg. bor.*

« *subsp. turfaceum Kindb. in Bot. Not. 1896.*

*— Eur. : Alp. Norveg.*

*subrotundum Mitt. — Acidodontium.*

*subrutilum Limpr. mss. ; Hagen in op. cit. p. 194.*

2. *Ad rup. calcar. — Eur. : Norveg. bor.*

*subsmaragdinum C. M. — Brachymenium.*

*subsplachnobryoides Par.*

*Br. splachnobryoides C. M. in Engler's Bot. Jahrb. 1883, p. 77 et in Forschungsreise... Bot. p. 13* (1889); *Par. Ind. p. 215* (1894).

*Ster. — Afr. : Ins. Kerguelen.*

*subtomentosum (Hpe.) Mitt. in Trans. a. Proced. of the roy. Soc. of Victoria, p. 73.*

*Br. truncorum H. f. et W. Fl. of the N. Zeal. (1855).*

*Rhodobryum subtomentosum Hpe. in Linn. 1869-70, p. 516 ; Par. op. et l. cc.*

2. *— Pac. : Austral. or. (Nov. Vales., Victoria).*

*subtumidum Limpr. ap. Joergens. in Christiania Vid. Selsk. Foerh. 1894, p. 78.*

*— Eur. : Norveg.*

*subturbيناتum C. M. — Br. turbيناتum var. imbricatum.*

*subulatum L. — Barbula.*

*var. β Weis. — Barbula muralis.*

*sudeticum Ludw. — Webera Ludwigii.*

*Sullivanii C. M. mss. ; Broth. in Oefv. af Finnska Vet.-Akad. Foerh. 1893, p. 48.*

2. ♂ *ign. Ad. rup. humid. — Pac. : Austral. or. (Victoria), Tasman.*

*superpensum C. M. in Bull. hb. Boissier, 1893, p. 32 et in Ule Coll. M. brasil. n. 1770.*

2. *Terr. in silv. — Am. merid. : Brasil. or. (S. do Itatiaia).*

surculo	{	<i>brevi ramoso, setis contortis, capsulis ovalibus erectis, operculis arcuatis</i> Stroem. — <i>Dicranella cerviculata (fid. Hagen).</i>
		<i>filiformi ramoso, foliis subulatis, capsulis piriformibus sessilibus, operculis arcuatis conicis</i> Stroem. — <i>Amphoridium lapponicum (fid. Sw.).</i>
		<i>bas. procumbente, foliis capillaribus erectis, operculo rostrato</i> Stroem. — <i>Ditrichum tortile var. pusilla (fid. Hagen).</i>
		<i>stricto</i> { <i>procumbente, foliis ex ovato linearibus, capsulis elongatis erectis operculis aristatis</i> Stroem. — <i>Trematodon ambiguus (fid. Hagen).</i>

*synoico-caespitium* C. M. et Kindb. in *Bidr. p.* 454 (nom., 1890) et in *Mac. Cat. p.* 128 (1892).

Syn. Ad terr. arenac. — *Am. sept.* : Columb. brit.

*synoico-crudum* C. M. — Webera.

*synoicum* C. M. in *Hedwig. 1898, p.* 86.

Syn. — *Pac.* : Austral. or. et merid.

*syriacum* Ltz. M. *Ehrenb. p.* 45, t. XII (1868).

Webera sacra ej. (*fid. Jur. et Milde, Beitr. p.* 597).

W. sinaitica ej. pl. ♂ (forsan ! *fid. iisd.*).

2. — *As.* : Pers., Caucas., Taurus, Syr.

*sysphinctum* Limpr. *Einig. neue Art.... 1883, p.* 240 et *Laubm. Deutschl.... II, p.* 352 (1892).

Syn. In arena humid. — *Eur.* : Norveg.

*systylium* C. M. — Brachymenium.

*Taitae* C. M. in *Fl. 1879, n.* 24.

Ster. — *Afr.* : M. Kenia.

*tasmanicum* Hpe. in *Linn. 1852, p.* 714.

Syn. — *Pac.* : Tasman.

*Tauriscorum* Limpr. — Br. archangelicum.

*Taylori* C. M. — Acidodontium subrotundum.

*tectorum* C. M. in *Nuov. Giorn. bot. ital. 1896, p.* 95.

Ster. Ad tecta. — *As.* : China centr. (prov. Schen-Si).

*tectorum* de Not. { — Br. caespitium var. imbricatum (*fid. C. M.*).

{ — Br. hadium (*fid. Limpr.*).

*tegulare, humile, pilosum et incanum* Dill. — Tortula muralis.

var. non pilosum ej. — Tortula marginata.

*tenellum* C. M. — Brachymenium.

*Teneriffae* Hpe. mss.; C. M. in *Bot. Zeit. 1832, p.* 12.

2. — *Afr.* : Ins. Teneriff.

tenue	{	Dicks. — Ceratodon purpureus.	
		Rav. — Br. Funckii var. tenue.	
		barbatum, foliis angustioribus et rarioribus Dill. — Barbula fallax.	
		imberbe et pallidum, foliis crebrioribus Dill.	{ <i>fid. C. M.</i> .) — Br. fallax.
			{ ( <i>fid. Limpr.</i> ) — Didymodon rigidulus.
		stellatum, setis purpureis Dill. — Ceratodon purpureus.	

*tenuicaule* Mont. in *Ann. sc. nat. 1845, IV, p.* 104 et *Syll. p.* 32 (1856);

C. M. *Syn. I, p.* 293 (1849); *Mitt. M. austr. am. p.* 307 (1869).

2. — *Am. merid.* : Chile.

*tenuifolium* H. f. et W. — Webera ?

*tenuirete* Dusen M. *Am. austr. n.* 651.

Saxic. — *Am. merid.* : Patagon. occid.

*tenuisetum* Limpr. in *LXXXIII Jahresb. d. schles. Ges. f. Vaterl. Kult. 1896.*

Polyg. Ad terr. humid. — *Eur.* : Stiria (Leibnitz).

*tenuissimum* Tayl. — Eccremidium arcuatum.

*terebellum* Hpe. in *Linn. 1863, p.* 130; *Mitt. op. cit. p.* 311.

Syn. (Hpe.), 1 (Mitt.). Terr. — *Am. merid.* : And. Nov. Granat.

*terebellum* W. P. Sch. — Br. genucaule.

*teres* Lindb. in *Oefr. af K. Sv. Vet.-Akad. Foerh.* 1866, p. 545.

Br. nitidulum var. *teres* Berggr. M. Spetsb. n. 76 b et in K. Sv. Vet.-Akad. Handl. XIII, n. 7, p. 64.

1. Terr. — Eur. : Spitzberg., ins. Ursor. Am. sept. : Frantz Joseph Fjord, fret. Scoresby, ins. Sabine.

*teretiussculum* Hook. — Br. alpinum.

*terminale* C. M. — Anomobryum.

*terskeiense* Par.

Br. paradoxum Philib. in *Rev. bryol.* 1900, p. 74.

1. — As. : M. Terskei Alatau.

*tessellatum* Philib. in *Rev. bryol.* 1899, p. 60.

Syn. — As. : M. Alatau.

*tetragonum* Dicks. — Conostomum boreale.

*Therioti* Philib. in *Rev. bryol.* 1897, p. 17.

Syn. Terr. — Eur. : Alp. Delphinat.

*Thomsoni* Mitt. M. Ind. or. p. 75 (1859).

Br. cuspidatum Wils. in *Keur Journ. bot.* IX, p. 364 (1857).

2. — As. : Tibet. occid., Kashmir.

*tibetanum* Mitt. op. cit. p. 72.

Br. caespiticium Wils. in op. cit. p. 332.

1. Ad rup. irror. — As. : Tibet. occid., Kashmir.

*timmiostomoides* Philib. in *Rev. bryol.* 1898, p. 49.

1. — As. : M. Alatau.

*tjiburrumense* Fleisch. M. Archip. ind. n. 144 (1900).

Ster. Ad rup. irror. — As. : Java.

*tomentosulum* Par.

Br. tomentosum Kindb. Eur. a. N. Amer. Bryin. p. 361 (1897) ; Par. Suppl. Ind. p. 74 (1900).

— Am. sept. : M. Rupestre.

*tomentosum* Limpr. mss.

Br. arcticum var. *tomentosa* (Limpr.) Joergens. in *Christiania Vid. Selsk. Forh.* 1894, p. 76.

— Eur. : Norveg.

*tomentosum* { Brid. — Br. capillare var. domingense.  
Kindb. — Br. tomentosulum.  
Sw. — Breutelia.

*tophaceum* DR. et Mont. in *Ann. sc. nat.* XII, p. 319 et *Mont. Syll.* p. 33 (1856) ; C. M. Syn. II, p. 574 (1851).

2. Ad rivul. — Afr. : Alger.

*torquescens* Bruch Fl. Sard. exs. (1828) ; de Not. Syll. n. 163 (1838) ; *Bryol. eur. fasc.* 6-9 (Vol. IV), *Mon.* p. 49, t. 358 (1839) ; C. M. Syn. I, p. 277 (1849) ; W. P. Sch. op. cit. p. 431.

Br. capillare var. *torquescens* Husn. *Musc. gall.* p. 240 (1889).

Syn. Ad terr. lapid., mur. et rup. — Eur. : Suec., Angl., Hibern., vall. Rhen., Helvet. occid., zon. merid. As. : Japon., Himalaya

occid., Kashmir, Pers., Caucas., Minor. Afr.: Alger., ins. Canar.,  
Madeir., Kerguelen, Transvaal, prom. B. Sp. Am. sept.: Nov.  
Scot.? Gaspes.? Nevada, Oregon, Washington, ins. Vancouver,  
Texas, Calif., Nov. Mexic. Am. merid.: Monte Video, ins. Chiloe.  
Pac.: Austral. or. (Victoria), Nov. Seeland, Tasman.

« *var. brevifolium* Lindb. in *Bol. d. soc. Broter*. 1889, VII.  
— *Eur.* : Lusit.

" *var. gracile* Besch. Cat. M. Alger. p. 24 (1882).  
— Afr. : Alger.

\* *f. monoica* Spr. *M. Pyren.* n. 134 et in *Trans. of the Linn. Soc. of Edinb.* 1850, III, p. 158.

*Br. brunnescens* Ren. in *Rev. bryol.* 1883, p. 81.

*Br. fuscescens* Spr. in op. et l. cc.

Br. systicum *Ren. in op. cit. p. 82.*

1. — *Eur.* : Pyren.

*torquescens* W. P. Sch. — Br. pseudo-torquescens.

torquescentulum C. M. in. *Hedwig.* 1899, p. 71.

2. Loc. umbr. inter frutic. — *Äfr.* : Prom. B. Sp.

*var. nutans* ej. in op. et l. cc.  
Ut et ubi typ.

*tortifolium* Brid. — Br. cyclophyllum.

*tortuosum* L. — Tortella.

*torulosicolum* C. M. — Br. Bescherellei.

*Tozeri* Grev. — Epipterygium.

*trachyticola* C. M. — *Brachymenium*.

*transvalo alpinum* C. M. — Philonotis.

*trematodonleum* C. M. — Webera.

Drumm. — *Meesea longiseta* var. *alpina*.

Hedw. — Webera nutans.

L. — Meesea uliginosa.

*acaulon palustre minimum,*  
*setis et capsulis* { *brevissimis* Dill. — Seligeria  
  *pusilla.*  
  *oblongis* Dill. — Ceratodon  
  *purpureus.*

*aureum* { *capitulis piriformibus nutantibus* Dill. —  
*Leptobryum piriforme.*  
*capsulis incurvis obtusis in setis longis* Dill. —  
*Meesca uliginosa.*

***trichodes***

*capitulis* { *erectis, pediculis intortis tenuibus virentibus*  
Dill. — *Campylopus flavosus.*  
*reflexis, pediculis ima medietate rubris...*  
Dill. — *Br. atro-purpureum.*

*capsulis rubris cernuis* Dill. — *Dicranella varia*.

*erectis capitulis* } *fusco-nigris* Dill. — *Didymodon tophaceus*.  
                              } *lanuginosum* ej. — *Rhacomitrium canescens*.

*hirsutiae canescens, capitulis subrotundis* } *erectis, in pediculis breviss.*  
Dill. — *Grimmia leucophaea.*  
} *reflexis in perbrevisibus . . .*  
Dill. — *Gr. pulvinata.*

*laete virens capitulis cernuis oblongis* Dill. — *Webera nutans*.

var.  $\beta$  Lam. — *Ditrichum pallidum*.

var. *longiseta* Drumm. — *Meesea uliginosa*.

var. *minor* Drumm. — *Meesea uliginosa* var. *minor*.

- trichophorum* *Kindb. Eur. and N. Am. Bryin. p. 359* (1897).  
 Ster. — *Am. sept.* : Columb. brit.
- trichopodium* *Hagen in Tromsø Mus. Aarsheft. 1898-99, p. 131.*  
 Syn. Ad litt. mar. — *Eur.* : Norveg. bor.
- trifarium* Gmel. — *Didymodon luridus.*
- triquetrum* { Drumm. — *Meesea Albertini.*  
 { H. et T. — *Meesea triquetra.*
- triste* *de Not. in Cron. briol. ital. in Comm. II, n. 14* (1866); *W. P. Sch. op. cit. p. 457.*  
*Br. capillare* var. *triste* *Limpr. Laubm. Deutschl.... II, p. 377* (1892).  
 2. *Ad saxa granit. madid.* — *Eur.* : Alp. Rhaet., Ital. super. (pr. lac. Maj.).
- triste* { C. M. — *Br. pertriste.*  
 { Jur. — *Br. pallens* var. *angustifolium.*
- truncatulum* { Ehrh. — *Pottia lanceolata* var. *gymnostoma.*  
 { L. — *Pottia truncata.*
- truncatum* Gmel. — *Pottia truncata.*
- truncorum* *Brid. Mant. M. p. 119* (1819); *C. M. Syn. I, p. 254* (1849).  
*Br. Polla truncorum* *Brid. Bryol. univ. I, p. 699* (1826).  
*Mnium giganteum* *Schwaegr. Sp. M. p. 28 p.p.*  
*Mn. truncorum* *Brid. Sp. M. III, p. 50* (1817).  
*Rhodobryum truncorum* *Par. Ind. p. 1121* (1897).  
 2. *Ad terr. humid.* — *Afr.* : Ins. Bourbon., Franciae, Tristao da Cunha,  
 Inaccessible. *Am. sept.* : Mexic. ?
- truncorum* { H. f. et W. } — *Br. leptothecium.*  
 { } — *Br. subtomentosum* ?  
 { W. P. Sch. — *Br. Lechleri.*  
 { Wils. — *Br. ramosum.*
- Tryoni* *Broth. in Oefv. af Finska Sv. Vet.-Akad. Foerh. 1893, p. 49.*  
 2. fr. ign. — *Pac.* : Austr. or. (Queensland).
- Tsania* *C. M. in Nuov. Giorn. bot. ital. 1897, p. 248.*  
 — *As.* : China centr. (prov. Schen-Si).
- tumidum* *J. O. Romanss. in Rev. bryol. 1899, p. 12 et in Act. Soc. p. Faun. et Fl. fenn. 1899, n. 4.*  
 Syn. Ad scopul. marit. — *Eur.* : Ins. Aland.
- turbinatum* (*Hedw.*) *Schwaegr. Suppl. I, P. II, p. 109* (1816); *Brid. Bryol. univ. I, p. 681* (1826); *Bryol. eur. fasc. 6-9 (Vol. IV), Mon. p. 65, t. 372 excl. var. β et γ* (1839); *C. M. Syn. I, p. 259* (1849);  
*W. P. Sch. op. cit. p. 461.*  
*Br. appressum* *Wils. in Kew Journ. bot. IX, p. 365* (1857).  
*Br. cuspidatum* *ej. in op. cit. p. 361 p.p.*  
*Br. distans* *ej. in op. et l. cc.*  
*Br. nigricans* *Dicks. Pl. crypt. fasc. IV, p. 15, t. II, f. 10* (1801).  
*Br. palustre complicatum rubens, capsulis turbinatis pendulis* *Dill. Hist. M. p. 406, t. 51, f. 74* (1741).  
*Br. turbinatum* var. *nigricans* *H. et T. Musc. brit. p. 122* (1818).  
*Hypnum turbinatum* *W. et M. Bot. Taschb. p. 232* (1807).  
*Mnium nigricans* *Brid. Musc. rec. II, P. III, p. 90* (1803).  
*Mn. turbinatum* *Hedw. Deser. III, p. 22, t. 8* (1792).  
 2. *In loc. aren., glareos. et turf.* — *Eur.* : Ex Ital. super. usq. in Scand. As. : Tibet, occid., Kashmir, Panjab, Tchitral, Kurdistan, Caucas. occ. et centr., Arab. petr. *Am. sept.* : Groenland., Canada merid., Minnesota, Columb. brit., ins. Vancouver, Utah. *Am. merid.* : And. Quitsens.

- var. *gracilescens* Bryol. eur. — Br. Schleicheri.
- \* var. *gracilifolium* Besch. Cat. M. Alger. p. 26 (1882).  
— Afr. : Alger.
- \* var. *imbricatum* W. P. Sch. in W. Sch. M. Abyss. II ed. n. 46.  
Br. subturbinatum C. M. in Bot. Zeit. 1858, p. 155.  
— Afr. : Alp. Abyss.
- var. *latifolium* Bryol. eur. — Br. Schleicheri var. *latifolium*.  
var. *minus* Kth. — Miehllichhoferia Jamesonii.  
var. *nigricans* H. et T. — Br. turbinatum.
- var. *pallens* Drumm. } — Br. uliginosum (fid. C. M.).  
                              } — Br. pallens (fid. Kindb.).
- var. *praelongum* C. M. — Br. Schleicheri.  
var. H. et T. — Br. intermedium.  
var. eor. — Br. longisetum.  
var. Schwaegr. — Br. versicolor.  
forma Jur. — Br. Klinggraeffii.
- turbinatum { Drumm. p.p. } — Br. bimum.  
                  { H. et T. } — Br. pallens.  
                              } — Br. Schleicheri.
- turbinitiforme W. P. Sch. in W. Sch. M. Abyss.  
— Afr. : Abyss.
- turgens Hagen in Det. K. Norske Vid. sells. Skript. 1897, p. 2, f. 1.  
Mn. hymenophyllum (fid. Limpr.).  
2. ♂ ign. l'aludic. — Eur. : Norveg.
- turgidum J. O. Bomanss. in Act. Soc. pr. Faun. et Fl. fenn. n. 4.  
Syn. Terr. — Eur. : Ins. Aland.
- turgidum Sw. — Aulacomnium.
- uber Hagen in Tromsø Mus. Aarsheft. 1898-99, p. 143 (in clav. dichot.).  
— Am. sept. : W. Groenland.
- uliginosum (Bruch) Bryol. eur. fasc. 6-9 (Vol. IV), Mon. p. 18, t. 385  
(1839) ; C. M. Syn. I, p. 271 (1849) ; W. P. Sch. op. cit. p. 425.  
Br. caespitium var. Schwaegr. Sp. M. p. 60 (1830).  
Br. cernuum Lindb. in Act. Soc. sc. Fenn. X, p. 75 (1871).  
Br. turbinatum var. pallens Drumm. M. bor. am. I ed. n. 267 (fid. C. M.).  
Cladodium uliginosum Brid. Bryol. univ. I, p. 841 (1826).  
Cynodontium cernuum Hedw. Sp. M. p. 58, t. 9 (1811) (fid. Limpr.).  
Didymodon cernuus Sw. M. Suec. pp. 28 et 85, t. I, f. 2 (1799).  
Pohlia stricta Schultz in Syll. Ratisb. II, p. 181 (1828).  
P. uliginosa Bruch in Brid. Bryol. univ. l. c.
- Syn. Loc. uligin. et ad rivul. — Eur. : Anglia, Suec., Horuss. sept.,  
Dania merid., Hannov., Harz, Westphal., Rhoengeb., Hassia,  
Siles., Hungar., Austr. infer., Salisb., Tirol., alp. Rhaet., vall.  
Rhen., Gall. centr. (Morvan), Normann. As. : Saghalien, Amour,  
Pamir, Caucas. Am. sept. : Labrador, Anticosti, Terra nov.,  
Canada merid. or., Wyoming, Montana, Columb. brit., reg. sept.,  
occid. et or.
- \* var. *regulare* Warnst. in Schrift. d. Natur. Ges. in Danzig, 1896.  
— Eur. : Westphal.
- \* var. *rivale* Limpr. in LXI Jahresb. d. schles. Ges. f. Vaterl. Kult. 1884,  
p. 224 et Laubm. Deutschl. .... II, p. 325 (1892).  
Arenic. — Eur. : Alp. Norveg. (Dovre).

- ? *umbonatum* Wallr. — *Webera nutans*.  
*uncinatum* Dicks. — *Dicranella cerviculata* var. *pusilla*.  
*undulatum* { L. — *Atrichum*.  
                   { Roth. — *Mnium*.  
*unguiculatum* { Huds. — *Barbula*.  
                   { et *barbatum* { *surculis in summitate* { *crassioribus* Dill. —  
   { *tenuius et stellatim* { *Barb. unguiculata*.  
*unilaterale* Gmel. — *Ditrichum homomallum*.  
*urnigerum* Neck. — *Pogonatum*.  
*usambaricum* Broth. in *Engler's Bot. Jahrb.* 1894, p. 189.  
     2. Ad terr. aprie. — *Afr.* : Usambara.  
*utriculatum* C. M. in *Bot. Centralbl.* 1883, n. 2-4.  
     Syn. — *As.* : Penins. Tschutschica.  
*vagans* H. f. et W. — *Philonotis*.  
*vaginale* Dicks. — *Dicranella crispa*.  
*Valdiviae* Ltz. Moosst. p. 157 (1864).  
     2. — *Am. merid.* : Valdiv.  
*Valenciae* C. M. in *Linn.* 1878-79, p. 479.  
     2. — *Am. merid.* : Venezuela.  
*validinervium* C. M. in *Engler's Bot. Jahrb.* 1883, p. 77 et in *Forschungsreise...* Bot. p. 15 (1889).  
     Syn. — *Afr.* : Ins. Kerguelen.  
*validius* Hpe. Enum. M. Brasil. p. 32 (1879).  
     2. — *Am. merid.* : Brasil. austr. or.  
*vancouveriense* Kindb. in *Bull. Torr. Bot. Club*, 1889, p. 95.  
     2. Ad decliv. humid. — *Am. sept.* : Ins. Vancouver.  
*varians* C. M. in *Engler's Bot. Jahrb.* 1883, p. 87 et in *Forschungsreise...* Bot. p. 60 (1889).  
     *Br. varium* H. f. et W. ?  
     2. Ad terr. humid. — *Pac.* : Nov. Seeland.  
     ♂ *var. mucronatum* C. M. in op. et l. cc.  
     Ut et ubi typ.  
*varium* Hook. f. et Wils. *Fl. of the N. Zeal.* II, p. 85, t. 85, f. 4 (1855)  
     et *Handb.* p. 444 (1867).  
     — *Pac.* : Nov. Seeland.  
*velutinum* C. M. — *Brachynemium*.  
*ventricosum* { Dicks. — *Br. pseudo-triquetrum*.  
                   { H. et T. — *Br. bimum*.  
                   { Sibth. — *Aulacomnium palustre*.  
     var. *Schwaegr.* — *Br. Duvalii*.  
*Venturii* de Not. *Epil.* p. 409 (1869); *W. P. Sch. op. cit.* p. 469.  
     *Br. elegans* Vent. et Bott. *Enum. critt. M. ital.* p. 35 p.p. (1884).  
     Fl. et fr. ign. In irrig. editior. — *Eur.* : Tirol. merid.  
*vernum* Philib. in *Rev. bryol.* 1899, p. 64.  
     — *As.* : M. Alatau.  
*veronense* de Not. et W. P. Sch. op. et ll. cc.  
     *Br. claviger* Kaur. (fid *Limpr. Laubm. Deutschl.*... II, p. 418).  
     *Br. virescens* Kindb. in *Rev. bryol.* 1884, p. 22 (id.).  
     Fl. et fr. ign. In arenos. — *Eur.* : Norveg., Ital. super. (Verona).  
     *Am. sept.* : Fret. Scoresby.

*verrucosum* C. M. — Brachymenium.

*versicolor* A. Br. in *Bryol. eur. fasc. 6-9 (Vol. IV), Mon. p. 75, t. 379* (1839); C. M. *Syn. I*, p. 308 (1849); W. P. Sch. *op. cit.* p. 439.

Br. *turbinatum* var. *Schwaegr. Sp. M. p. 56* (1830).

2. In aren. ad rip. fluv. — Eur.: Westphal., vall. Rhen., Helvet., Sardin., Pedemont., Bavar., Tirol. merid., Carinth., Stir., Salisb., Austr. infer. Am. sept.: Florid.

*versicolor* Kaur. — Br. *excurrens*.

*versisporum* J. O. Bomanss. in *Rev. bryol. 1896, p. 91 et in Act. Soc. pr. Faun. et Fl. fenn. 1899, n. 4.*

2. Ad terr. arenos. — Eur.: Norveg. bor., ins. Aland.

*verticillatum* C. M. in *Bull. hb. Boissier, 1898, p. 30 et in Ule Coll. M. brasil. n. 1769.*

2. Terr. — Am. merid.: Brasil. or.

*verticillatum* { Dicks. — *Hymenostylium curvirostre*.  
Lightf. — *Blindia acuta*.  
L. — *Eucladium*.

var.  $\beta$  Huds. — *Blinuia acuta*.

*vinosulum* Besch. in *Ann. sc. nat. VII ser., II, p. 91.*  
Ster. — Afr.: Ins. Mayotte.

*virens* { Dicks. — *Weisia viridula*.  
Sw. — *Oncophorus*.

*virescens* Kindb. — Br. *veronense*.

*viridans* Wils. (1) { — Br. *inclinatum* var.  
— Br. *intermedium* var. } (*fd. Mitt.*).  
— Br. *pendulum*

*viridatum* C. M. — *Webera*.

*viride* Philib. in *Rev. bryol. 1886, p. 25.*

Syn. — Eur.: Norveg. bor.

*viridescens* Welw. et Duby in *Mém. . . Genève, 1870, p. 4, t. II, f. 2.*

1. In prat. — Afr.: Angola.

*viridissimum* Dicks. — *Zygodon*.

*viridulum* C. M. in *Rev. bryol. 1896, p. 3 (nom.) et in Hedwig, 1898, p. 105.*

*Rhodobryum viridulum* Par. *Suppl. Ind. p. 301* (1900).

2. — Pac.: Austral. or. (Nov. Vales.).

*viridulum* { Dicks. — *Fissidens pusillus*.  
Sm. — *Fissidens exilis*.  
L. — *Weisia*.

*vulcanicolum* C. M. in *Bull. hb. Boissier, 1897, p. 184.*

2. — Am. sept.: And. vulc. Guatemat.

*vulcanicum* Dusen M. Am. austr. n. 876.

Terr. — Am. merid.: Chile austr.

(1) Cl. Mitt. (M. Ind. or. p. 71) has tres spec. ad unam, contra fere omn. bryolog. sententiam, reducens, cui earum nom. *Br. viridantis* imposuit cl. Wilson mihi latet.



*Wagneri* C. M. *Syn.* II, p. 570.

2. — *Am. merid.* : Columb.

*Wahlenbergii* Schwaegr. — *Mniobryum albicans*.

var. *glaciale* Bryol. eur. — *Mniobryum albicans* var. *glacialis*.

*Wallaceum* C. M. in *Hedwig*. 1898, p. 89.

*Syn.* — *Pac.* : Austral. or. (Victoria).

*warneum* Bland. M. *exs. fasc. IV, n.* 184 (1805); *Bryol. eur. fasc. 6-9* (Vol. IV), *Mon. p.* 20, t. 340 (1839); C. M. *Syn. I, p.* 272 (1849); W. P. Sch. *op. cit. p.* 425.

*Br. caespitium* var. *warneum* Schwaegr. *Suppl. III, P. I, II, p.* 109 (1828).

*Br. flagellare* Brid. *Mant. M. p.* 118 (1819).

*Hypnum caespitium* W. et M. *Bot. Taschb. p.* 285 p.p. (1807).

*Pohlia cirrifera* de Not. *Mant. M. n.* 9.

*P. warnensis* Schwaegr. *Suppl. III, P. I, t.* 236 (1828).

1 rar. *syn.* In udis aren. et limos. — *Eur.* : Angl., Suec., German. sept., Siles., Hungar., Ital. super. As. : Himalaya occid. Afr. : Ins. Kerguelen. Am. sept. : Calif.

*warneum* Mac. — *Br. Biddlecomiae*.

*Warnstorffii* Ruthe in *Hedwig*. 1899, *Beibl. p.* 17.

*Syn.* Ad terr. argill. arenac. — *Eur.* : Pomeran.

*Wattsi* Broth. in *Oefv. af Finska Vet.-Akad. Foerh.* 1899, p. 101.

2. Terr. — *Pac.* : Austral. or. (Nov. Vales.).

*weberaceum* Besch. in *Bull. Soc. bot. Fr.* 1898, p. 67.

*Saxic.* in torrent. — *Pac.* : Taïti.

*Weigeli* Brid. — *Webera Ludwigii*.

*Weisia* { C. M. — *Brachymenium*.

{ Dicks. — *Ditrichum homomallum*.

{ Hoffm. — *Dicranella heteromalla*.

*Welovitschii* Dy. — *Brachymenium*.

*Weslayi* Par. *Ind. p.* 223 (1894).

*Br. origanum* Bosw. in *Lond. Journ. of bot. Febr.* 1880 et in *Rev. bryol* 1880, p. 45.

*Ster.* — *Eur.* : Anglia (Teesdale).

*Wichurae* Broth. in *Hedwig*. 1899, p. 219.

*Rhodobryum Wichurae* Par. *Suppl. Ind. p.* 301 (1900).

2. Ad rivul. in silv. — *As.* : Japon.

*Wightii* Mitt. — *Br. strigosum*.

*Willdenowii* Jur. — *Br. longisetum*.

*Willdii* Broth. in *Oefv. af Finska Vet.-Soc. Foerh.* 1900.

*Plagiobryum Willdii* Par. *Ind. p.* 958 (1896).

2. — *Pac.* : Austral. or. (Queensland).

*Wilmsii* C. M. in *Hedwig*. 1899, p. 74.

*Ster.* — *Afr.* : Transvaal.

*Wilsoni* Mitt. — *Leptobryum*.

*Winkelmanni* Ruthe. — *Br. finmarkicum*.

*Wrightii* Sull. in *Proced. of the Amer. Acad.*... 1859, p. 278.  
 1 vel syn. — *Am. sept.* : Fret. Behring.

*Wrightii* Mitt. — *Leptotheca*.

*Zierni* Dicks. — *Plagiobryum*.

*Zollingeri* Wils. — *Rhodobryum roseum*.

*zonatiforme* Par. *Ind.* p. 223 (1894).

*Br. lonchophyllum* Broth. in *Engler's Bot. Jahrb.* 1897, p. 245.

*Br. zonatum* C. M. in *Dusen M. Camer. n.* 340.

Ster. Ad rup. basalt. humid. — *Afr.* : Cameroon.

*zonatiforme* Kindb. — *Br. dubiosum*.

*zonatum* W. P. Sch. *Syn. II ed.* p. 472 (1876); *Hagen in Det. K. Norske Vid. Sells. Skrift.* 1897, n. 2.

Ster. In humid. — *Eur.* : Norveg. bor.

*zonatum* C. M. — *Br. zonatiforme*.

*zygodontoides* C. M. in *Engler's Bot. Jahrb.* 1883, p. 83 et in *Forschungsreise*.... p. 40 (1889).

2. — *Afr.* : Ins. Ascension.

BRYUM POLLA *Adans.*; *Brid. Bryol. univ. I*, p. 688 (1826).

Spec. omn. non indicatæ sunt MNIA ejusd. nom.

*Auberti* Schwaegr. — *Bryum*.

*binervis* Brid. — *Lepidopilum polytrichoides*.

*Commersonii* Schwaegr. — *Rhodobryum*.

*cruda* Brid. — *Webera*.

*cylindrica* Hsch. — *Mnium* ? (*Cf. Brid. op. cit.* p. 712).

*densifolia* Brid. — *Br. schistoneuron* var. *densifolium*.

*elliptica* Brid. — *Mnium punctatum*.

*erythroculon* Schwaegr. — *Bryum*.

*latifolia* Schwaegr. — *Br. Schleicheri* var. *latifolia*.

*ligulata* Schreb. — *Mnium undulatum*.

*rosea* Brid. — *Rhodobryum*.

var. *dominghensis* Brid. — *Rhodobryum Antillarum*.

γ *gigantea* Schwaegr. — *Rhodobryum giganteum*.

*serrata* Schrad. var. *pensilvanica* Brid. — *Mnium serratum* var. *apiculatum* ?

*spinosa* Voit p. p. { — *Mnium*.

{ — *Mnium affine*.

*subernevis* Schwaegr. — *Eriopus remotifolius*.

*truncorum* Brid. — *Bryum*.

*umbraculum* Bursh. — *Rhodobryum*.

BRYUM WEBERA *Brid. Bryol. univ. I*, p. 264 (1826).

Spec. omn. non indicatæ sunt BRYA ejusd. nom.

*alpina* Brid. — *W. longicolla*.

*bicolor* Brid. — *W. nutans* var. *bicolor*.

*caespitosa* Brid. — *W. nutans* var. *caespitosa*.

*erecta* Brid. — *Brachymenium mexicanum*.

*fasciculata* Brid. — *W. nutans*.

*flexuosa* Brid. — *W. Brideliana*.

*grimsulana* Brid. — *W. longicolla* var. *grimsulana*.

- longicolla* Brid. — *W. longicolla*.  
*longiseta* Thom. — *W. nutans* var. *longiseta*.  
*macrocarpa* Brid. — *W. elongata* var. *macrocarpa*.  
*nitens* Kindb. — *W. cruda* var. *minor*.  
*nutans* Brid. — *Webera*.  
 $\beta$  *minor* Brid. — *Webera nutans* var. *minor*.  
*pendula* Brid. — *W. nutans* var. *subdenticulata*.  
*piriformis* Brid. — *Leptobryum*.  
*sphagnicola* Bryol. eur. — *Webera*.  
*subdenticulata* Brid. — *W. nutans* var. *subdenticulata*.

**BUXBAUMIA** Haller Enum. Stirp. Helv. I, p. 10 (1742). — 5 spec.

*aphylla* L. *Diss. Buxb.* § II, p. 10 et § VIII, p. 15 (1757); *Bryol. eur. fasc. I* (Vol. IV), *Mon.* p. 5, t. 427 (1837) et *Mon. Suppl.* p. 4, t. 641 ( ); *C. M. Syn.* I, p. 151 (1849); *W. P. Sch. Syn.* II ed. p. 519.

*B. caulescens* Schmid. *Diss. Buxb.* p. 25, t. I, f. 1-25 (1758).

*B. caulescens aphylla* Hall. *op. cit.* III, p. 25 (1758).

*B. curiosa* Gray, *Nat. arr. brit. pl.* I, p. 750 (1821).

*B. vulgaris* Brid. *Bryol. univ.* I, p. 329 (1826).

*Hippopodium aphyllum* Fabr. *Prim. fl. butisbac.* p. 31 (1743).

*Muscus capillaceus aphyllus*, capitulo crasso bivalvi *Buxb. Pl. min. cogn. Cent. II*, pl. 8, t. 4, f. 2 (1728); *Dill. Hist. M.* p. 477, n. 5 et p. 554, t. 68, f. 5 (1741) et *hb.*

*Saccophorus aphyllus* P. B. *Prodr.* p. 30 (1805).

2. In ericet. — *Eur.*: E reg. campestr. usq. in alp. per zon. interm.  
*As.*: Volga, Jenissei super., *Amour. Am. sept.*: Nov. Scot., Quebec, Nov. Angl., Massachussets, Columb. brit.

*aphylla* Schwaegr. p. p.  
 var. *indusiata* Wahlenb. } — *B. indusiata*.  
 var. *viridis* Moug. }  
*caulescens* Schmid. } — *B. aphylla*.  
*curiosa* Gray. }  
*foliosa* L. — *Diphyscium*.

*indusiata* Brid. *Bryol. univ.* I, p. 331 (1826) et II, p. 741, t. 2, *Suppl.* f. 1-8 (1827); *Bryol. eur. fasc. I* (Vol. IV), *Mon.* p. 6, t. 428 (1837) et *Mon. Suppl.* p. 5, t. 641 ( ); *C. M. Syn.* I, p. 152 (1849) et II, p. 553 (1851); *W. P. Sch. op. cit.* p. 550.

*B. aphylla* { *Schwaegr. in L. Sp. Pl. V* ed., V, II, p. 24, n. I, p. p.

{  $\beta$  *indusiata* Wahlenb. *Fl. Suec. II*, ed., II, p. 760 (1833).

{  $\beta$  *viridis* Moug. in *DC. Fl. fr. III* ed., V, p. 227 (1815).

*B. viridis* Brid. in Moug. et Nestl. *Stirp. crypt. vog.-rhen. fasc. VIII*, n. 724 (1823).

2. Lignic. — *Eur.*: Ubi praeced.: sporadica. *As.*: Caucas., China centr. *Am. sept.*: Columb. brit.

*javanica* C. M. *Syn.* I, p. 152; *Bryol. jav.* I, p. 34, t. 24 (1).

1. Lignic. et rupic. — *As.*: Java.

(1) Cf. *W. P. Sch. Syn.* II ed. p. 550.

*Piperi* Best in *Bull. Torr. bot. Club*, 1893, p. 116.

2. Lignic. — *Am. sept.* : Washington, Idaho.

*sessilis* Schmid. — *Diphyscium foliosum*.

*tasmanica* Mitt. in *Journ. of the Linn. Soc.* 1859, p. 96.

— *Pac.* : Tasman.

*viridis* }  
*vulgaris* } Brid — *B. aphylla*.

**Cacodon** Lindb. in *Act. Soc. pr. Faun. et Fl. fenn.* 9<sup>bri</sup> 1882 (subgen. POHLIÆ = WRBERA).

**CALLIBRYON** Zenk. et Dietr. *M. Thur. fasc. XI*, n. 41 (1+22).

*polytrichoides* Wieb. — *Atrichum undulatum*.

*undulatum* Z. et D. — *Atrichum*.

**CALLICOSTA** C. M. in *Linn.* 1848, p. 787.

Spec. omn. non indicatae sunt *PILOTRICHA* ejusd. nom.

*evanescens* C. M. — *Pilotrichum bipinnatum*.

*filiformis* C. M. — *Hookeria*.

**CALLICOSTELLA** C. M. *Syn. II*, p. 216 (sect. *HOOKERIÆ*, postea gen. propr.) (1851).

Spec. omn. non indicatae sunt *HOOKERIÆ* ejusd. nom.

*daltonioides* Jaeg. — *Schizomitrium*.

*heterophylla* R. C. — *Hookeria Perrotti*.

*Leprieurii* Jaeg. — *Schizomitrium*.

*obliqua* ej. — *Schizomitrium*.

*obscura* ej. — *Schizomitrium*.

*papillata* Mitt. (Sam. M.) — *Hookeria oblongifolia*.

*viridissima* C. M. — *Hookeria pontianacensis*.

**CALLIERGON** Lesq. et Jam. *Man.* p. 404 (subgen. *HYPNÆ*) (1884); *Kindb. Eur. and N. Am. Bryin.* p. 79 (gen. propr.) (1897).

Spec. omn. non indicatae sunt *HYPNÆ* ejusd. nom.

*nivale* Kindb. — *H. stramineum var. nivale*.

*submolle* ej. — *H. molle*.

**CALOMNION** Hook. f. et Wils. *Fl. of the N. Zeal.* II (1855). — 3 spec.

*denticulatum* Mitt. in *Journ. of the Linn. Soc.* 1868, p. 192.

*Cortic.* — *Pac.* : Ins. Samoa.

*laetum* H. f. et W. *op. cit.* p. 87, f. 5 et *Handb. of the N. Zeal. Fl.* p. 490 (1867).

*Gymnostomum* (Eucladon) *complanatum* *cor.* in *Journ. of bot.* 1814, p. 538;  
*C. M. Syn. I*, p. 190 (1849).

2. Filicic. et lignic. — *Pac.* : Nov. Seeland. sept. et med.

*Nadeaudii* Besch. in *Rev. bryol.* 1898, pp. 42 et 43.

*Nadeaudia schistotegiellacea* ej. in *op. cit.* p. II.

2. Ad rhizom. *Filicum.* — *Pac.* : Taiti.

**CALYMPERES** Sw. in Web. Tab. exh. calyptr. opercul... gen. (1813) und in Jahrb. d. Gewächsk. 1818, p. 1; Besch. Essai sur le g. *Calymperes* in Ann. sc. nat. 1895-96, pp. 247-308. — 196 spec. (1)

*aduncifolium* Besch. in Bull. Soc. bot. Fr. 1898, p. 61.

Ster. Cortic. — Pac.: Taïti.

*aeruginosum* Hpe. mss.; Lac. Sp. nov. M. Archip. ind. p. 7, t. 5 A (1872).

2. Cortic. — As.: Ins. Philipp., Celebes.

*africanum* Par. — Syrrhopodon.

*Afzelii* Sw. in Jahrb. d. Geswächsk. 1818, p. 3; Brid. Bryol. univ. II, p. 86 (1827); C. M. Syn. II, p. 523 (1851).

*Dicranum ustulatum* Brid. op. cit. I, p. 458 (1826).

2. Cortic. — Afr.: Sierra Leone.

{	<i>Afzelii</i>	{	Mitt. (ins. Princip.). — C. occidentale.
			— C. Aongstromii.
			— C. Hookeri.
			— C. Perrottetii.
			— C. Richardi.
{	W. P. Sch. {	— C. Hombroni.	
		— C. Jardini.	

var. *moluccense* Brid. — C. moluccense.

*albo-vaginatatum* W. Arn. — Syrrhopodon.

*andamense* Besch. in hb. British. Mus. et op. cit. p. 276.

— As.: Ins. Andaman.

*andamense* C. M. — Syrrhopodon.

*androgynum* Mont. — Syrrhopodon rigidus.

*Aongstroemii* Besch. Fl. bryol. Taïti, p. 24 (1894).

C. *Afzelii* Mont. in hb. Mus. Paris.

C. *rotundatum* Besch. in Ann. sc. nat. 1894, p. ; Par. Ind. p. 232 (1893).

C. *taïtense* Aongstr. mss.; Lindb. hb.

2. — Pac.: Taïti.

*arcuatum* C. M. in Engler's Bot. Jahrb. 1883, p. 85 et in Erforschungsreise.... Bot. p. 51 (1889).

Ster. Cortic. — Pac.: Nov. Guinea.

*Ascensionis* C. M. in op. cc. pp. 84 et 42.

C. *pulvinatum* Mitt. in Voy. H. S. M. Challenger, 1885, III, p. 42.

Ster. — Afr.: Ins. Ascensionis.

*asperipes* Besch. op. cit. p. 277.

— Am. sept.: Guadalup., Martinic., Dominic.

Am. merid.: Ins. Trinitat.

*asperum* C. M. in Dusen M. Camer. n. 400 et in K. Sv. Vet.-Akad. Foerh. 1895, n. 3.

Ster. Cortic. et lignic. — Afr.: Cameroon.

*asterigerum* ej. in Coll. cit. n. 82 et in op. et l. cc.

2. Muric. — Afr.: Liberia (Monrovia).

(1) De speciebus indo-malay. et (pr. p.) novo-guineens. cf. Fleischer, *Die Musci der Flora von Buitenzorg*. 1, pp. 236-278.

- asteristylum* *ej. in Coll. cit. n. 751 et in op. et l. cc.*  
 Ster. Ad ramul. — *Afr.* : Cameroon.
- Aurantium* *Hpe. — Syrrhopodon croceus.*
- australe* *Besch. in op. cit. p. 277.*  
 Cortic. — *Pac.* : Ins. Kermadec, Raoul.
- bahiense* *C. M. in Hedwig. 1900, p. 261.*  
 — *Am. merid.* : Brasil. or. (Bahia).
- Beccarii* *Hpe. in Nuov. Giorn. bot. ital. 1872, p. 279 (1).*  
 — *As.* : Horneo.
- Bescherellei* *Fleisch. M. Archip. ind. n. 65 (1899).*  
 2. Cortic. — *As.* : Sumatra, Java, Singapore.
- Bodeni* *C. M. in Hedwig. 1900, p. 262.*  
 Ster. — *Am. merid.* : Brasil. or. (Bahia).
- Borgenii* *Kiaër in Wright M. of Mad. in Journ. of bot. 1883; Besch. in op. et l. cc.*  
*C. Richardi* *Hb. Mont.*  
 2. — *Afr.* : Madagascar.
- *var. congolense* *Ren. et Card. in Bull. Soc. roy. bot. Belg. 1896, I, p. 311.*  
 Cortic. — *Afr.* : Congo.
- Boulayi* *Besch. in op. cit. p. 278.*  
*C. Dozyanum* *Mitt. (fid. Fleisch.).*  
*C. moluccense* *Hb. Mus. Lugdun. Batav.*  
*C. Palisoti* *Hb. Mus. Paris.*  
 Cortic. — *As.* : Borneo, Singapore, Java.
- brachypelma* *C. M. in Dus. M. Camer. n. 85 et in K. Sv. Vet.-Akad. Foerh. 1895, n. 3.*  
 Cortic. — *Afr.* : Liberia, Cameroon.
- Breutelii* *Besch. in op. et l. cc.*  
 — *Am. sept.* : Ins. S. Thomas.
- Brittoniae* *Besch. in op. et l. cc.*  
 — *Am. sept.* : Florida.
- Brotheri* *Besch. in op. cit. p. 279.*  
*C. moluccense* *Broth. in Micholitz M. Nov. Guin. n. 22.*  
 — *Pac.* : Nov. Guinea (Skrae, pr. Mac Cloer Golf).
- bryaceum* *Besch. in op. et l. cc. et in Thwaites M. Ceylan, n. 62 p. p.*  
 — *As.* : Ceylan.
- burmense* *Hpe. hb. n. 4436.*  
*C. crocatum* *ej. eod. loco.*  
 — *As.* : M. Barma (Moulmein).
- cacazouense* *Besch. in op. cit. p. 280.*  
 — *Afr.* : Ins. Cacazou, Mayotte.
- campylopodioides* *C. M. in Dus. M. Camer. n. 151 et in K. Sv. Vet.-Akad. Foerh. 1895, n. 3.*  
 Ster. Ad rup. littor. — *Afr.* : Cameroon.

(1) *Berterii* *Spreng. Neue Entdeck. III, p. I (1820-22); C. M. Syn. I, p. 528.*  
 — Deleatur.

Cardoti (*Fleisch.*) *Par.*

Thyridium Cardoti *Fl. M. Archip. ind. n.* 73 (1899).

Cortic. — *As.* : Java.

Carionis *C. M. in Bull. hb. Boissier*, 1897, p. 189.

— *Am. sept.* : Guatemala.

caudatum *C. M. in Fl.* 1879, n. 24.

Ster. — *Afr.* : Monbaza.

cernense *Mitt. mss.*; *Besch. in op. cit.* p. 268, n. 105.

— *Afr.* : Madagascar.

ceylanicum *Besch. in op. et l. cc.*

*C. Kurzianum* *Hpe. in hb. Melbourne*, n. 313.

*C. porrectum* *Mitt. in Thwaites M. Ceylan*, n. 60.

*C. salakense* var. ? (*fid. Fleisch.*)

— *As.* : Ceylan, Java (m. Salak).

chamaeleonteum *C. M. in Engler's Bot. Jahrb.* 1883, p. 86.

— *Pac.* : Nov. Guinea.

« var. nanum *ej. in op. et l. cc.*

Cortic. — *Pac.* : Nov. Hannoveria.

Chamissonis *Besch. in op. et l. cc.*

*C. Palisoti de Chamisso in hb. Schwaeagr.*

— *Pac.* : Ins. Corall. (pr. ins. Marquis.).

chlorosum *Hpe. Enum. M. Brasil.* p. 6 (1879).

Ster. Cortic. et lignic. — *Am. merid.* : Brasil. austr. or.

chrysoblastum *C. M. in Engler's Bot. Jahrb.* 1883, p. 88 et in *Forschungsreise.... Bot.* p. 63 (1889).

Ster. Ad terr. humid. — *Afr.* : Monrovia.

constrictum *Sull.* — Syrrhopodon.

contractum *Besch. in op. et l. cc.*

— *As.* : Ceylan.

couguiense *Besch. Fl. bryol. Nouv. Caled.* p. 206 (1873).

2 ? Cortic. — *Pac.* : Nov. Caled.

crassilimbatum *Ren. et Card. in Bull. Soc. roy. bot. Belg.* 1893, 11, p. 18.

Ster. — *Afr.* : Ins. Borbon., Madagasc., ins. S. Marie.

crassinerve *Jaeg.* — Syrrhopodon.

? crispum *Aust.* — Syrrhopodon.

cristatum *Hpe. in Nuov. Giorm. bot. ital.* p. 278.

2. Cortic — *As.* : Borneo (Sarawak), Java.

crocatum *Hpe.* — *C. burmense.*

croceum *Mitt.* (ins. Samoa). — Syrrhopodon subcroceus.

Crügeri *C. M. Syn. I*, p. 527 (1849); *Mitt. M. austr. am.* p. 128 (1869).

2. — *Am. sept.* : Guadalup. *Am. merid.* : Ins. Trititat.

cuspidatum *Besch. mss.*

Thyridium cuspidatum *Fleisch. M. Archip. ind. n.* 72 (1899).

2. Ster. — *As.* : Java.

*cymbifolium* C. M. *mss.*; *Besch. in op. cit. p. 264, n. 4.*

— *Pac.* : Nov. Guinea (Stephensort).

*Daemelii* Hpe. — *Syrrhopodon taiitensis.*

*decolorans* C. M. in *Besch. Fl. bryol. Réunion, etc. p. 58 (1880-81).*

-- *Afr.* : Madagasc., ins. Mayotte, S. Marie.

*Delesserti* *Besch. in Ann. sc. nat. 1895-96, I, p. 282.*

C. Dozyanum *Kurz M. ins. Andaman.*

C. Richardi *Mont. hb.*

— *As.* : Ins. Andaman, Poulo Pinang.

*denticulatum* C. M. in *Engler's Bot. Jahrb. 1883, p. 86 et in Erforschungsreise.... Bot., p. 52 (1889)*; *Geheeb in Bibl. bot. 1898, p. 12, t. VII B.*

C. orientale (*fd. Fleisch.*).

2. Cortic. — *Pac.* : Nov. Guinea.

*dilatatum* C. M. in *Hildebr. M. Mad.*

— *Afr.* : Ins. Nossi Be.

*disciforme* C. M. in *Linn. 1848, p. 183 et Syn. I, p. 525 (1849)*; *Mitt. M. austr. am. p. 126 (1869).*

C. Richardi *Aust. M. Appal. n. 489.*

Cortic. — *Am. sept.* : Martinic. *Am. merid.* : Ins. Trinitat., Surinam, Cayenne.

*disciforme* Lesq. et Jam. — *Syrrhopodon Smithii (fd. Besch. in litt. 1896).*

*disjunctum* *Besch. in op. et l. cc.*

Cortic. — *Afr.* : Ins. Nossi-Be.

? *dissitifolium* Broth. in *Micholitz M. Nov. Guin.*

— *Pac.* : Skraë, pr. Mac Cloer Golf.

*Donnelli* *Aust. in Bot. Gaz. IV, p. 151.*

C. Richardi *var. Donnellii Lesq. et Jam. Man. p. 184 (1884).*

— *Am. sept.* : Florid.

*Dozyanum* *Mitt. M. Ind. or. p. 42 (1859).*

C. moluccense *Bryol. jav. I, p. 47, t. 38 (fd. Mitt.).*

2. Cortic. — *As.* : Ceylan, Sumatra, Java, Batavia, Singapore, Borneo.

<i>Dozyanum</i>	{	Besch. (non Mitt. ! <i>fd. Fleisch.</i> ). — C. tenerum.
		Broth. — C. flaviusculum.
		M. Griffith. — C. malayanum.
		M. Kurz. { — C. Delesserti.
		{ — (n. 1650) — C. linguatum.
		M. Powell. — C. samoanum.
	{	Thwaites n. 62 p.p. { — C. C. bryaceum.
		{ — C. Mittenii.
		{ — C. Thwaitesii.

*Dusenii* C. M. in *Dus. M. Camer. n. 917.*

— *Afr.* : Cameroon.

*emersum* C. M. in *Bull. hb. Boissier, 1897, p. 189.*

— *Am. sept.* : Guatemala.

*erosum* C. M. in *Linn. 1848, p. 182 et Syn. I, p. 524 (1849)*; *Mitt. M. austr. am. p. 127 (1869).*

2. Cortic. — *Am. merid.* : Surinam, Cayenne, Amazon.



*erosum* Mitt. — C. Sprucei.

*eutrichostomum* C. M. mss.; *Besch. in op. cit.* p. 265, n. 40.

C. Dozyanum (*fid. Fleisch.*).

— As. : Singapore.

*exlimbatum* C. M. — C. Manii.

*fasciculatum* *Bryol. jav. I*, p. 50, t. 41 (1855-61).

2. Cortic. — As. : Java.

*fasciculatum* Mitt. { (M. Ind. or.) — Syrrhopodon.  
{ (M. Thwaites) — C. Leucoloma.

var. *heterophyllum* Lac. — C. salakense.

*flaviusculum* Broth. in *Micholitz M. Nov. Guin.*

C. Dozyanum (*fid. Fleisch.*).

— Pac. : Ins. Great Natunas.

*Fordii* *Besch. in op. cit.* p. 284.

— As. : Hong-Kong.

« var. *javanica* *Fleisch. in litt.*

— As. : Java.

*Garciae* Mitt. in *Journ. of the Linn. Soc. XXII*, p. 338.

— Afr. : Ins. Diego-Garcia.

*Gardneri* Hook. — Syrrhopodon.

*Geppii* *Besch. in op. et l. cc.*

C. Palisoti *Wils. hb. n.* 1258.

2. Cortic. — As. : Java.

*Gilleti* *Ren. et Card. in Bull. Soc. roy. bot. Belg.* 1900, II, p. 109.

— Afr. : Congo belg.

*Glaziowii* *Hpe. Symb.* 1877, p. 716.

Ster.

— Am. merid. : Brasil. austr. or.

*Graeffeanum* C. M. M. *Polyn. p.* 64 (1874).

2. Cortic. — Pac. : Ins. Samoa.

*Griffithii* C. M. mss.; *Besch. op. cit.* p. 269, n. 113.

— As. : Nepal.

*guadalupense* *Besch. in op. cit.* p. 85.

— Am. sept. : Guadalup.

*Guiltingii* Hook. et Grev. in *Brewster Edinb. Journ. III*, p. 223 (1824) ;

C. M. *Syn. I*, p. 528 (1849) Mitt. *M. austr. am. p.* 128 (1869).

Cortic. — Am. sept. : Ins. S. Vincent., Guadalup., Martinic., Dominic.

*Hampei* *Bryol. jav. I*, p. 48, t. 39 (1855-61) et II, p. 223 (1861-70).

2. Cortic. et rupic. — As. : Banca ? Java, Sumatra, ins. Malay. Pac. : Timor ?

*Hampei* C. M. — C. Naumanni.

*heterophyllum* (Mitt.) *Besch. in op. et l. cc.*

Syrrhopodon *heterophyllum* Mitt. *M. Ind. or. p.* 40 (1859).

S. serrulatus *Wils. hb.*

— As. : Ceylan.

*heterophyllum* { A. Br. — Syrrhopodon *erocaeus*.  
                           { Hpe. } (Bryol. jav.). — *C. salakense*.  
                           { (Mus. Lugdun. batav.). — *C. scalare*.

*Heudeloti* Besch. in op. et l. cc.

Ad *Elaim guineensem*. — Afr. : Senegamb.

*hexagonum* Besch. in op. et l. cc.

*C. Palisoti* Rich.

Ad saxa humid. silvul. et torr. — Am. sept. : Ins. S. Thoma.

*Hildebrandti* C. M. in Linn. 1876, p. 246.

Ster. Cortic. — Afr. : Ins. Anjouan.

*hispidum* Ren. et Card. in Bull. Soc. roy. bot. Belg. 1893, II, p. 14.

*C. Mariei* Ren. M. M. M. n. 17.

2. Afr. : Madagasc., ins. S. Maria.

*Hobsoni* Grev. — Syrrhopodon.

*Hombroni* Besch. in op. cit. p. 287.

*C. Afzelii* W. P. Sch. in hb. Mus. Paris.

— Pac. : Archip. Gambier.

*Hookeri* Besch. in op. et l. cc.

*C. Afzelii* hb. Hook. et hb. Mont.

— Am. sept. : Ins. Antigoa.

*hyalinoblastum* C. M. mss.; Besch. in op. cit. p. 265, n. 45.

*C. cymbifolium* (fid. Fleisch.).

— Pac. : Nov. Guinea (ins. Mola).

*hyophilaceum* C. M. mss.; Besch. in op. cit. p. 265, n. 32.

2. Cortic. — As. : Ins. Philippin., Java, Sumatra.

« var. *timorense* Besch. in op. et l. cc.

*C. moluccense* hb. Geheeb.

— As. : Timor Laut.

*inaequifolium* C. M. in Bot. Zeit. 1834, p. 348.

Ster. — As. : Poulo-Pinang.

*incurvatum* C. M. M. Polyn. p. 64 (1874).

Ster. — As. : Ins. Samoa.

*integrifolium* C. M. in Fl. 1886, p. 514.

Cortic. — Afr. : Kiulu.

*intra-limbatum* C. M. in op. cit. p. 513.

2. Muscic. — Afr. : Mossamedes.

*involutum* W. P. Sch. — Trichostomum microcarpum.

*Isleanum* Besch. Fl. bryol. Réunion, etc. p. 56 (1880-81).

2. Cortic. — Afr. : Ins. Seychell., Nossi-Be.

*japonicum* Besch. in Journ. de bot. 1898, p.

Ster. — As. : Japon.

« var. *rufescens* Broth. in Hedwig. 1899, p. 211.

Ubi typ.

- Jardini *Besch. in Ann. sc. nat.* 1895-96, p. 287.  
*C. Afzelii* W. P. Sch. in *Jardin Enum.*  
 Cortic. — *Afr.* : Senegamb. (ins. Loss).
- Kaërbachii *Broth. in Engler's Bot. Jahrb.* 1893, p. 477.  
 Ster. — *Pac.* : Nov. Guinea.
- \* *var. timorenses* *Besch. in op. et l. cc.*  
 — *As.* : Timor Laut.
- Kennedyanum *Hpe. in Linn.* 1876, p. 303.  
 Ster. — *Pac.* : Austral. or. (Buckingham Bay).
- Kurzianum *Hpe. in Fl.* 1878, n. 6.  
 — *As.* : Ins. Andaman.
- Kurzianum* *Hpe. hb.* { (n. 313). — *C. ceylanicum.*  
 { (n. 2550). — *C. linguatum.*
- laevifolium *Mitt. in Phil. Trans. of the royal. Soc.* 1879 (*extr.-vol.*),  
 p. 388.  
 — *Afr.* : Ins. Rodriguez.
- lanceolatum *Hpe. Enum. M. Brasil.* p. 6 (1879).  
 Ster. — *Am. merid.* : Brasil. austr. or.
- latifolium *Hpe. in Pl. Preiss. n.* 116; *C. M. Syn. I*, p. 525 (1849)  
 2? — *Pac.* : Austral. occ. et or.
- Lecomtei *Besch. in Journ. de bot.* 1895, n. 12.  
 — *Afr.* : Congo gall. (Kitabi).
- leucocoleos *C. M. in Dus. M. Camer. n.* 76 et in *K. Sv. Vet.-Akad.*  
*Foerh.* 1895, n. 3.  
 2. *Ad Raphiam vinifer.* — *Afr.* : Cameroon.
- Leucoloma *Besch. in Ann. sc. nat.* 1895-96, p. 289.  
*C. fasciculatum* *Thwaites in M. Ceylan*, n. 59.  
 — *As.* : Ceylan.
- leucomitrium *C. M. in Fl.* 1886, p. 512.  
 2. Cortic. — *Afr.* : Niger, Cameroon.
- Levyianum *Besch. in op. cit.* p. 290.  
 2. ♂ ign. — *Am. sept.* : Nicaragua.
- ligulare *Mitt. in Journ. of the Linn. Soc.* 1886, p. 303.  
 — *Afr.* : M. Usagara.
- ligulina *C. M. in Micholitz M. Nov. Guineens. n.* 196.  
*C. moluccense* ? (*fid. Fleisch.*).  
 — *Pac.* : Archip. Duc d'York.
- Lindigii* *Hpe.* — *Streptopogon Hampeanus.*
- Lindmanni *Broth. in Bihang till K. Sv. Vet.-Akad. Foerh.* 1900, p.  
 2. Saxic. — *Am. merid.* : Brasil. centr. (Matto Grosso).
- linealifolium *C. M. in Dus. M. Camer. n.* 756 et in *K. Sv. Vet.-Akad.*  
*Foerh.* 1895, n. 3.  
 Ster. *Ad rup. marit.* — *Afr.* : Cameroon.
- linearifolium *C. M. M. Polyn. p.* 65 (1874).  
 Ster. — *Pac.* : Ins. Samoa? Fidji, Tonga.

linguatum *C. M. mss.*

*C. Dozyanum* *Hpe. hb. n.* 1650.

*C. Kurzianum* *ej. ibid. n.* 2550.

*C. moluccense* *Kurz M. ins. Andaman.*

— *As.* : *Ins. Andaman, Mangrove Bny.*

lonchophyllum *Schwaegr. Suppl. I, P. II, p.* 133, *t.* 98 (1816); *Brid. Bryol. univ. II, p.* 88, *excl. syn.* *C. Palisoti* (1827); *C. M. Syn. I, p.* 526 (1849); *Mitt. M. austr. am. p.* 129 (1869).

2. *Cortic. et ad rup. calcar.* — *Am. sept.* : *Guadalup. Am. merid.* : *Venezuela, Guian. gall.*

longifolium *Mitt. in Journ. of the Linn. Soc.* 1868, *p.* 173.

*C. loreum* *Sande Lac. M. nov. Archip. ind. p.* 7 (1872).

— *As.* : *Borneo (Labuan).*

longifolium *Broth.* — *C. thomeanum.*

loreum *Lac.* — *C. longifolium.*

lorifolium *Mitt. Sam. M. p.* 173 (1868).

2? *Filicic.* — *As.* : *Borneo? Pac.* : *Ins. Fidji, Samoa.*

lorifolium var. *neo caledonicum* *Besch.* — *C. Novae Caledoniae.*

loucoubense *Besch. in op. et l. cc.*

— *Afr.* : *Ins. Nossi Be.*

malayanum *Besch. in op. cit. p.* 291.

*C. Dozyanum* *Mitt. in hb. Kew.*

— *As.* : *Malacca.*

Malimbae *C. M. in Fl.* 1886, *p.* 513.

2. — *Afr.* : *Gabon.*

mamosum *Besch. in op. et l. cc.*

— *As.* : *Ins. Philippin.*

Manii *C. M. mss.*

*C. exlimbatum* *ej. prius.*

*Thyridium Manii* *Fleisch. M. Archip. ind. n.* 71 (1899).

*In arena marit.* — *As.* : *Ins. Andaman (Pytler's Ghat), Singapore.*

Mariei *Besch. Fl. bryol. Réunion, etc. p.* 57 (1880-81).

— *Ins. Nossi Be.*

Mariei *Ren.* — *C. hispidum.*

mayottense *Besch. in Ann. sc. nat.* 1895-96, *p.* 292.

— *Afr.* : *Ins. Mayotte.*

megamitrium *C. M. in Dus. M. Camer. n.* 495 *et in K. Sv. Vet.-Akad. Foerh.* 1895, *n.* 3.

2? *Ad Palmas.* — *Afr.* : *Cameroon, Dahomey.*

Melinoni *C. M. mss.*; *Besch. in op. et l. cc.*

*C. Richardi* *hb. Mont. pr. p.*

— *Am. merid.* : *Guian. gall.*

menadense *Besch. in op. et l. cc.*

*C. Dozyanum* (*fid. Fleisch.*).

*C. moluccense* *Hb. Dozy.*

— *As.* : *Celebes (Minetrage van Menado, Amboine).*

microblastum *C. M.* in *Dus. M. Camer. n.* 807 et in *K. Sv. Vet.-Akad. Foerh.* 1895, n. 3.

2. Cortic. — *Afr.* : Cameroon.

minus *Besch. in op. cit. p.* 293.

— *Afr.* : Nossi Comba.

minutifolium *C. M.* in *Micholitz M. Nov. Guineens. n.* 199.

— *Pac.* : Nov. Britann.

Mittenii *Besch. in op. et l. cc.*

*C. Dozyanum Mitt. in M. Thwaites n.* 62 p.p.

— *As.* : Ceylan.

moluccense *Schwaegr. Suppl. II, P. II, p.* 99, t. 127 (1826); *C. M. Syn. I, p.* 523 (1849); *Bryol. jav. I, p.* 47, t. 37 (1855-61); *Besch. in op. et l. cc.*

*C. Afzelii* β moluccense *Brid. Bryol. univ. II, p.* 87 (1827).

*C. Palisoti Gaudich. n.* 15 (29) in *hb. Mus. Paris.*

2. Lignic. — *As.* : Ins. Rauwack Molucc., Mindanao.

<i>moluccense</i>	{	Broth. — <i>C. Brotheri.</i>
		Dz. et Mk. — <i>C. menadense.</i>
		Hb. Geheeb. } — <i>C. hyophilaceum var. timorens.</i>
		} — <i>C. Visheri.</i>
		Hpe. — <i>C. peguense.</i>
		Kurz. — <i>C. linguatum.</i>
		Hb. Lugdun. Batav. } — <i>C. platycinclis.</i>
		} — <i>C. Vriesei.</i>
		Mitt. — <i>C. fasciculatum.</i>
		Sande Lac. — <i>C. sandeanum.</i>
		var. <i>heterophyllum</i> Par. — <i>C. salakense.</i>

Motleyi *Mitt. in Bryol. jav. I, p.* 48, t. 38 (1855-61).

2. — *As.* : Ins. Labuan, Borneo, Ceylan.

Mülleri *Mitt.* — *Syrrophodon.*

nanum *C. M. in Linn.* 1875, p. 398.

Ster. — *Afr.* : Monbottou.

Naumannii *Besch. in op. cit. p.* 294.

*C. Dozyanum (fid. Fleisch.).*

*C. Hampei C. M. in Engler's Bot. Jahrb.* 1883.

— *As.* : Ins. Timor.

nicaraguense *Ren. et Card. in Bull. Soc. roy. bot. Belg.* 1894, II, p. 117.

2. — *Am. sept.* : Nicaragua.

nicobarense *Hpe. hb.* ; *Besch. in op. cit. p.* 295.

— *As.* : Ins. Nicobar.

Nietneri *C. M. in Bot. Zeit.* 1864, p. 348.

2. Cortic. — *As.* : Ceylan, Java.

nigrescens *Broth. et Geh. in Oefv. af Finska Vet.-Akad. Foerh.* 1899, p. 97.

Ster. — *Pac.* : Austral. or. (Queensland).

Nossi Combae *Besch. Fl. bryol. Réunion, etc. p.* 57 (1880-81).

Ster. — *Afr.* : Ins. Nossi Comba.

*Novae Caledoniae Besch. in op. et l. cc.*

*C. lorifolium var. Neo-Caledonicum ej. Fl. bryol. Nouv. Caled. p. 207 (1873); Par. Ind. p. 230 (1893).*

— *Pac.* : Nov. Caledon.

*nukahivense Besch. op. et l. cc.*

— *Pac.* : Ins. Marquis.

*obliquatum C. M. M. Polyn. p. 65 (1874).*

*C. Dozyanum Mitt. Sam. M. p. 172 (1868).*

*Ster. Cortic. — Pac.* : Ins. Admiral., Fidji, Samoa, Tonga.

*occidentale Besch. in op. et l. cc.*

*C. Afzelii Mitt. in Journ. of the Linn. Soc. 1883.*

— *Afr.* : Ins. Princip.

*omanicum Besch. in op. et l. cc.*

— *As.* : Ins. Laquediv.

*orientale Mitt. mss.; Besch. in op. cit. p. 272, n. 159.*

2. *Rupic. — As.* : Borneo, Labuan, Java.

*orthophyllaceum C. M. in Abhandl. bot. Ver. d. Prov. Brandenb. XXXI, p. 67.*

— *Afr.* : Gabon.

*pachyloma Hpe. in Linn. 1876, p. 247.*

*Ster. — Afr.* : Ins. Anjouan.

*pachyneuron C. M. M. Polyn. p. 65 (1874).*

*Ster. — Afr.* : Ins. Samoa, Tonga.

*Palisoti Schwaegr. Suppl. I, sect. poster. p. 334, p.p. (1811); Besch. in op. cit. p. 296.*

*Cryphaea vaginans P. B. mss.*

— *Afr.* : Oware.

<i>Palisoti</i>	{	Dz. et Mk. — C. Boulayi.
		Schwaegr. p.p. } — C. Chamissonis.
		Wils. } — C. platycinclis.
		Wils. } — C. Geppii.
		— C. Hampei.
		— C. punctulatum.

*pallidum Mitt. in Phil. Trans of the roy. Soc. 1879 (extra-vol.), p. 383.*

*Ster. — Afr.* : Ins. Rodriguez.

*palmicola Besch. in op. cit. p. 297.*

— *Afr.* : Ins. Nossi Comba.

*Panamae Besch. in op. cit. p. 298.*

— *Am. sept.* : Panama.

*Pandani C. M. in Engler's Bot. Jahrb. 1883, p. 87 et in Forschungsreise... Bot. p. 58 (1889).*

*Ad Pandanos. — As.* : Amboine.

*panduraefolium Broth. in Oefv. af Finska Vet.-Akad. Foerh. 1898, p. 165.*

*C. Motleyi (fid. Fleisch.).*

2. — *Pac.* : Fret. Torres (prom. York).

*parasiticum Hook. et Grev. — Syrrhopodon.*

- peguense *Besch. in op. cit. p. 299.*  
*C. moluccense Hpe. hb. n. 2928 b.*  
 — *As. : Pegu.*
- perinvolutum *C. M. in Fl. 1897, p. 232.*  
*Ster. — Am. merid. : Venezuela.*
- Perrottetii *Besch. in op. et l. cc.*  
*C. Afzelii Hb. Mont.*  
*Cortic. — Afr. : Senegamb.*
- Pintasii *C. M. mss.; Besch. in op. cit. p. 299.*  
*C. thomeanum Pintas in hb. C. M.*  
 — *Afr. : Ins. Sao Thome.*
- platycinclis *Besch. in op. et l. cc.*  
*C. moluccense Hb. Lugdun. Batav.*  
*C. moluccense var. (fid. Fleisch.).*  
*C. Palisoti Hb. Schwaegr.*  
 2. *Cortic. — As. Java.*
- platyloma *Mitt. M. austr. an. p. 128 (1869).*  
 — *Am. merid. : Guian brit.*
- Polii *Besch. in op. et l. cc.*  
 — *Afr. : Nossi Comba.*
- porrectum *Mitt. Sam. M. p. 172 (1868).*  
*Ster. Lignic. — Pac. : Ins. Samoa.*
- Principis *Broth. in Boll. du Soc. Broter. VIII, 1890, p.*  
 2. *fr. ign. Cortic. — Afr. : Ins. Princip.*
- prionotum *Besch. in op. cit. p. 300.*  
*Syrhophodon Kurzii Hpe. mss.*  
 — *As. : Barma (Nathung).*
- proligerum *C. M. in K. Sv. Vet.-Akad. Foerh. 1895, n. 3.*  
*Ster. Muric. — Afr. : Liberia (Monrovia).*
- pulvinatum *Mitt. — C. Ascensionis.*
- punctulatum *Hpe. in M. Kurz. n. 3961 et 3863.*  
*C. Palisoti Wils. in M. Wallich. n. 1250.*  
 — *As. : Ins. Nicobar, Singapore.*
- pungens *C. M. in Engler's Bot. Jahrb. 1883, p. 87 et in Forschungsreise.... Bot. p. 57 (1889).*  
*Ster. Cortic. — As. : Amboine.*
- Quintasii *Broth. in Boll. da Soc. Broter. VIII, 1890.*  
 2. *Cortic. — Afr. : Ins. Sao Thoma.*
- Rabenhorsti *Hpe. et C. M. in Fl. 1886, p. 512.*  
 2. — *Afr. : Sinus Guineens.*
- recurvifolium (*Wils.*) *Besch. in op. cit. p. 301.*  
*Syrhophodon Campylopus C. M. in Engler's Bot. Jahrb. 1883, p. 86 et in Forschungsreise.... Bot. p. 53 (1889); Par. Ind. p. 1245 (1900).*  
*S. recurvifolius Wils. hb. n. 3823 et in M. Wallich. n. 48.*  
*Terr. — As. : Singapore. Pac. : Archip. d'Entrecasteaux (ins. For-*  
*gusson).*

reduncum *Kiaër* in *M. Borgens.*; *Besch. in op. et l. cc.*  
— *Afr.* : Madagasc.

*repens* C. M. — *Syrrophodon*.

retusum *Besch. in op. et l. cc.*

*Syrrophodon heterophyllum Bryol. jav. Suppl. p. 223.*

— *As.* : Ins. Banca.

*rhypariophyllum* C. M. in *Fl.* 1886, p. 512.

2. *Cortic.* — *Afr.* : Niger.

*Richardi* C. M. *Syn. I*, p. 524 (1849) et *II*, p. 620 (1851); *Mitt. M. austr. am.* p. 127 (1869).

C. *Afzelii* *Hb. Mont. p.p.*

C. *Donnelli* *Aust. hb.*

2. *Cortic.* -- *Am. sept.* : Florid. *Am. merid.* : Guian. gall., Amazon.

*Richardi* { *Aust.* — C. *disciforme*.  
                  { *Mont.* } — C. *Borgenii*.  
                  {           } — C. *Delesserti*.  
                  {           } — C. *Melinoni*.

var. *Donnelli* Lesq. et Jam. — C. *Donnelli*.

*Robillardii* *Besch. in op. cit. p. 302.*

— *Afr.* : Ins. Franciae.

*rotundatum* C. M. in *Dus. M. Camer. n. 53* et in *K. Sv. Vet.-Akad. Foerh.* 1895, n. 3.

2. *Cortic.* — *Afr.* : Cameroon.

*rotundatum* *Besch.* — C. *Aongstroemii*.

*rufescens* *Besch. in op. et l. cc.*

*Ad Cassiam officin.* — *Am. sept.* : Antill.

*rupestre* C. M. in *Dus. M. Camer. n. 53* et in *K. Sv. Vet.-Akad. Foerh.* 1895, n. 3.

*Ster.* *Ad rup. inadid.* — *Afr.* : Cameroon.

*salakense* *Besch. in op. et l. cc.*

C. *fasciculatum* var. *heterophyllum Bryol. jav. II*, p. 223 (1861-70).

C. *heterophyllum* *Hpe. in Bryol. jav. l. c.*

C. *moluccense* var. *heterophyllum Par. Ind. p. 231* (1893).

C. *porrectum* ? (*fid. Fleisch.*).

2. *Cortic.* — *As.* : Java (m. Salak), Singapore, ins. Philippin.

*samoanum* *Besch. in op. et l. cc.*

C. *Dozyanum* *Mitt. Sam. M. p. 172* (1868).

— *Pac.* : Ins. Samoa.

*Sanctae Mariae* *Besch. Fl. bryol. Réunion, etc. p. 58* (1880-81).

*Ster.* — *Afr.* : Madagasc., ins. S. Maria.

« var. *mayottensis* *Besch. in Ann. sc. nat. VII ser. II*, p. 89.

*Ster.* — *Afr.* : Ins. Mayotte.

*Sandeanum* *Besch. in op. cit. p. 303.*

C. *Hampei* (*fid. Fleisch.*).

C. *moluccense* *Sande Lac. hb.*

— *As.* : Borneo.

*saxatile* C. M. in *Dus. M. Camer. n. 94* et in *K. Sv. Vet.-Akad. Foerh.* 1895, n. 3.

*Ster. Rupic.* — *Afr.* : Liberia (Monrovia).



scaberrimum *Broth. in Oefv. af Finska Vet.-Akad. Foerh.* 1893.

*C. subchamaeleonteum C. M. mss.*

2. *Cortic. — Pac. : Nov. Guinea.*

scalare *Besch. in op. et l. cc.*

*C. heterophyllum Hpe. mss.*

*C. salakense (fid. Fleisch.).*

— *As. : Ins. Basilan Philippin.*

secundulum *C. M. in Dus. M. Camer. n. 154 et in K. Sv. Vet.-Akad. Foerh.* 1895, n. 3.

2. *Ad Elain guineens. et rup. marit. — Afr. : Côte d'Ivoire, Cameroon.*

semi-liberum *Mitt. — Syrrhopodon.*

semi-limbatulum *C. M. in Hedwig. 1900, p. 261.*

*Ster. Cortic. — Am. merid. : Brasil. or. (Nov. Friburg.).*

semi-marginatum *C. M. in Engler's Bot. Jahrb. 1883, p. 87 et in Forschungsreise.... Bot. p. 57 (1889).*

*Ster. Cortic. — As. : Amboine.*

Semperi *Hpe. hb. ; Besch. in op. cit. p. 304.*

— *As. : Philippin.*

serratum *A. Br. in C. M. Syn. I, p. 527 (1849); Mitt. M. Ind. or. p. 41 (1859); Bryol. jav. I, p. 49, t. 40 (1855-61) et II, p. 223 (1861-70).*

1. *Filicic. — As. : Borneo, Labuan, Celebes, Java, Sumatra, Ceylan. Pac. : Ins. Samoa.*

setifolium *Hpe. hb. ; Besch. in op. et l. cc.*

— *As. : Ins. Basilan Philippin.*

setosum *C. M. M. Polyn. p. 64 (1874).*

*Ster.*

— *Pac. : Ins. Samoa.*

Seychellarum *Besch. in op. et l. cc.*

— *Afr. : Ins. Mahe Seychell.*

Sprucei *Besch. in op. et l. cc.*

*C. erosum Mitt. M. austr. am. p. 127 (1869).*

— *Am. merid. : Brasil.*

spurio-denticulatum *C. M. in Micholitz M. Nov. Guineens. n. 201.*

— *Pac. : Nov. Britann.*

stenogaster *Besch. in op. cit. p. 305.*

2? *Cortic. — As. : Java.*

stylophyllum *C. M. in Engler's Bot. Jahrb. 1883, p. 85 et in Forschungsreise.... Bot., p. 51 (1889); Geheeb. in Bibl. bot. 1898, p. 10, t. IV A.*

— *As. : Sumatra (ins. Engano).*

subchamaeleonteum *C. M. — C. scaberrimum.*

subdisciforme *C. M. — Syrrhopodon.*

taitense (*Sull.*) *Mitt. Sam. M. p. 172 (1868).*

*C. Daemelii Hpe. mss.*

*Syrrhopodon taitensis Sull. in U. St. Exp. Expl. Wilkes, 1859, M. p. 6, t. I ; Par. Ind. p. 1256 (1897).*

*Ster. Rupic. et terr. — Pac. : Ins. Fidji, Samoa, Taiti, Nuka Hiva, archip. Gambier, Nov. Hebrid.*

- taitense* Lindb. — *C. Aongstroemii*.  
*tenellifolium* C. M. — Syrrhopodon.  
*tenellum* C. M. in *Dus. M. Camer.* n. 419 et in *K. Sv. Vet.-Akad. Foerh.* 1885, n. 3.  
 Ster. Cortic. — *Afr.* : Cameroon.  
*tenerum* C. M. in *Linn.* 1871-73, p. 174.  
 2. Cortic. — *As.* : Calcutta, Sumatra, Java, Singapore, Saigon.  
 « *var. edamense* *Fleisch. M. Archip. ind.* n. 63 (1899).  
 Cortic. — *As.* : Java.  
*tenuifolium* Sull. — Syrrhopodon.  
*thomeannm* C. M. in *Fl.* 1886, p. 286.  
*C. longifolium* Broth. in *sched.*  
 Ster. Muscic. — *Afr.* : Ins. Sao Thoma.  
*thomeanum* Pintas. — *C. Pintasii*.  
*Thompsoni* Besch. in *op. cit.* p. 306.  
 — *Afr.* : Madagasc..  
*Thwaitesii* Besch. in *op. et l. cc.*  
*C. Dozyanum* Mitt. in *Thwaites M. Ceylan.* n. 62 p.p.  
 — *As.* : Ceylan.  
*Uleanum* Broth. in *Hedwig.* 1895, p. 423.  
 — *Am. merid.* : Brasil. or. (Goyaz) et centr.  
 (Matto-Grosso).  
*usambaricum* Broth. in *Engler's Bot. Jahrb.* 1894, p. 184.  
 — *Afr.* : Usambara.  
*varium* Mitt. in *Burmah its People and Prod. II ed.*, 1883.  
*C. Hampei* ? (*fil. Fleisch.*).  
 — *As.* : Barma (Maulmein).  
*Visheri* Besch. in *op. et l. cc.*  
*C. moluccense* Hb. *Geheeb.*  
 — *As.* : Ins. Banca Neera.  
*Vriesei* Besch. in *op. cit.* p. 307.  
*C. moluccense* Hb. *Lugdun. Batav.*  
 2. Cortic. — *As.* : Celebes (Menahaye van Menado), Java.  
*Welwitschii* Duby. — Syrrhopodon.  
*Wulfschlaegelii* Ltz. *Moosst.* p. 160 (1864).  
 Ster. — *Am. sept.* : Haïti. *Am. merid.* : Guian. batav.  
*zanzibarense* Besch. in *op. et l. cc.*  
 — *Afr.* : Zanzibar.
- CALYMPERIDIUM *Bryol. jav.* I, p. 51 (1855-61).  
*Mülleri* *Bryol. jav.* — Syrrhopodon.  
***Calymperopopsis*** C. M. in *Dus. M. Camer.* (sect. SYRRHOPODONTIS).

**CALYPTOPOGON** Mitt. in Journ. of the Linn. Soc. 1859, p. 78 (sect. STREPTOPOGONIS) et in *Proc. of the Linn. Soc.* 1879 (extra-vol.), p. 33, gen. propr. (1). — 5 spec.

*crispatulus* (C. M.) *Broth. Natürl. Pflanzenfam.* p. 419.

*Barbula crispatula* C. M. in *Hedwig.* 1897, p. 104.

*Streptopogon crispatulus* *Par. Ind.* p. 1240 (1897).

Ster.

— *Am. merid.* : Patagon.

*crispatus* (Hpe.) *Broth. op. et l. cc.*

*Barbula crispata* Hpe. in *Linn.* 1876, p. 304.

*Streptopogon crispatus* *Jaeg. Ad. II*, p. 670 (1877-78) ; *Par. op. et l. cc.*

Cortic. — *Pac.* : Austral. or., Tasman.

*Hookeri* (R. Br.) *Broth. op. et l. cc.*

*Streptopogon Hookeri* R. Br. in *Trans. of the N. Zeal. Inst.* 1897, p. 417 ;

*Par. Suppl. Ind.* p. 313 (1900).

1. Cortic. et rupic. — *Pac.* : Nov. Seeland.

*mnioides* (Schwaegr.) Mitt. in *Proc. of the Linn. Soc.* 1879 (extr. vol.), p. 33.

*Barbula mnioides* Schwaegr. *Suppl. IV*, p. 310 (1842) ; *C. M. Syn. I*, p. 632 (1849).

*Streptopogon mnioides* Mitt. in *Journ. of the Linn. Soc.* 1859, p. 78 ; *Par. Ind.* p. 1240 (1897).

*Tortula mnioides* Hook. f. *Handb. of the N. Zeal. Fl.* p. 420 (1867).

2. Cortic. — *Am. merid.* : And. Quitens., Chile. *Pac.* : Nov. Seeland. sept., Tasman.

*Wilhelmi* (C. M.) *Broth. op. et l. cc.*

*Barbula Wilhelmi* C. M. in *Hedwig.* 1897, p. 101.

*Streptopogon Wilhelmi* *Par. Ind.* p. 1241 (1897).

Cortic. — *Pac.* : Austral. or. (Victoria), Tasman.

**Calypotroporus** Lindb. *M. Scand.* p. 28 (sect. ORTHOTRICH) (1879).

**CALYPTOTHECIUM** Mitt. *M. austr. am.* p. 427 (sect. POROTRICH) (1869) et in *Journ. of the Linn. Soc.* 1868, p. 190 (gen. propr.). — 14 spec.

*acutifolium* (Brid.) *Broth. in litt.*

*Neckera acutifolia* C. M. *Syn. II*, p. 48 (1851) ; *Par. Ind.* p. 842 (1896).

*N. Distichia acutifolia* Brid. *Bryol. univ. II*, p. 48 (1827).

Ster. Ad saxa irror. — *Afr.* : Ins. Borbon.

« *var. antitrichioides* (Besch.) *Par.*

*N. acutifolia var. antitrichioides* Besch. *Fl. bryol. Réunion, etc.* p. 133 (1880-81) ; *Par. op. et l. cc.*

Ubi typ.

*acutum* (Mitt.) *Broth. in litt.*

*Pterobryum acutum* Mitt. in *Trans. and Proc. of the roy. Soc. of Victoria*, 1883, p. 81.

Ster.

— *Pac.* : Austral. or. (Queensland).

---

(1) « *Saemtliche Arten sind dem C. mnioides sehr nahe verwandt und werden wahrscheinlich bei reichlicherem materiale z. T. eingehen.* » (*Broth. Natürl. Pflanzenfam.* p. 419).

*africanum* Mitt. in *Journ. of the Linn. Soc.* 1886, p. 312.

*Trachyloma africanum* Rehm. *M. Afr. austr.* n. 332; *Broth. in Engler's Bot. Jahrb.* 1894, p. 198; *Par. Ind.* p. 1301 (1897).

2. *Cortic.* — *Afr.* : M. Usagara, Kilima N'Djaro, Usambara; Austral. *australinum* (Mitt.) Par.

*Pterobryum australinum* Mitt. in *op. et l. co.*; *Par. Ind.* p. 1043 (1897).

Ster. — *Pac.* : Austral. or. (ins. of Moreton Bay).

*breviusculum* (C. M.) *Broth. in litt.*

*Neckera breviuscula* C. M. in *Dus. M. Camer.* n. 242 et in *K. Sv. Vet.-Akad. Foerh.* 1895, n. 3; *Par. Ind.* p. 814 (1896).

Ster. Ad arbor. — *Afr.* : Cameroon.

*Bufloni* *Broth. in Engler's Bot. Jahrb.* 1895, p. 65.

*Trachyloma Bufloni* Par. *Ind.* p. 1301 (1897).

2. — *Pac.* : Tasman.

*Duseni* (C. M.) *Broth. in Engler's Bot. Jahrb.* 1897, p. 255.

*Garovaglia Duseni* C. M. in *Dus. M. Camer.* n. 289; *Par. Ind.* p. 508 (1895).

2. Ad arbor. — *Afr.* : Cameroon, ins. Fernando Po.

*humile* (Mitt.) *Broth. in litt.*

*Porotrichum humile* Mitt. *M. austr. am.* p. 466 (1869); *Par. Ind.* p. 1014 (1896).

Ster. — *Am. merid.* : And. Quitens.

*longiusculum* (C. M.) *Broth. in litt.*

*Neckera longiuscula* C. M. in *Dus. M. Camer.* n. 241 et in *K. Sv. Vet.-Akad. Foerh.* 1895, n. 3; *Par. Ind.* p. 852 (1896).

Ster. Ad arbor. — *Afr.* : Cameroon.

*philippinense* *Broth. in Warburg, Monsonia*, 1900, I, p. 48.

*Neckera philippinensis* Par. *Suppl. Ind.* p. 255 (1900).

— *As.* : Ins. Philippin. (Mindanao).

*praelongum* Mitt. — *C. taïtense*.

*recurvulum* (C. M.) *Broth. in litt.*

*Trachyloma recurvulum* C. M. in *Bailey Syn. of the Queensland Fl. IV, Suppl.* p. 98 (nom., 1891); *Broth. in Oesfv. af Finska Vet.-Akad. Foerh.* p. 53; *Par. Suppl. Ind.* p. 322 (1900).

2. — *Pac.* : Austral. (Queensland).

*subacutifolium* *Broth. in Engler's Bot. Jahrb.* 1897, p. 254.

*Neckera Brotheri* Par. *Suppl. Ind.* p. 254 (1900).

2. fr. ign. *Cortic.* — *Afr.* : Pondo Land.

*taïtense* (W. P. Sch.) Par.

*C. praelongum* Mitt. *Sam. M.* p. 130 (1868).

*Neckera Eugeniae* Lindb. in *Reichdt. Voy. Novara*, 1870 (nom.).

*N. taïtensis* W. P. Sch. in *hb. Mus. Paris* (1850); *Par. Ind.* p. 860 (1896).

2. Ad arbor. — *Pac.* : Taïti, ins. Fidji, Samoa, Nov. Caledon.

*tumidum* (Dicks.) *Fleisch. M. Arch. ind. exs.* n. 222.

*Meteorium tumidum* Mitt. *M. Ind. or.* p. 86 (1859); *Bryol. jav. II*, p. 85, t. 199 (1861-70); *Par. Ind.* p. 801 (1896).

*Neckera tumida* Dicks. *ms. in Hook. M. ex. in adn.* t. 158 (1820).

*Pilotrichum tumidum* Brid. *Bryol. univ.* II, p. 265 (1827); *C. M. Syn.* II, p. 153 (1851).

2. *Filicic.* — *As.* : Celebes, ins. Sarapaea, Ceram, Sumbawa, M. Java, Sumatra, Ceylan, Madras, Mysore, Nepal. (*fid. C. M.*).

**CAMPTOCHÆTE** Reichdt. in Novara Exped. Bot. I, p. 190 (1870). — 11 spec.

*angustata* (Mitt.) Jaeg. *Ad.* II, p. 214 (1874-75).

*Isothecium angustatum* H. f. *Handb. of the N. Zeal. Fl.* p. 465 (1867).

*Ptilocladus angustatus* Lindb. in *Act. Soc. sc. Fenn.* 1872, p. 278.

*Stereodon angustatus* Mitt. in *Journ. of the Linn. Soc.* 1859, p. 88.

2. — *Pac.* : Nov. Seeland.

*Arbuscula* (Hook.) Jaeg. *op. cit.* p. 213 (1874-75).

*Hookeria Arbuscula* Sm. in *Trans. of the Linn. Soc.* IX, p. 280, t. 23.

*Hypnum Arbuscula* Hook. *M. ex.* II, p. 20, t. 112 (1818-20); *C. M. Syn.* II, p. 228 (1851).

*Isothecium Arbuscula* Brid. *Bryol. univ.* II, p. 372 (1827).

*Pterobryum Arbuscula* Mitt. in *Trans. and Proceed. of the roy. Soc. of Victoria*, 1883, p. 83.

*Ptilocladus Arbuscula* Lindb. in *op. et l. cc.*

*Thamniella* ? *Arbuscula* Besch. *Fl. bryol. Nouv. Calédon.* p. 240 (1873).

2. — *Pac.* : Nov. Seeland., Tasman.

*arbusculans* Broth. in *Dusen M. Am. austr.* n. 359.

*Cortic.* — *Am. merid.* : Chile austr.

*Beckettii* Broth. in *Oefv. af Finska Vet.-Akad. Foerh.* 1899, p. 114.

2. *Cortic.* — *Pac.* : Nov. Seeland.

*dendroclada* C. M. in *Helms M. N. Seeland.* n. 14.

— *Pac.* : Nov. Seeland.

*deflexa* (Wils.) Jaeg. *op. et l. cc.*

*Hypnum deflexum* Wils. in *Stephens. M. Nov. Seeland.* ; *C. M. Syn.* II, p. 680 (1851).

*Isothecium Arbuscula*  $\beta$  *deflexa* Handb. of the N. Zeal. Fl. p. 465 (1867).

*Porotrichum deflexum* Mitt. in *op. et l. cc.*

*Ptilocladus deflexus* Lindb. in *op. et l. cc.*

*Stereodon deflexus* Mitt. in *op. et l. cc.*

2. *Cortic. et terr.* — *Pac.* : Austral. or. (Victoria), Nov. Hebrid., Tasman., ins. Auckland, Campbell.

*excavata* (Tayl.) Jaeg. *op. cit.* p. 214.

*Hypnum excavatum* Tayl. in *Phytolog.* 1844, p. 1095; *C. M. Syn.* II, p. 229.

*Thamnium perpusillum* C. M. in *Rev. bryol.* 1877, p. 43 (nom.).

2 ? — *Pac.* : Austral. or. (ins. Five).

*fruticosa* (Mitt.). *Par. Suppl. Ind.* p. 86 (1900).

*Porotrichum fruticosum* Mitt. in *op. et l. cc.*

— *Pac.* : Tasman.

*gracilis* (Hook. f. et Wils.) *Par. Ind.* p. 234 (1894).

*Hypnum gracilescens* C. M. *Syn.* II, p. 271 (1851).

*H. gracile* H. f. et W. in *Lond. Journ. of bot.* 1844, p. 553 et *Fl. antarct.* I, p. 140 (1844).

*H. micans* eor. *op. cit.* II, p. 421 (1847) ?

*Isothecium gracile* Fl. of the N. Zeal. II, p. 106 (1855) et *Fl. Tasman.* II, p. 205 (1860).

*Microthamnium gracile* Jaeg. *Ad. II*, p. 492 (1876-77).

*Porotrichum gracile* Mitt. *in op. cit.* p. 84.

*Ptilocladus gracilis* Lindb. *in op. et l. cc.*

*Stereodon gracilis* Mitt. *in op. et l. cc.*

2. *Ad rup. madid.* — *Am. merid.* : Chile. *Pac.* : Austral. or. (Nov. Vales., Victoria), Nov. Seeland., Tasman, ins. Auckland.

*pulvinata* (Hook. f. et Wils.) Jaeg. *op. cit.* p. 213.

*Hypnum pulvinatum* H. f. et W. *in op. cit.* p. 155 ; *C. M. Syn. II*, p. 500.

*Isoetecium pulvinatum* H. f. et W. *Fl. of the N. Zeal. II*, p. 105, t. 38 (1855) *et Handb.* p. 465 (1867).

*Ptilocladus pulvinatus* Lindb. *in op. et l. cc.*

*Thamniella* ? *pulvinata* Besch. *in op. et l. cc.*

— *Pac.* : Nov. Seeland. sept.

*ramulosa* (Mitt.) Jaeg. *op. et l. cc.*

*Isoetecium ramulosum* Mitt. *in Handb. of the N. Zeal. Fl.* p. 465 (1867).  
*Porotrichum ramulosum* *ej.* *in Trans. and Proceed. of the roy. Soc. of Victoria*, 1883, p. 84.

*Ptilocladus ramulosus* Lindb. *in op. et l. cc.*

2. *Cortic.* — *Pac.* : Austral. or. (Victoria), Nov. Seeland., Tasman., ins. Chatham.

**Camptheciopsis** Kindb. *Check List. of Eur. and N. Am. Bryin.* p. 74 (sect. **BRACHYTHECI** sensu Kindb.).

**CAMPTOTHECIUM** Bryol. eur. fasc. 46-47 (1853). — 15 spec.

*acuminatum* Kindb. — *Brachythecium*.

*alsioides* Kindb. *in Pittonia*, 1892, p. 243.

2. *Rupic.* — *Am. sept.* : Calif.

*Amesiae* Ren. et Card. *in Bot. Gaz.* 1888, p. 202, t. 20.

2. — *Am. sept.* : Ins. Vancouver, Columb. brit., Washington, Oregon, Calif.

*arenarium* (Lesq.) Jaeg. *Ad. II*, p. 380 (1876-77).

*Hypnum arenarium* Lesq. *in Trans. of the Amer. Phil. Soc. XIII*, p. 13 ; *Lesq. et Jam. Man.* p. 333 (1884).

2. *Arenic.* — *Am. sept.* : Ins. Vancouver, Calif.

*aureolum* Kindb. — *Homalothecium sericeum*.

*aureum* (Lagasca., Garc. et Cl.) *Bryol. eur. fasc. 52-54 (Vol. V), Mon. p. 7, t. 559* (1853) ; *W. P. Sch. Syn. II ed.* p. 636 (1876).

*Brachythecium aureum* de Not. *Epil.* p. 116 (1869).

*Hypnum aurescens* *C. M. Syn. II*, p. 371 (1851).

*H. aureum* Lag., Garc. et Cl. *in Ann. de Cienc. nat. n. 14*, p. 204 (1802) ; *Brid. Bryol. univ. II*, p. 462 (1827).

*Isoetecium aureum* Spr. *M. Pyren. n. 85* (1847) *et in Ann. and Mag. of nat. Hist.* 1849.

2 *et ps. 1.* *Loc. sicc. et ad rup. arid.* — *Eur.* : Galloprov., Etrur., Sardin., Sicil., Pyren., Hispan. *As.* : Palaestin. *Afr.* : Alger. *Am. sept.* : Ins. Vancouver ; Cuba (*fd. Brid.*).

*caucasicum* (Lindb.) *Limpr. Laubm. Deutschl.... III*, p. 58 (1895).

*Hypnum* (*Pleuropus*) *caucasicum* Lindb. *in sched.* ; *Broth. in Act. Soc. sc. fenn.* 1892, n. 12, p. 117.

*Homalothecium caucasicum* Par. *Ind.* p. 567 (1894).

2. *Ad arbor. et rup.* — *As.* : Caucas.

*cladoneuron* Jaeg. — *Brachythecium*.

*corticola* Kindb. — *Homalothecium*.

*dolosum* Ren. et Card. in *Bot. Centralbl.* 1890, n. 51 (*subsp.* *C. ænei*).

Ad rup. umbros. — *Am. sept.* : Ins. Vancouver, Columb. brit., Washington, Idaho, Calif.

*fallax* Philib. — *Homalothecium*.

*fulgenscens* Mitt. in *Bot. Zeit.* 1878, p. 170.

Lignic. — *Am. sept.* : M. Selkirk, Columb. brit.

*Geheebii* Kindb. — *Brachythecium*.

*hamatidens* Kindb. — *C. Nuttallii*.

*lamprochryseum* Kindb. — *Brachythecium*.

*lutescens* (Huds.) *Bryol. eur. fasc.* 52-54 (*Vol.* V), *Mon.* p. 6, t. 558 (1853); *W. P. Sch. op. cit.* p. 635.

*Brachythecium lutescens* de Not. *Epil.* p. 115 (1869).

*Climacium lutescens* Voit, *M. Herbip.* p. 79 (1812).

*Hypnum cassubicum* Scop. *Fl. Carn. II ed.* p. 1327 (1760).

*H. lutescens* Huds. *Fl. angl.* p. 421 (1762); *Hedw. Descript. IV*, p. 40, t. 16 (1794); *Brid. Bryol. univ. II*, p. 466 (1827); *C. M. Syn. II*, p. 370 (1851).

*H. myosuroides* Lag. *Ann. d. Cienc. nat.* n. 24.

*H. myosuroides*  $\beta$  *praelongum* Web. *Spic. Fl. goett.* n. 163  $\delta$  (1778).

*H. sericeum*, surculis longioribus et rectoribus, capsulis incurvis *Dill. Hist. M.* p. 345, t. 42, f. 60 (1741).

*H. sericeum*  $\beta$  *praelongum* Weis *Crypt. Goett.* p. 225 (1770).

*H. sericeum*  $\beta$  *ramosum* Leers *Fl. Herb.* (1775).

*Isothecium lutescens* Spr. *M. Pyren.* n. 88 (1847).

*Leskea longifolia* Ehrh. *Crypt. exs.* n. 281.

*Neckera lutescens* Willd. *Prodr. Fl. berol.* n. 939 (1787).

*N. sericea* var.  $\beta$  *Leyss. Fl. Halens.* n. 1075 (1783).

2 et ps. 1. In petros. sicc. et ad silv. marg. — *Eur.* : Fere tota. *As.* : Boreal., Caucas. occ., Taurus cilic. *Afr.* : Ins. Canar. *Am. sept.* : Alaska, ins. Vancouver, Columb. brit., Washington, Oregon, Idaho, Calif.

« var. *condensatum* Del. *Fl. crypt. Belg.* p. 241 (1883).

— *Eur.* : Belg.

var. *fallax* Breidl. — *Homalothecium*.

« var. *gracile* C. Jens. *Consp. fl. Groenland.* p. 338 (1887).

— *Am. sept.* : Groenland.

« var. *occidentale* Ren. et Card. in *op. et l. cc.*

— *Am. sept.* : Washington, Calif.

« var. *subpinnatum* (Brid.) *Bryol. eur. l. c.*

*Hypnum lutescens* var. *subpinnatum* Brid. *Sp. M. II*, p. 176 (1812).

— *Eur.* : Ital. merid.

*megaptilum* Sull.  *Ic. M. Suppl.* p. 102, t. 78 (1874).

*Hylocomium megaptilum* Kindb. *Eur. and N. Am. Bryin.* p. 48 (1897).

*Hypnum megaptilum* Lesq. et Jam. *Man.* p. 331 (1884).

*Pleurozium megaptilum* Kindb. *Check List of Eur. and N. Am. Bryin.* p. 19 (1894).

2. Terr. in silv. conifer. — *Am. sept.* : Oregon, Idaho, Calif.

*megaptilum* Mac. — *Brachythecium rivulare*.

nevadense (Lesq.) Par. Ind. p. 235 (1893).

Homalothecium nevadense R. C.

Hypnum nevadense Lesq. in Mem. Calif. Acad. I, p. 33; Lesq. et Jam. Man. p. 332 (1884).

H. pseudo-sericeum Mac. Canad. M. p. 135.

2. Rupic. — Am. sept. : Ins. Vancouver, Columb. brit., Nevada, Calif.

var. subulatum R. C. — Homalothecium sericeoides.

nitens (Schreb.) W. P. Sch. Syn. I ed. p. 530 (1880) et II ed. p. 637.

Hypnum nitens Schreb. Spicil. p. 92, n. 1062 (1771); C. M. Syn. II, p. 381 (1851); Bryol. eur. fasc. 57-61 (Vol. VI, Mon. p. 53, t. 622 (1854).

H. palustre erectum trichodes, ramulis crebris, luteo et rufo-virentibus glabris Dill. Cat. Giss. p. 220 (1718) et Hist. M. p. 303, t. 39, f. 37 (1741).

H. trichoides Neck. Del. Gallo-belg. p. 483, n. 35 (1768); Lindb. in Act. Soc. sc. fenn. 1872, p. 274.

H. Stereodon nitens Brid. Bryol. univ. II, p. 560 (1827).

Neckera nitens Mönch Fl. Marp. p. 738 (1794).

2. In pratis palud. et turf. — Eur. : E reg. campestr. usq. in subalp., praecip. zon. sept. : Spitzberg., ins. Ursor. — As. : Jenissei, Taimyr, Ochotsk, Amour. Am. sept. : Groenland, fret. Hudson et Kotzebue, ins. Melville, Anticosti, Terra nov., Ottawa, Ontario, Sackatchewan; Alaska, M. Rupestr., Columb. brit., reg. sept. et centr., Calif.

• var. stricha Kindb. Eur. and N. Am. Bryin. p. 87 (1897).  
Ster. — Eur. : Spitzberg., Norveg.

• var. insignis (Mohl et Schlecht.) Milde in Ltz. Bryol. Notizb. p. 65 (1865).

Hypnum insigne M. et Schl. in Bot. Zeit. 1861.

Ut typ. — Eur. : Silesia.

• var. involuta Limpr. Laubm. Deutschl.... III, p. 60 ( ).

Orthothecium chryseum Helborn in Un. crypt. itin. 1867.

— Eur. : Suecia (Herjedal.).

Nuttallii (Wils.) Bryol. eur. fasc. 52-54 (Vol. VI), Mon. p. 6 (1853).

C. hamatidens Kindb. in Mac. Cat. p. 189 (fid. R. C.).

Homalothecium Nuttalli Jaeg. Ad. II, p. 375 (1876-77).

Hypnum hamatidens Kindb. in Bull. Torr. bot. Club, XVI, p. 97 (1889).

H. Nuttallii Wils. Bryol. brit. pp. 334 et 339 (1855); Sull. Ic. M. p. 211, t. 128 (1864); Lesq. et Jam. Man. p. 332.

H. pinatifidum Mac. Canad. M. n. 280.

Leskea californica Hpe. in Linn. 1860, p. 460.

L. campylocarpa ej. in sched.

2. Cortic. et rupic. — Am. sept. : Ins. Vancouver, Columb. brit., Washington, Oregon, Idaho, Calif.

• var. tenue (Kindb.) R. C. in Rev. bryol. 1893, p. 16.

C. hamatidens  $\beta$  tenue Kindb. in Bull. Torr. bot. Club, XVII, p. 277 (1890).

Ad rup. vertic. — Am. sept. : Columb. brit.

œneum (Mitt.) Jaeg. Ad. II, p. 381 (1876-77).

Hypnum œneum Mitt. in Journ. of the Linn. Soc. 1864, p. 277; Lesq. et Jam. Man. p. 331 (1884).

2. — Am. sept. : Ins. Vancouver, Columb. brit., Washington, Oregon, Idaho, Montana, Sierra Nevada.



*oxycladon* Kindb. — *Brachythecium*.

*Philippi* ej. — *Homalothecium Philippianum*.

*pinnatifidum* (Sull. et Lesq.) Jaeg. op. cit. p. 383.

*Hypnum pinnatifidum* Sull. *Io. M. Suppl.* p. 101, t. 77 (1874); *Lesq. et Jam. Man.* p. 333 (1884).

2. *Rupic.* — *Am. sept.* : Ins. Vancouver, Calif.

*pseudo-lutescens* (Hpe.) Jaeg. op. cit. p. 381.

*Hypnum pseudo-lutescens* Hpe. in *Ann. sc. nat.* 1865, p. 314; *Mitt. M. austr. am.* p. 564 (1869).

*Cortic.* — *Am. merid.* : And. Nov. Granat.

*sericeoides* Kindb. — *Homalothecium*.

*sericeum* ej. — *Homalothecium*.

*spurio-acuminatum* ej. — *Brachythecium*.

*stenocarpum* ej. — *Brachythecium acuminatum subspec. stenocarpum*.

**Campyliadelphus** Lindb. *M. Scand.* p. 32 (sect. *HYPNI*) (1879).

**Campylidium** Kindb. *Eur. and N. Am. Bryin.* p. 119 (sect. *HYPNI sensu Kindb.*) (1897).

**CAMPYLIUM** Sull. *M. and Hepat. of U. St.* p. 677 (1856); *W. P. Sch. I ed.* p. 599 (sect. *HYPNI*); *gen. propr. nonnull.*

*Spec. omn. non indicatae sunt HYPNA ejusd. nom.*

*bergenense* Kindb. — *H. chrysophyllum var. tenellum*.

*Duriei* ej. — *Orthothecium*.

*hygrophilum* ej. — *Amblystegium*.

*Fitz-Geraldii* ej. — *Plagiothecium*.

*protensum* ej. — *H. stellatum var. protensum*.

*striatellum* ej. — *Plagiothecium Mühlenbeckii*.

**CAMPYLOCHÆTIUM** Besch. *Prodr. bryol. mexic.* p. 24 (1871).

*mexicanum* ej. — *Campylopodium*.

**CAMPYLODONTIUM** Schwaegr. *Suppl. III, P. I, I* (1827-28).

*Spec. omn. non indicatae sunt ENTODONTES ejusd. nom.*

*flavescens* Bryol. jav. } — *Fabronia flavescens*.  
*striatum* Dz. et Mk. }

**Campylodryptodon** W. P. Sch. *Syn. I ed.* p. 226 (sect. *RHACOMITRII*) (1860).

**Campylophyllum** W. P. Sch. *Syn. II ed.* p. 721 (subgen. *HYPNI*) (1876).

**CAMPYLOPODIUM** C. M. *Syn. I*, p. 429 (subgen. *AONGSTROEMIAE*) (1849);  
Besch. *Fl. bryol. Nouv. Caled.* p. 189 (1873). — 17 spec.

*Buchanani* (C. M.) *Par. Suppl. Ind.* p. 88 (1900).

*Aongstroemia Buchanani* C. M. in *Hedwig*. 1898, p. 116.

2. *Terr.* — *Pac.* : Nov. Seeland. (Wellington).

*curvisetum* (Hpe.) *Par. Ind.* p. 237 (1894).

*Aongstroemia curviseta* Hpe. *Prodr. Fl. Nov. Granat.* p. 19 (1866)

*Dicranella curviseta* Mitt. *M. austr. am.* p. 38 (1869).

— *Am. merid.* : And. Nov. Granat.

*cygnaeum* (Aongstr.) *Par. op. et l. cc.*

2. — *Afr.* : Ins. S. Helen.

- euphorocladum (*C. M.*) *Besch. Fl. bryol. Nouv. Caled. p.* 189 (1873).  
*Dicranella cygnaea Aongstr. in Oefv. af K. Vet.-Akad. Foerh.* 1878, n. 5, p. 47.  
*Dicranum proscriptum Hsch.?*  
*Aongstroemia euphoroclada C. M. Syn. I, p.* 429 (1849).  
*Campylopus euphorocladus Bryol. jav. I, p.* 79, t. 66 (1855-61).  
*C. nanus Bryol. jav. I, p.* 74, t. 61  
*C. pseudo-nanus Jaeg. Ad. I, p.* 120 } *fid. Fleisch.*  
*Dicranella euphoroclada Jaeg. Ad. II, p.* 637 (1877-78).  
*Dicranum pseudo-nanum C. M. in Bot. Zeit.* 1859, p. 190 (*fid. Fleisch.*).  
*Didymodon grimmoides Duby in Moritzi Verz. d. Zolling. Pfl.* p. 134?  
*D. medius ej. l. c.?*  
*Micro-Campylopus pseudo-nanus C. M. in Hedwig.* 1899, p. 78 (*fid. Fleisch.*).  
 2. *Rupic. et terr. — As. : Java.*  
*var. β squarrosus Besch. — C. Novae Caledoniae.*
- Fendleri (C. M.) Par. Ind. p.* 237 (1894).  
*Aongstroemia Fendleri C. M. in Linn.* 1878-79, p. 470.  
*Dicranella Fendleri Jaeg. Ad. II, p.* 637 (1877-78).  
 2. — *Am. merid. : Venezuela.*
- flexipes (Mitt.) Broth. Natur. Pflanzenfam. p.* 312.  
*Dicranella proscripta Mitt. in Melliss's S. Helena, p.* 357 ( ).  
*Dicranodontium flexipes ej. mss. ; Handb. of the N. Zeal. Fl. II, p.* 413 (1867); *Par. Ind. p.* 338 (1894).  
*Dicranum proscriptum H. f. et W. Fl. of the N. Zeal. II, p.* 67 (1855).  
*Grimmia Buchanani Stirton in Proceed. of the nat. hist. Soc. of Glasgow,* 1876, p. 87.  
 2. — *Pac. : Nov. Seeland. sept.*
- « *var. β* }  
 « *var. γ* } *Handb. l. c. (sub Dicranodontie).*  
*Ubi typ.*
- integrum (C. M.) Par.*  
*Aongstroemia integra C. M. in Bot. Zeit.* 1857, p. 777.  
*Microdus integer Par. Ind. p.* 804 (1895).  
 2. — *Pac. : Ins. Fidji.*
- itatiaiense Broth. in Ule Bryoth. brasil. n.* 102.  
*Aongstroemia itatiaiensis C. M. in Bull. hb. Boissier,* 1898, p. 43 *et in Ule Coll. M. brasil. n.* 1908.  
*Terr. — Am. merid. : Brasil. subtropic. or.*
- khazianum (Griff.) Par. op. et l. cc.*  
*Campylopus tenuis Wils. in sched. et in Kew Journ. bot. IX, p.* 296 (1857).  
*Dicranella khaziana Jaeg. op. et l. cc.*  
*Dicranum khasianum Griff. Not. p.* 418 *et Ic. pl. asiat. II, p.* 94, f. 2 (1849).  
*Leptotrichum khazianum Mitt. M. Ind. or. p.* 8 (1859).  
 2? — *As. : Khasia, Sikkim, Ceylan.*
- micro-Campylopus (C. M.) Par. Suppl. Ind. p.* 88 (1900).  
*Aongstroemia micro-Campylopus C. M. in Fl.* 1886, p. 417.  
 — *Pac. : Ins. Hawai.*
- mexicanum Besch. in litt.*  
*Campylochaetium mexicanum ej. Prodr. bryol. mexic. p.* 24 (1871).  
*Dicranella mexicana Jaeg. op. et l. cc.*  
 — *Am. sept. : Mexic.*

*Novae Caledoniae* (C. M.) *Par. Ind.* p. 237 (1894).

*Aongstroemia Novae Caledoniae* C. M. in *Hedwig.* 1898, p. 115.

*C. euphorocladum* var. *squarrosus* *Besch. Fl. bryol. Nouv. Caled.* p. 189 (1873).

2. *Ad terr. argill.* — *Pac.* : Nov. Caledon.

*phascoides* (C. M.) *Par. Ind.* p. 237 (1894).

*Aongstroemia phascoides* C. M. in *Bot. Zeit.* 1853, p. 39; *Mitt. M. Ind. or.* p. 8 (1859).

*Dicranella phascoides* *Jaeg. Ad. II*, p. 637 (1877-78).

2. — *As.* : M. Neilgherr.

*Pilopogon* (C. M.) *Par. op. et l. cc.*

*Aongstroemia Pilopogon* C. M. in *Linn.* 1874, p. 630.

*Dicranella Pilopogon* *Jaeg. op. et l. cc.*

2. — *Am. sept.* : Mexic.

*Sartorii* (C. M.) *Par. op. et l. cc.*

*Aongstroemia Sartorii* C. M. in *op. cit.* p. 629.

*Dicranella Sartorii* *Jaeg. op. et l. cc.*

2. — *Am. sept.* : Mexic.

*taïtense* *Besch. in Ann. sc. nat.* 1894, p.

1. *In argill.* — *Pac.* : Taïti.

*trichophyllum* (Mitt.) *Par. op. et l. cc.*

*Dicranella trichophylla* Mitt. in *Fl. Vit.* p. 386 (1871).

*Leptotrichum trichophyllum* Mitt. in *Seem. Bonpl.* 1861, p. 336.

2. — *Pac.* : Ins. Fidji, Ovalau.

**CAMPYLOPUS** Brid. *Mant. M.* p. 71 (1819). — 437 spec. (1).

Spec. omn. non indicatae sunt *PILOPOGONES* ejusd. nom.

*abyssinicus* de Not. in *Nuov. Giorn. bot. ital.* 1872, p. 21.

— *Afr.* : Abyssin.

*abyssinicus* W. P. Sch. — *C. subflexuosus*.

*acervatus* Mitt. *M. austr. am.* p. 87 (1869).

— *Am. merid.* : Absque loco.

*acicularis* Wahlenb. — *Rhacomitrium*.

*acrocaulon* (C. M.) *Par. Ind.* p. 228 (1894).

*Dicranum acrocaulon* C. M. in *Fl.* 1888, p. 110.

Ster. — *Afr.* : Kilima N'Djaro.

*acuminatus* Mitt. *M. austr. am.* p. 90 (1869).

*Dicranum acuminatum* C. M. in *Fl.* 1885, p. 409.

Ster. — *Am. merid.* : Fuegia, ins. Eremitae.

*adustus* de Not. *Epil.* p. 649 (1869); W. P. Sch. *Syn. II ed.* p. 107 (1876).

*C. atro-virens* var. *adustus* Husn. *Muscol. gall.* p. 46 (1881).

Ster. — *Eur.* : Ital. super. (ad Verbanum), Pyr. or.

*aethiops* Welw. et Duby in *Mém.... Genève*, 1870, p. 7, t. 3, f. 2.

Ster. — *Afr.* : Angola.

---

(1) De speciebus Novo Seelandicis Brownianis (16) cf. p. 34 in adn.

*afro-concolor* C. M. in *Dus. M. Camer. n.* 653; *Broth. in Engler's Bot. Jahrb.* 1897, p. 235.

Ster. Ad arbor. junior. — *Afr.* : Cameroon.

*albescens* (C. M.) *Par. Ind.* p. 238 (1894).

*Dicranum albescens* C. M. in *Bot. Zeit.* 1853, p. 36; *Mitt. M. Ind. or.* p. 16 (1859).

— *As.* : M. Neilgherr.

*albescens* Kindb. — *Dicranum albicans.*

*alopecurus* (C. M.) *Par. op. et l. cc.*

*Dicranum alopecurum* C. M. in *Linn.* 1880-82, p. 401.

Ad arbor. — *Am. merid.* : Argentin. subtropic.

*alpicola* Jaeg. — *C. torfaceus* var. *alpicola.*

*alpinus* { W. P. Sch. } — *Dicranodontium longirostre* var. *alpina.*  
           { var. *pachyneuron* Mol. }

*altissimus* (C. M.) *Jaeg. Ad. II,* p. 645 (1877-78).

*Dicranum altissimum* C. M. in *Fl.* 1875, n. 34-35.

— *Am. merid.* : And. Nov. Granat.

*alto-filifolius* C. M. in *Ule Bryoth. brasil. n.* 113 et in *Hedwig.* 1900, p. 253.

Terr. — *Am. merid.* : Brasil. austr. or.

*alto-virescens* (C. M.) *Par. Suppl. Ind.* p. 88 (1900).

*Dicranum alto-virescens* C. M. in *Hedwig.* 1899, p. 152.

— *Afr.* : Natal

*amabilis* (C. M.) *Par. Ind.* p. 239 (1894).

*Dicranum amabile* C. M. in *Linn.* 1830-82, p. 399.

Ad arbor. — *Am. merid.* : Argentin. subtropic.

*ampliretis* Rehm. *M. Afr. austr. n.* 60.

*Dicranum amplirete* C. M. in *Hedw.* 1899, p. 81.

Ster. — *Afr.* : Pr. B. Sp. (Montagu.).

*Anderssonii* (C. M.) *Jaeg. Ad. I,* p. 140 (1869-70).

*Dicranum Anderssonii* C. M. in *Bot. Zeit.* 1856, p. 159.

— *Am. merid.* : Ins. Gallapagos.

*Aongstroemii* C. M. mss.; *Aongstr. in Oef. af K. Vet.-Akad. Foerh.* 1876, n. 4, p. 54.

*C. aureo-nitens* *Aongstr. in op. cit.* 1873, p. 43.

— *Afr.* : Ins. Franciaie

*angustiretis* (Aust.) *Lesq. et Jam. Man.* p. 80 (1884).

*Dicranum angustirete* Aust. in *Coult. Bot. Gaz. IV,* p. 150.

— *Am. sept.* : Florid. merid.

*annotinus* *Mitt. M. austr. am.* p. 80 (1869).

— *Am. merid.* : And. Nov. Granat.

*appressifolius* *Mitt. in Handb. of the N. Zeal. Fl.* p. 414 (1867).

*C. clavatus* H. f. et W. *Fl. of the N. Zeal. II,* p. 69 p.p. (1855).

— *Afr.* : Ins. Kerguelen. *Pac.* : Austral. or. (Nov. Vales.), Nov. Seeland.

- Araucarieti* C. M. in *Ule Bryoth. brasil.* n. 51 et in *Hedwig.* 1900, p. 254.  
Terr. — *Am. merid.* : Brasil. or.
- Arbogasti* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1891, II, p. 183;  
*Ren. M. Masc. Mad.* n. 57.  
Ster. Arenic. — *Afr.* : Madagasc., ins. S. Maria.
- arctocarpus* (Hsch.) Mitt. *M. austr. am.* p. 88 (1869).  
Dicranum *arctocarpum* Hsch. *Fl. brasil.* I, p. 12 (1840); *C. M. Syn.* I, p. 398 (1849).  
*D. stenocarpum* C. M. in *op. cit.* p. 404 (*fid. Mitt.*).  
— *Am. merid.* : Venezuela, Brasil. or., Monte Video.
- arctocarpus* Mitt. p.p. — *C. crispicoma.*  
*arcuatus* Mitt. — *C. Echernieri.*  
*Arduennas* Lib. — *C. flexuosus.*
- arenicola* (C. M.) Mitt. *op. cit.* p. 85.  
Dicranum *arenicola* C. M. in *Bot. Zeit.* 1855, p. 762.  
In arena marit. — *Am. merid.* : Brasil. austr. or.
- areodictyon* (C. M.) Mitt. *op. cit.* p. 85.  
Dicranum *areodictyon* C. M. *Syn.* I, p. 394 (1849).  
Lignic. — *Am. sept.* : Guatemala. *Am. merid.* : And. Növ. Granat. et Quitens.
- areodictyon* W. P. Sch. — *C. leucognodes.*  
var. *gracilescens* ej. — *C. leucognodes* var. *gracilescens.*
- asperifolius* Mitt. *op. cit.* p. 79.  
Ster. — *Am. merid.* : And. Quitens. et Chilens.
- asperulus* Kindb. — *Dicranodontium aristatum.*
- atro-luteus* (C. M.) Rehm. *M. Afr. austr.* n. 63.  
*C. brevis* Rehm. in *Coll. cit.* n. 44; *Par. Ind.* p. 241.  
Dicranum *atro-luteum* C. M. in *Hedwig.* 1899, p. 80.  
Ster. — *Afr.* : Pr. B. Sp.
- atro-sordidus* C. M. in *Dusen M. Cam.* n. 347; *Broth. in Engler's Bot. Jahrb.* 1897, p. 233.  
Rupic. — *Afr.* : Cameroon.
- atro-virens* de Not. — *C. longipilus.*  
var. *adustus* Husn. — *C. longipilus.*  
var. *falcata* Braithw. } — id. varr. ejusd. nom.  
var. *mutica* Milde }
- aureo-nitens* (C. M.) Jaeg. *Ad. II,* p. 435 (1876-77).  
*C. introflexus* Brid. *Bryol. univ.* I, p. 472 p.p. (1827).  
Dicranum *aureo-nitens* C. M. *Syn.* I, p. 406 (1849).  
— *Afr.* : Ins. Borboniae.
- aureo-nitens* Aongstr. — *C. Aongstroemii.*
- aureo-viridis* W. P. Sch. in *Breutel M. capens*; *Jaeg. Ad. I,* p. 139 (1870-71).  
Dicranum *aureo-viride* C. M. in *Hedwig.* 1899, p. 85.  
— *Afr.* : Pr. B. Sp.
- aureus* *Bryol. jav.* I, p. 80, t. 67 (1855-61).  
Ad arbor. — *As.* : Celebes, Java, ins. Nicobar, Ceylan.

*auribrunneus* (C. M.) Par.

*Dicranum auribrunneum* C. M. in *Hedwig*. 1900, p. 260.

Ster. Rupic. — *Am. merid.* : Brasil. or. (Goyaz).

*auriculatus* Wils. — C. Schwarzii.

*aurifciscus* (C. M.) Par. *Ind.* p. 240 (1894).

*Dicranum aurifciscum* C. M. in *Linn.* 1880-82, p. 399.

Ster. — *Am. merid.* : Uruguay.

♣ *var. flavescens* (C. M.) Par. *op. et l. cc.*

*Dicranum aurifciscum var. flavescens* C. M. in *op. et l. cc.*

Ubi typ.

*australiensis* Duby in *Mém.... Genève*, 1869, p. 9, t. 4, f. I.

— *Pac.* : Austral.

*austro-alpinus* (C. M.) Par. *op. et l. cc.*

*Dicranum austro-alpinum* C. M. in *Linn.* 1880-82, p. 398.

Ster. — *Am. merid.* : Alp. Argentin. subtropic. Tucuman.

*austro-subulatus* Broth. et Geh. in *Oefv. af Finska Vet.-Akad. Foerh.* 1895, p. 52.

Ster. — *Pac.* : Nov. Guinea.

*azoricus* Mitt. in *Godman's Nat. hist. of I. Azor.* p. 292 (1870).

Ster. — *Afr.* : Ins. Azor.

*Balansacanus* Besch. *Fl. bryol. Nouv. Caled.* p. 199 (1873).

C. aureus (fid. *Fleisch.*).

Ad arbor. in silv. — *Pac.* : Nov. Caledon.

*bartramiaeus* C. M. — C. leucobasis *var. bartramiaeus*.

*basalticolus* (C. M.) Par. *Suppl. Ind.* p. 89 (1900).

*Dicranum basalticolum* C. M. in *Hedwig*. 1899, p. 82.

Ster. In fiss. rup. basalt. — *Afr.* : Pr. B. Sp.

*batjunensis* Broth. — *Dicranodontium*.

*Berteroanus* Duby in *Mém.... Genève*, 1875, p. 12, t. I, f. 4.

C. introflexus Mont. in *Ann. sc. nat.* 1835, p.

Silvic. — *Am. merid.* : Ins. Juan Fernandez.

*Bessoni* R. C. — C. polytrichoides *var. Bessoni*.

*Beyrichianus* Duby, *Choix de Crypt. ex.* 1869, p. 9, t. I, f. 3.

Locis irrig. — *Am. merid.* : Brasil. austr. or.

♣ *var. elatius* (Hpe.) Jaeg. *Ad. II*, p. 760 (1877-78).

*Dicranum Beyrichianum var. elatius* Hpe. *Symb.* 1877, p. 719.

Ubi typ.

*bicolor* (Hsch.) *Handb. of the N. Zeal. Fl.* p. 115 (1867).

*Dicranum bicolor* Hsch. in *Sieber M. exs. n. 9*; *C. M. Syn. I*, p. 392 (1819); *Fl. of the N. Zeal. II*, p. 69 (1855).

— *Pac.* : Austral., fret. King's Georges, Nov.

Seeland., Tasman.

*bicolor* Besch. — C. pachycomus.

*Boivinianus* Besch. *Fl. bryol. Réunion, etc.* p. 32 (1880-81).  
— *Afr.* : Ins. Franciae.

*Boryanus* Besch. *op. cit.* p. 31.  
— *Afr.* : Usambara, ins. Borbon.

*Boswellii* Hpe. *hb.*

Dicranum *Boswellii* C. M. in *Hedwig*. 1897, p. 442.

Ster. — *Pac.* : Ins. Hawaï.

*brachymastix* C. M. in Besch. *Fl. bryol. Réunion, etc.* p. 36.  
Fr. ign. — *Afr.* : Ins. Franciae.

*brachymitrius* (Geh. et Hpe.) Par. *Ind.* p. 241 (1894).

Dicranum *brachymitrium* G. et H. in *Fl.* 1881, p. 341.

— *Am. merid.* : Brasil. austr. or.

*brachyphyllulus* (C. M.) Par.

Dicranum *brachyphyllulum* C. M. in *Hedwig*. 1900, p. 260.

Ster. Rupic. — *Am. merid.* : Brasil. or. (Minas Geraës).

*brachyphyllus* Wils. in sched.

Dicranum *brachyphyllum* Mitt. in *Lond. Journ. of bot.* 1851, p. 52 et *M. austr. am.* p. 81 (1869) ; *C. Syn. II*, p. 597 (1851).

Lignic. et terr. — *Am. merid.* : And. Quitens.

*brachyphyllus* C. M. (*Afr.*). — *C. subbrachyphyllus*.

*brachythysanos* (C. M.) Par.

Dicranum *brachythysanos* C. M. in *Hedwig*. 1897, p. 97.

— *Am. merid.* : Monte-Video.

*brevifolius* W. P. Sch. *Bryol. eur. Suppl. fasc.* 1-2, t. 2 (1864) et *Syn. II ed.* p. 106 (1876).

*C. subulatus* ej. in *Rab. Bryoth. eur.* n. 451 et *Un. crypt. itin.* 1863, n. 9.  
*Orthopus subulatus* Wulfsb. in *Christian. Vid.-Selsk. Foerh.* 1875, p. 351.

Fl. ! et fr. ? ign. — *Eur.* : Ardenn., Belgia, Carinth., Bavar., Tirol.  
merid., Helvet., Simplon, Gall. centr. et occid.

« *var. elongatus* Boswell in the *Naturalist*, 1883, p. 28.

In uliginosis. — *Eur.* : Anglia (Vales. c. fr. !), Stir.

*brevifolius* de Not. — *C. Mildei*.

*var. Schimperii* Husn. — *C. Schimperii*.

*brevipilus* *Bryol. eur. fasc.* 41 (*Vol. I*), *Mon.* p. 7, t. 92 (1847) ; *W. P. Sch. Suppl. fasc.* 1-2, t. 2, f. 1 et 2 (1844) et *Syn. II ed.* p. 106 (1876).

Dicranum *brevipilum* C. M. *Syn. I*, p. 412 (1849).

*C. decipiens* Van der Sande Lac. mss.

*C. Molkenboëri* Van der Sande Lac. mss. ; *Milde Bryol. siles.* p. 79 (1869)  
ut *var.* ; *Jaeg. Ad. I*, p. 142 (*id.*).

♂ et fr. ign. In ericet. et subturfac. — *Eur.* : Dania, Westphal.,  
Neerland., Britann., Gall. centr. et occ. *Afr.* : Alger.

« *var. auriculatus* Ferg. mss. ; *Dix. et Jam. Stud. Handb.* p. 98 (1896).

— *Eur.* : Anglia.

« *var. compactus* Card. in *Bull. Soc. roy. bot. Belg. XXIV, II*, p. 86.

— *Eur.* : Belgia, Gallia.

« *var. elatus* Card. in *Del. Fl. crypt. Belg. M.* p. 281 (1883).

— *Eur.* : Belgia.

- brevis* Rehm. — *C. atro-luteus*.  
*Brotherianus* Par. *Suppl. Ind.* p. 89 (1900).  
*C. procerus* Broth. in *Ule Bryoth. brasil.* n. 110.  
 Rupic. — *Am. merid.* : Brasil. subtropic. or.  
*brunneus* (*C. M.*) Par. *op. et l. cc.*  
*Dicranum brunneum* *C. M.* in *Hedwig.* 1897, p. 352.  
 Ster. — *Pac.* : Tasman.  
*Cacti* (*C. M.*) Par. *Ind.* p. 241 (1894).  
*Dicranum Cacti* *C. M.* in *Linn.* 1880-82, p. 403.  
 — *Am. merid.* : Uruguay.  
*caecuminis* (*C. M.*) Par. *Suppl. Ind.* p. 90 (1900).  
*Dicranum caecuminis* *C. M.* in *Hedwig.* 1898, p. 227.  
 — *Am. sept.* : M. Jamaic.  
*caespiticius* Brid. — *Grimmia sulcata*.  
*caespitosus* Jaeg. — *Dicranodontium*.  
*Cailleae* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* I (1893), p. 114 ;  
*Ren. M. Masc. Mad.* n. 205.  
 Ster. Ad terr. argill. — *Afr.* : Nossi Comba, Madagasc.  
*caldensis* Aongstr. in *Oefv. af K. Vet.-Akad. Foerh.* 1870, n. 4, p. 5.  
 Ster. — *Am. merid.* : Brasil. austr. or.  
*calvus* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1894, II, p. 113 ; *Ren. M. Masc. Mad.* n. 160.  
 Ster. Ad terr. arenac. subterfos. — *Afr.* : Madagascar.  
*calymperidictyon* (*Geh. et Hpe*) Par. *Ind.* p. 242 (1894).  
*Dicranum calymperidictyon* G. et H. in *Fl.* 1881, p. 242.  
 — *Am. merid.* : Brasil. austr. or.  
*Cambouei* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1891, II, p. 182.  
 Ster. Ad terr. argill. — *Afr.* : Madagascar.  
 « *var. amblyacron* Ren. *Prodr. fl. bryol. Madag. etc.* p. 292 (1897).  
 Ubi typ.  
*campiadelphus* (*C. M.*) Par.  
*Dicranum campiadelphus* *C. M.* in *Hedwig.* 1900, p. 252.  
 Terr. — *Am. merid.* : Brasil. or. (Minas Geraës).  
*canaliculatus* (*Geh. et Hpe.*) Par. *Ind.* p. 242 (1894).  
*Dicranum canaliculatum* G. et H. *Enum. M. Brasil.* p. 20 (1879).  
 — *Am. merid.* : Brasil. austr. or.  
*canescens* W. P. Sch. in *Leckler Pl. ins. Maclov.* n. 90.  
*Dicranum canescens* *C. M.* in *Bot. Zeit.* 1859, p. 220.  
 — *Am. merid.* : Ins. Maclov.  
*capensis* W. P. Sch. — *C. lepidophyllus*.  
*capillaceus* (*Brid.*) Jaeg. *Ad. II*, p. 429 (1876-77).  
*Dicranum capillaceum* *Brid. Bryol. univ.* I, p. 160 (1826) ; *C. M. Syn. I*,  
 p. 396 (1849).  
 — *Afr.* : Madagasc.



- capillaceus* H. f. et W. — C. Holomitrium.  
*capillatus* Hook. f. et Wils. *Fl. Tasman.* II, p. 172, t. 172, f. 4 (1860).  
 — Pac. : Austral., Tasman.  
*capitiflorus* Mont. *Syll.* p. 42 (1856).  
 Dicranum capitiflorum C. M. *Syn.* II, p. 598 (1851).  
 — Afr. : Ins. Borbon.  
 « *var. pachycomus* Besch. *Fl. bryol. Réunion, etc.* p. 34 (1880-81).  
 Ad arbor. — Afr. : Ins. Franciae.  
*carassensis* (Broth.) Par. *Ind.* p. 242 (1894).  
 Thysanomitrium carassense Broth. in *Act. Soc. sc. fenn.* 1891, n. 5.  
 Ster. Terr. — Am. merid. : Brasil. or. (Minas Geraës).  
*cataractarum* Fleisch. mss.  
 — As. : Java.  
 « *var. rufidulum* ej. in *M. Archip. ind.* n. 111 (1900).  
 Ad cataract. inter *Sphagna*. — As. : Java.  
*catarratilis* (C. M.) Par. *Suppl. Ind.* p. 90 (1900).  
 Dicranum catarratile C. M. in *Rehm. M. Afr. austr.* n. 64 et in *Hedwig.* 1899, p. 79.  
 Ster. — Afr. : Prom. R. Sp.  
*catharinensis* Par.  
 Dicranum perfalcatum C. M. in *Hedwig.* 1900, p. 250.  
 Ster. Terr. ? — Am. merid. : Brasil. austr. or. (ins. S. Catharina).  
*catumbemis* Broth. in *Bihang till. K. Sw. Vet.-Akad. Foerh.* 1895, n. 3.  
 In arenos. sicc. — Am. merid. : Brasil. austr. or.  
*caudatus* (C. M.) *Bryol. jav.* I, p. 78, t. 65 (1855-61).  
 C. flexuosus Mont. in *Ann. so. nat.* 1842, *Crypt. Neilgh.* n. 52.  
 Dicranum caudatum C. M. *Syn.* I, p. 401 (1849) ; *Mitt. M. Ind. or.* p. 18 (1859).  
 Cortic., lignic., rupic. et terr. — As. : M. Neilgherr., Java, Horneo, Molucc., Halmaheira.  
*cavifolius* Mitt. M. *austr. am.* p. 87 (1869).  
 — Afr. : Ins. Kerguelen, Marion. Am. merid. : And. Quitens.  
*chalarobasis* C. M. in *Dusen M. Camer.* n. 262.  
 — Afr. : Cameroon.  
*chilensis* de Not. in *Mitt. M. austr. am.* p. 85 (1869).  
 — Am. merid. : Chile.  
*chionophilus* (C. M.) *Mitt. op. cit.* p. 81.  
 Dicranum chionophilum C. M. *Syn.* I, p. 398 (1849).  
 — Am. merid. : And. Nov. Granat.  
*chlorophyllosus* (C. M.) *Jaeg. Ad.* I, p. 132 (1870-71).  
 Dicranum chlorophyllosum C. M. *op. cit.* p. 395.  
 Ster. — Afr. : Transvaal, pr. B. Sp.  
 « *var. compactus* } C. M. in *Rehm.* { n. 49 } — Afr. : Pr. B. Sp.  
 « *var. rivularis* } M. Afr. austr. { n. 48 }  
 « *var. tristis* } { n. 47. — Afr. : Pr. B. Sp., Natal,

- chlorotrichus *C. M.* in *Coll. cit. n.* 53 et 449.  
— *Afr.* : Pr. B. Sp., Natal.
- Chrismari (*C. M.*) *Mitt. M. austr. am.* p. 88 (1869).  
Dicranum Chrismari *C. M.* in *Bot. Zeit.* 1855, p. 761.  
— *Am. sept.* : Mexic.
- chryseolus *C. M.* in *Besch. Fl. bryol. Réunion, etc.* p. 37 (1880-81).  
An Pilopogon ? (*Broth.*)  
— *Afr.* : Comor. magn., Madagasc.
- chrysodictyon (*Hpe.*) *Mitt. M. austr. am.* p. 78 (1869).  
Dicranum chrysodictyon *Hpe.* in *Ann. sc. nat.* 1866, p. 338.  
— *Am. merid.* : And. Nov. Granat.
- Cinchonae *Par. Suppl. Ind.* p. 90 (1900).  
Thysanomitrium jamaïcense *C. M.* in *Bull. hb. Boissier*, 1897, p. 552.  
Ster. Ad Cinchonas. — *Am. sept.* : Jamaïc.
- cirratus *Hsch.* — *Cynodontium gracilescens var. inflexa.*
- Civa *Ltz. Moosst.* p. 150 (1864) (1).  
Ster. — *As.* : India (absque loc. spec.)
- clavatus (*R. Br.*) *Hook. f. et Wils. Fl. of the N. Zeal. II*, p. 69 p.p. (1855).  
Dicranum clavatum *R. Br.* in *Schwaegr. Suppl. III, P. II, I, t.* 255 (1829-30); *C. M. Syn. I*, p. 412 (1849).  
*D. pudicum Hsch.* in *Sieber M. Nov. Holl. n.* 24 p.p. (*fid. Mitt.*).  
— *Afr.* : Ins. S. Paul. Pac. : Austral., Nov. Seeland., Tasman.
- clavatus { *H. f. et W.* p.p. } — *C. appressifolius.*  
                  { *Rehm.* — *C. pseudo-clavatus.*  
                      — *C. insititius.*
- collinus (*C. M.*) *Par. Suppl. Ind.* p. 90 (1900).  
Dicranum collinum *C. M.* in *Bull. hb. Boissier*, 1898, p. 38 et in *Ule Coll. M. Brasil.* n. 948.  
— *Am. merid.* : Brasil. austr. or.
- comatulus *Besch. Not. M. des I. St-Paul et Amsterdam*, p. 4 (1875).  
Ster. — *Afr.* : Ins. Amsterdam.
- comatus *Ren. et Card. in Bull. Soc. roy. bot. Belg.* 1893, II, p. 183;  
*Ren. M. Masc. Mad. n.* 55.  
Ster. Silvic. — *Afr.* : Madagasc., ins. S. Maria.
- Commersoni *Besch. mss.*; *Ren. Prodr. fl. bryol. Madag. etc.* p. 88 (1897).  
Ster. — *Afr.* : Ins. Borbon.
- comosus (*Hsch. et Reinw.*). — *Sphaerothecium.*
- compactus { *W. P. Sch.* — *C. Schimper.*  
                  { *Wils.* p.p. — *C. Goughii.*
- concolor (*Hook.*) *Brid. Bryol. univ. I*, p. 476 (1826); *Mitt. M. austr. am.* p. 82 (1869).  
*C. Jamesoni Jaeg. Ad. II*, p. 442 (1876-77).  
Dicranum concolor *Hook. M. ex.* p. 10, t. 138 (1818-20); *C. M. Syn. I*, p. 391 (1849).  
*D. Jamesoni Hook. Ic. pl. rar.* t. 179 (1841); *C. M. op. et l. cc.*  
Lignic. — *Am. merid.* : And. Nov. Granat.

---

(1) « Forsan *C. nodiflori* forma sterilis » *Ltz.* l. c.

- contortus* Brid. — *Grimmia*.
- controversus* (Hpe) Jaeg. *Ad. II*, p. 649 (1877-78).  
*Dicranum controversum* Hpe. *Symb.* 1872, p. 42.  
 Silvic. — *Am. merid.* : Brasil. austr. or.
- corralensis* Broth. in *Dusen M. Am. austr. n.* 87.  
 Ad rupes littor. — *Am. merid.* : Chile austr.
- crassinervis* Wils. — *Dicranum albicans*.
- crassissimus* Besch. in *Bull. Soc. bot. Fr.* 1885, p. LVIII et in *Miss. sc. Cap Horn*, V, *Bot. p.* 265, t. 2, f. 7.  
 Ster. — *Am. merid.* : Patagon. occid.
- crateris* Besch. *Fl. bryol. Réunion, etc. p.* 39 (1880-81).  
 Ster. Ad rup. vulcan. — *Afr.* : Ins. Borbon (1).
- Crishna Ltz. Moost. p.* 159 (1864) (2).  
 Ster. — *As.* : India (absq. loc. spec.).
- crispatus* Broth. in *sched.*  
*Dicranum crispatum* C. M. in *Bull. hb. Boissier*, 1898, p. 36 et in *Ulc Coll. M. brasil.* n. 1346, 1799, 1800.  
 Cortic. — *Am. merid.* : Brasil. or.
- crispicoma* (C. M.) Jaeg. *Ad. II*, p. 647 (1877-78).  
*C. arctocarpus* Mitt. *M. austr. am. p.* 87 p.p. (1869).  
*Dicranum crispicoma* C. M. in *Fl.* 1875, n. 34-35.  
 Ster. — *Am. merid.* : Venezuela.
- Crügeri* (C. M.) Par. *Suppl. Ind. p.* 91 (1900).  
*Dicranum Crügeri* C. M. in *Hedwig*. 1898, p. 226.  
 Ster. — *Am. merid.* : Ins. Trinitat.
- cryptopodioides* Broth. in *Bihang till K. Sv. Vet.-Akad. Foerh.* 1900, p. 2. fr. ign. — *Am. merid.* : Brasil. centr. (Matto Grosso).
- cubensis* Sull. — *C. filifolius*.
- Cummingii* Duby in *Mém.... Genève*, 1869, p. II, t. II, f. 3.  
 — *Am. merid.* : Chile.
- curvifolius* Brid. — *Grimmia commutata*.
- cuspidatus* (Hsch.) Mitt. *M. austr. am. p.* 90 (1869).  
*Dicranum cuspidatum* Hsch. *Fl. Brasil. I*, p. 13 (1840); *C. M. Syn. I*, p. 414 (1849).  
 Ster. — *Am. merid.* : Brasil. austr. or.
- cygnaeus* (Hedw.) Brid. *Bryol. univ. I*, p. 475 (1826); *Mitt. op. cit. p.* 76.  
*Dicranum cygnaeum* Hedw. *Sp. M. p.* 148, t. 37 (1801); *C. M. op. cit. p.* 388.  
 — *Am. sept.* : Haïti.

---

(1) *cribrosus* Brid. *Mant. M. p.* 76. { — *Grimmia leucophaea* } Deleatur.  
 Cf. C. M. *Syn. I. p.* 810 et *Dryptodon cribrosus*.  
 (2) « Forsan *C. erythroglyphato* nimis aff. » Ltz. l.c.

*deciduus* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1894, II, p. 11;  
*Ren. M. Masc. Mad.* n. 161.

— *Afr.* : Madagasc.

*decipiens* Lac. — *C. brevipilus*.

*declivium* Par.

*Dicranum nano-filifolium* C. M. in *Hedwig.* 1900, p. 254.

*Terr.* — *Am. merid.* : Brasil, austr. or.

*Delagoae* (C. M.) Par. *Suppl. Ind.* p. 91 (1900).

*Dicranum Delagoae* C. M. in *Hedwig.* 1899, p. 86.

*Ster.* — *Afr.* : Prom. Sin. Delagoae.

*densicoma* (C. M.) Par. *Suppl. Ind.* p. 91 (1900).

*Dicranum densicoma* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 33.

— *Am. merid.* : And. Boliv.

*densifolius* Aongstr. in *Oefv. af K. Sv. Vet. Akad. Foerh.* 1872, n. 4,  
 p. 18.

*C. plumosus* Sull. in *Expl. Exped. Wilkes*, p. 4? (C. M.).

— *Pac.* : Ins. Hawai.

*densus* Bryol. eur. }  
 var. *fragilis* Wils. } — *C. fragilis*.

*denudatus* Kindb. — *Dicranodontium longirostre*.

*destructilis* (C. M.) Jaeg. *Ad. I*, p. 131 (1870-71).

*Dicranum destructile* C. M. in *Bot. Zeit.* 1859, p. 220.

*Ster.* — *Am. sept.* : Mexic.

*detonsus* (Hpe.) Par. *Ind.* p. 245 (1894).

*Dicranum detonsum* Hpe. in *Fl.* 1881, p. 341.

— *Am. merid.* : Brasil, austr. or.

*dichotomus* Brid. *Mant. M.* p. 77. — Cf. *Dryptodon dichotomus*.

*dichrotis* (C. M.) Par.

*Dicranum dichrotis* C. M. in *Hedwig.* 1900, p. 255.

— *Am. merid.* : Brasil, or. (Minas Geraës.).

*dicnemoides* (C. M.) Par. *Suppl. Ind.* p. 91 (1900).

*Dicranum dicnemoides* C. M. in *Bull. hb. Boissier*, 1898, p. 37 et in *Ule Coll. M. brasil.* n. 1788.

In *decliv. rupestr.* — *Am. merid.* : Brasil, or. (S. do Itatiaia).

*dictycion* (false : *dictyon*) Jaeg. — *Dicranodontium*.

*Didrichseni* (C. M.) Jaeg. *Ad. I*, p. 128 (1870-71).

*Dicranum Didrichseni* C. M. in *Bot. Zeit.* 1862, p. 329.

*Ster.* — *Pac.* : Ins. Hawai.

*Didymodon* Jaeg. — *Dicranodontium*.

*Dietrichiae* (C. M.) Jaeg. *Ad. II*, p. 647 (1877-78).

*Dicranum Dietrichiae* C. M. *M. Polyn.* p. 62 (1874).

*Ster.* — *Pac.* : Ins. Tonga.

- discriminatus* (Hpe.) Jaeg. op. cit. p. 760.  
*Dicranum discriminatum* Hpe. *Symb.* 1877, p. 718.  
 Cortic. et terr. — *Am. merid.* : Brasil. austr. or.
- dissitus* C. M. in *Dusen M. Camer. n.* 264; *Broth. in Engler's Bot. Jarhb.* 1897, p. 234.  
 Ster. Cortic. — *Afr.* : Cameroon.
- distractus* (C. M.). *Par. Suppl. Ind. p.* 91 (1900).  
*Dicranum distractum* C. M. in *Hedwig.* 1897, p. 350.  
 Ster. — *Pac.* : Nov. Seeland.
- gütrichoides* *Broth. in Act. Soc. sc. fenn.* 1891, n. 5.  
 Ster. Terr. — *Am. merid.* : Brasil. centr. (Minas Geraës).  
 « *var. robustior* ej. in *Bihang till K. Sv. Vet.-Akad. Foerh.* 1895, n. 3.  
 — *Am. merid.* : Brasil. austr. or.
- divaricatus* (Mitt.) Jaeg. *Ad. II, p.* 645 (1877-78).  
*Dicranum divaricatum* Mitt. in *Journ. of the Linn. Soc.* 1863, p. 149.  
 Ster. — *Afr.* : Ins. Princip.
- divisus* (Geh. et Hpe.) *Par. Ind. p.* 245 (1894).  
*Dicranum divisum* Geh. et Hpe. in *Fl.* 1881, p. 315.  
 — *Am. merid.* : Brasil. austr. or.
- dolosus* Besch. *Fl. bryol. Réunion, etc. p.* 32 (1880-81).  
 Terr. — *Afr.* : Ins. Borbon.
- Donnellii* (Aust.) Lesq. et Jam. *Man. p.* 79 (1884).  
*Dicranum Donnellii* Aust. in *Coult. Bot. Gaz. IV, p.* 150.  
*D. sublencogaster* ej. *M. Appal. exs. Suppl. I, p.* 470.  
 Ster. — *Am. sept.* : Florid. merid.
- Dozyanus* Jaeg. — *Pilopogon Blumii.*
- Duseni* C. M. in *Dusen M. Camer. n.* 304.  
 — *Afr.* : Cameroon.
- Echernieri* Besch. *Fl. bryol. Réunion, etc. p.* 35 (1880-81).  
*C. arcuatus* Mitt. in *Phil. Trans. of the Linn. Soc.* 1879 (extra-vol.) p. 389?  
*Dicranum arcuatum* Brid. *Bryol. univ. I, p.* 463 p.p. (1826); *C. M. Syn. I, p.* 388 p.p. (1819).  
 Ster. Locis cav. humid. et aquat. — *Afr.* : Ins. Tristan da Cunha?, Borbon.
- echinatus* Rehm. *M. Afr. austr. n.* 67.  
 Ster. — *Afr.* : Pr. B. Sp.  
 « *var. leucobasis* ej. in *Coll. cit. n.* 71.  
 — *Afr.* : Pr. B. Sp. (Montagu).  
 « *var. turgescens* ej. in *Coll. cit. n.* 69.  
 — *Afr.* : Pr. B. Sp. (Campo Bay.).  
 « *var. umbrosus* ej. in *Coll. cit. n.* 72.  
 — *Afr.* : Pr. B. Sp. (Knysma).  
 « *var. Vallis Gratiae* ej. in *Coll. cit. n.* 70.  
 — *Afr.* : C. B. Sp. (Montagu).
- ellipticus* Brid. — *Rhacomitrium.*

- elongatus* Brid. *Bryol. univ.* I, p. 815 (1826).  
*Dicranum Grateloupii* C. M. *Syn.* I, p. 387 (1849) et II, p. 596 (1851).  
 ♂ et fr. ign. — *Eur.* : Gall. austr. occident.  
*erectus* (C. M.) Mitt. *M. austr. am.* p. 89 (1869).  
*C. strictus* C. M. in *Linn.* 1847, p. 207.  
*Dicranum erectum* C. M. *Syn.* I, p. 408 (1849).  
*Thysanomitrium inflexum* Hpe. in *Linn.* 1847, p. 74.  
 — *Am. merid.* : Venezuela, Columb.  
*ericeticola* C. M. in *Abhandl. Nat. Ver. Brem.* 1900, p. 496.  
 — *Pac.* : Ins. Chatham.  
 « *var. minor* ej. in op. cit p. 497.  
 Ubi typ.  
*ericetorum* Jaeg. — *C. polytrichoides*.  
*ericoides* (Griff.) Jaeg. *Ad.* I, p. 128 (1870-71).  
*Dicranum ericoides* Griff. *Not.* p. 420 (1849); Mitt. *M. Ind.* or. p. 18 (1859)  
 Muric. et terr. — *As.* : Khasia, Ceylan, Java.  
*erythrocaulon* Broth. in *Boll. da Soc. Broter.* 1890, p.  
 Ster. — *Afr.* : Ins. Sao Thoma.  
*erythrodontius* (Hpe.) Jaeg. *Ad.* II, p. 760 (1877-78).  
*Dicranum erythrodontium* Hpe. *Symb.* 1877, p. 720.  
 — *Am. merid.* : Brasil. austr. or.  
*erythrognaphalon* (C. M.) Jaeg. *Ad.* I, p. 143 (1870-71).  
*Dicranum erythrognaphalon* C. M. in *Bot. Zeit.* 1853, p. 37; Mitt. *M. Ind.*  
 or. p. 21 (1859).  
 — *As.* : M. Neilgherr.  
*erythropoma* Duby in *Mém.... Genève*, 1869, p. 10, t. 3, f. 2.  
 — *Pac.* : Austral. or. (Victoria).  
*eunanus* (C. M.) Par.  
*Dicranum eunanum* C. M. in *Hedwig.* 1897, p. 348.  
 — *Pac.* : Nov. Caledon.  
*euphoro-cladus* Bryol. jav. — *Campylopodium*.  
*exalaris* (Hpe.) Par. *Ind.* p. 246 (1894).  
*Dicranum exalare* Hpe. *Symb.* 1877, p. 721.  
 — *Am. merid.* : Brasil. austr. or.  
*exaltatus* (C. M.) Par. *op. et l. cc.*  
*Dicranum exaltatum* C. M. in *Linn.* 1878-79, p. 472.  
 Ster. — *Am. merid.* : Venezuela.  
*exfimbriatus* C. M. in *Fl.* 1897, p. 421.  
 — *Am. merid.* : And. Venezuel.  
*eximius* Reichdt. in *Novara Exped.* 1879, p. 167, t. 28.  
*C. introflexus* Mitt. *Fl. Vit.* p. 385 (1871).  
 — *Afr.* : Ins. Marion, St-Paul. *Pac.* : Nov. Caledon.  
*exustus* Mitt. *M. austr. am.* p. 83 (1869).  
 — *Am. merid.* : And. Peruv.  
*falcatus* Ferg. — *C. longipilus*.

*falcifolius* Mitt. in *Journ. of the Linn. Soc.* XIV, p. 78.

— *Afr.* : Ins. S. Paul.

*Fendleri* (C. M.) Par. *Ind.* p. 247 (1894).

*Dicranum Fendleri* C. M. in *Linn.* 1878-79, p. 472.

— *Am. merid.* : And. Venezuel.

*Ferromecocae* Par.

*Dicranum Kunerti* C. M. in *Hedwig.* 1900, p. 255.

Ster. — *Am. merid.* : Brasil. austr. (Rio Grande do Sul.)

*fibrobasis* Dus. M. *Am. austr. n. A.* 18.

Ad rup. littor. — *Am. merid.* : Patagon. occid.

*filescens* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1890, I, p. 169;

*Ren. M. Masc. Mad. n.* 9.

Lignic. — *Afr.* : Madagasc.

*filicaudatus* (C. M.) Par.

*Dicranum filicaudatum* C. M. in *Hedwig.* 1900, p. 260.

Ster. Rupic. — *Am. merid.* : Brasil. or. (Minas Geraës).

*filicaulis* (Hpe.) Mitt. M. *austr. am.* p. 74 (1809).

*Dicranum filicaule* Hpe. in *Ann. sc. nat.* 1866, p. 337.

Silvic. — *Am. merid.* : And. Nov. Granat.

*filifolius* (Hsch.) Mitt. *op. cit.* p. 76.

C. cubensis Sull. in *Proced. of the Amer. Acad. August.* 1861.

*Dicranum Mulleri* Ltz. ap. Geh. et Hpe. in *Fl.* 1881, p. 345.

*Dicranum filifolium* Hschs. *Fl. brasil.* I, p. 12 (1840); C. M. *Syn.* I, p. 403 (1849).

Lignic. et terr. — *Am. sept.* : Cuba, Porto Rico. *Am. merid.* : Brasil. austr. or.

*fimbriatus* Mitt. *op. cit.* p. 88.

Ad arbor. — *Am. merid.* : And. Quitens.

*flaccidus* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1893, II, p. II.

Ster. In sphagnetis. — *Afr.* : Madagasc.

*flagellaceus* (C. M.) Mitt. *op. cit.* p. 77.

*Dicranum flagellaceum* C. M. *Syn.* II, p. 597 (1851).

Ster. — *Am. sept.* : Mexic.

*Flageyi* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1893, II, p. 12; Ben.

*M. Masc. Mad. n.* 204.

Silvic. — *Afr.* : Madagascar.

*flavescens* Rehm. M. *Afr. austr. n.* 445.

— *Afr.* : Transvaal.

*flavicomma* C. M. in *Dusen M. Camer. n.* 299; Broth. in *Engler's Bot. Jahrb.* 1897, p. 235.

Fr. ign. Cortic. — *Afr.* : Cameroon.

*flavissimus* (C. M.) Par. *Ind.* p. 247 (1894).

*Dicranum flavissimum* C. M. in *Fl.* 1885, p. 409.

Ster. — *Am. merid.* : Fuegia.

*flavo-nigritus* Dusen M. *Am. austr. n.* 610.

Terr. — *Am. merid.* : Patagon. occid.

*flavo-viridis Dusen in Coll. cit. n. 103.*

Terr. — *Am. merid.* : Patagon occid.

*flexifolius (Hsch.) Bryol. jav. I, p. 76 (1855-61).*

*Dicranum flexifolium Hsch. in Schwaeagr. Suppl. II, P. II, II, p. 115, t. 185 (1826); C. M. Syn. I, p. 399 (1849).*

*D. flexuosum Hsch. in Nov. Act. Acad. Leop. XIV, II, Suppl. p. 703 ?*

— *As.* : Java.

*flexuosus (L.) Brid. Mant. M. p. 71 (1819) et Bryol. univ. I, p. 469 (1826); Bryol. eur. fasc. 41 (Vol. I), Mon. p. 3, t. 89 (1848); W. P. Sch. Syn. II ed. p. 102 (1876).*

*Bryum flexuosum L. Sp. pl. p. 1118 (1753).*

*Br. pilosum molle, setis intortis Dill. Hist. M. p. 373, t. 17, f. 33 A-E (1741) p.p.*

*Br. trichoides, capitulis erectis, pediculis intortis tenuibus, virentibus Dill. Cat. Giss. p. 225 (1718).*

*Campylopus Arduennae Lib. Pl. crypt. Ard. n. 106.*

*Dicranum flexuosum Brid. Musc. rec. II, P. I, p. 163 (1792) excl. syn.*

*D. palustre la Pyl. in Brid. Bryol. univ. I, p. 814 (1826) (pd. C. M.).*

*Thysanomitrium flexuosum W. Arn. Dispos. M. p. 33 (1825) excl. var.*

*Weisia immersa Brid. Sp. M. I, p. 121 (1806).*

Ad terr. et saxa silic. — *Eur.* : Reg. campestr. montos. silvat.

*As.* : Borneo ? *Afr.* : Alger., Abyss. *Pac.* : Austr. or. ? (cf. *C. torquatus*).

« *var. fayalensis Card. in Ann. Rep. of the Missouri bot. Gard. 1897, p. 54.*

— *Afr.* : Ins. Azor.

« *var. flagelliferus Ren. in Del. Fl. crypt. Belg. M., p. 281 (1883).*

— *Eur.* : Belg.

*var. major Boul. — C. flexuosus var. zonatus.*

« *var. pachyneuros Mol. Moosst. Alg. Alp. 1865.*

Ut typ. — *Eur.* : Alp. Algov., Bavar.

« *var. paludosus W. P. Sch. in Braithw. Brit. Moosst. I, p. 133 (1882).*

— *Eur.* : Anglia.

*var. paradoxus Husn. — C. paradoxus.*

« *var. uliginosus Ren. in Rev. bryol. 1897, p. 81.*

In turf. — *Eur.* : Gall. centr. (Arvernica), Pyren.

« *var. zonatus (Mol.) Limpr. Laubm. Deutschl. ... I, p. 391 (1887).*

*C. flexuosus* β *major Boulay M. Fr. p. 511 (1881).*

*C. zonatus Mol. Bayern's Laubm. p. 53 (1875).*

*flexuosus* { *Brid. p.p. — C. torfaceus.*  
 { *Mont. — C. caudatus.*  
 { *Sull. — Dicranodontium Millspaughii.*

*fragilis (Dicks.) Bryol. eur. fasc. 41 (Vol. I), Mon. p. 4, t. 90 (1848);*

*W. P. Sch. op. et l. cc.*

*Bryum fragile Dicks. Pl. crypt. fasc. III, p. 5 (1793).*

*C. densus* β *fragilis Wils. Bryol. brit. (1855).*

*C. penicillatus Brid. Mant. M. p. 73 (1819).*

*Dicranum densum Funck Cent. n. 631.*

*D. flexuosum* β *fragile Turn. Musc. hib. p. 74 (1801).*



*D. Funckii* C. M. *Syn.* I, p. 392 (1849).

*Thysanomitrium flexuosum* var. *saxicola* Rab. *Deutsch. Kryptfl.* II, 3, p. 149 (1848).

In fiss. rup. arenac. et granit. — *Eur.* : Britann., Voges., Silv. nigr., Hercyn., Thuring., Bohem., Salisb., Stir., Francon., Bavar., Tirol., Helvet., Armoric., Lusitan., Ligur. As. : Caucas. : M. Neilherr. ?  
*Afr.* : Madeir.

\* var. *densus* (Schleich.) W. P. Sch. in litt. (cf. *Mol. Bryog. Reisebild.* in *Fl.* 1863).

*C. densus* *Bryol. eur. l. c.* p. 6, t. 93.

*C. fragilis* var. *densus* Wulfsb. in *Christian. Vid. Sellsk. Foerh.* 1875 ?

*C. fragilis* forma *elatio* Ltz. *Bryol. Notizb.* p. 88 (1865).

*Dicranum densum* Schleich. *Exs. et Cat.* 1807.

*D. Schleicheri* C. M. *op. cit.* p. 393.

? *Thysanomitrium* piriforme  $\beta$  *penicillatum* Rab. *op. cit.* p. 150.

In turf. — *Eur.* : Helvetia, etc.

*friabilis* (Hpe.) Jaeg. *Ad. I*, p. 136 (1870-71).

*Dicranum friabile* Hpe. *mss.* ; C. M. in *Bot. Zeit.* 1859, p. 220.

— *Am. sept.* : Costa Rica.

*frigidus* Lesq. in *Port. et Coult. Fl. of Colorado*, p. 155 ; *Lesq. et Jam. Man.* p. 79 (1884).

Ster. — *Am. sept.* : Summ. m. Colorado.

*fulvus* Kindb. — *Dicranum*.

*Fumarioli* C. M. in *Abhandl. Nat. Ver. Brem.* 1900, p. 498.

— *Pac.* : Ins. Hawai.

*funalis* Brid. — *Grimmia*.

*fuscatus* Besch. in *Journ. de bot.* 1891, p.

Ster. — *Am. merid.* : Monte Video.

*fusco-croceus* (Hpe.) Jaeg. *Ad. II*, p. 646 (1877-78).

*Dicranum fusco-croceum* Hpe. *Symb.* 1872, p. 45.

Ster. — *Am. merid.* : Brasil. austr. or.

*Gallienii* Par. in *Rev. bryol.* 1900, p. 90.

Ster. — *Afr.* : S. Madagasc.

*Gardneri* (C. M.) Mitt. *M. austr. am.* p. 83 (1869).

*Dicranum flexuosum* var. *Wils. in Gardn. M. brasil. n.* 20.

*D. Gardneri* C. M. in *Bot. Zeit.* 1857, p. 379.

Rupic. — *Am. merid.* : Brasil.

*gastro-alaris* (C. M.) Par.

*Dicranum gastro-alare* C. M. in *Hedwig.* 1900, p. 256.

Ad Cataract. — *Am. merid.* : Brasil. or. (Minas Geraës).

*Gaudichaudi* Besch. in *Journ. de bot.* 1891, p.

Ster. — *Am. merid.* : Chile.

*gemmatus* (C. M.) Par. *Suppl. Ind.* p. 92 (1900).

*Dicranum gemmatum* C. M. in *Bull. hb. Boissier*, 1898, p. 31 et in *Ule Coll. M. brasil. n.* 1789.

— *Am. merid.* : Brasil. or. (S. do Itatiaia).

*geraënsis* Par.

*Dicranum caldense* C. M. in *Hedwig.* 1900, p. 250.

Ster. — *Am. merid.* : Brasil. or. (Minas Geraës).

- giganteus* Sull. — *C. penicillatus*.  
*glauco-viridis* (C. M.) *Par. Suppl. Ind.* p. 92 (1900).  
*Dicranum glauco-viride* C. M. in *Hedwig*. 1897, p. 350.  
 Ster. — *Pac.* : Ins. Kermadec, Sunday.  
*Glaziowii* (Hpe.) *Jaeg. Ad. II*, p. 649 (1877-78).  
*Dicranum Glaziowii* Hpe. *Symb.* 1874, p. 488.  
 — *Am. merid.* : Brasil. austr. or.  
*Goughii* (Mitt.) *Jaeg. Ad. I*, p. 128 (1870-71).  
*C. compactus* Wils. in *Kew Journ. bot. IX*, p. 297 p.p. (1857).  
*Dicranum Goughii* Mitt. *M. Ind. or.* p. 17 (1859).  
 — *As.* : Nepal., Sikkim, Khasia, Bhotan, M. Neil-  
 gherr., Ceylan.  
*gracilicaulis* Mitt. *M. austr. am.* p. 83 (1869).  
 Terr. — *Am. sept.* : Florid. merid. *Am. merid.* : Amazon., And.  
 Peruv.  
*gracilis* (Mitt.) *Jaeg. Ad. I*, p. 131 (1870-71).  
*Dicranum gracile* Mitt. *M. Ind. or.* p. 17 (1859).  
*D. sricum* Wils. in *Kew Journ. bot. IX*, p. 297 (1857).  
 — *As.* : Sikkim.  
*grimmiioides* (C. M.) *Par. Ind.* p. 249 (1894).  
*Dicranum grimmiioides* C. M. in *Linn.* 1880-83, p. 397.  
 Ster. — *Am. merid.* : Argentina. Cordob.  
*griseolus* (C. M.) *Par. Suppl. Ind.* p. 92 (1900).  
*Dicranum griseolum* C. M. in *Hedwig*. 1899, p. 80.  
 Ster. — *Afr.* : Transvaal.  
*griseus* (Hsch.) *Jaeg. Ad. I*, p. 147 (1870-71).  
*Dicranum griseum* C. M. *Syn. I*, p. 406 (1849).  
*Thysanomitrium griseum* Hsch. *Fl. brasil. I*, p. 16 (1840).  
 Rupic. — *Am. merid.* : Brasil. austr. or., Monte Video.  
*guadalupensis* (Brid.) Mitt. *M. austr. am.* p. 77 (1869).  
*Dicranum guadalupense* Brid. *Bryol. univ. I*, p. 463 (1826); *C. M. Syn. I*,  
 p. 386 (1849).  
 ♂ et fr. ign. — *Am. sept.* : Guadalup.  
*Guatucæ* Dusen M. *Am. austr. n.* 644.  
 Ad rup. littor. — *Am. merid.* : Patagon. occid.  
*Hallii* Lesq. in *Port. et Coult. Fl. of Colorado*, p. 155; *Lesq. et Jam.*  
*Man.* p. 79 (1884).  
 Ster. — *Am. sept.* : Summ. m. Colorado.  
*Harrisii* (C. M.) *Par. Suppl. Ind.* p. 92 (1900).  
*C. lamellinerve* Mitt. *M. austr. am.* p. 82 (*quoad pl. jamaicens.*).  
*Dicranum Harrisii* C. M. in *Bull. hb. Boissier*, 1897, p. 553.  
 Terr. — *Am. sept.* : Jamaic.  
*Hartmanni* W. P. Sch. — *Tortella fragilis*.  
*hawaïco-flexuosus* (C. M.) *Par. op. et l. cc.*  
*Dicranum hawaïco-flexuosum* C. M. in *Bull. hb. Boissier*, 1897, p. 143.  
 — *Pac.* : Ins. Hawaï.

*helenicus* (C. M.) Par.

C. Sanctae Helenae W. P. Sch. in sched.

Dicranum helenicum C. M. Syn. II, p. 599 (1851).

Filicic. — Afr. : Ins. Ascension., S. Helen.

*Hellerianus* (Hpe.) Jaeg. Ad. I, p. 121 (1870-71).Dicranum Hellerianum Hpe. Spec. nov. Mexic. in Verh. d. K.K. zool.-bot. Gesells. in Wien, 1<sup>a</sup> April. 1869.

Ster. — Am. sept. : Mexic.

*hemitrichius* (C. M.) Jaeg. Ad. II, p. 648 (1877-78).

Dicranum hemitrichium C. M. in Linn. 1874, p. 553.

Pilopogon ? (fid. Brotherus).

Ster. — As. : Ins. Philippin.

*Henrici* Ren. et Card. in Bot. Gaz. 1888, p. 197. t. 14.

Ster. — Am. sept. : Kansas.

*Hensii* Ren. et Card. in Bull. Soc. roy. bot. Belg. 1896, I, p. 301.

Ster. Ad terr. arenac. — Afr. : Congo belg.

*Heribaudi* Ren. et Card. in Bull. Soc. roy. bot. Belg. 1890, I, p. 168 ;

Ren. M. Masc. Mad. n. 105.

— Afr. : Madagasc.

*heterophyllus* Mitt. M. austr. am. p. 77 (1869).

— Am. merid. : And. Nov. Granat.

*heterostachys* Hpe. in Fl. 1865, n. 37.

— Am. merid. : And. Peruv.

*Hildebrandtii* (C. M.) Jaeg. Ad. II, p. 646 (1877-78).

Dicranum Hildebrandtii C. M. in Linn. 1876, p. 236.

Humic. — Afr. : Ins. Comor.

*hispidus* Ren. et Card. in Bull. Soc. roy. bot. Belg. 1890, I, p. 171.

Ster. Ad terr. argill. silic. — Afr. : Madagasc.

*Hoehnelii* (C. M.) Par. Ind. p. 250 (1894).

Dicranum Hoehnelii C. M. in Fl. 1890, p. 473.

Ster. In lapidos. — Afr. : Kilima N'Djaro.

*Holomitrium* (C. M.) Jaeg. Ad. I, p. 135 (1870-71).

C. caespiticius H. f. et W. in Lond. Journ. of bot. 1814, p. 543.

D. Holomitrium C. M. Syn. I, p. 389.

— Pac. : Nov. Seeland.

*Hoffmanni* (C. M.) Ren. et Card. in Bull. Soc. roy. bot. Belg. 1892, I, p. 147.

Dicranum Hoffmanni C. M. in Linn. 1874, p. 592 ; Par. Ind. p. 355 (1895).

Ster. Paludic. — Am. sept. : And. Costa Ricens.

*homalobolax* (C. M.) Par. Suppl. Ind. p. 92 (1900).

Dicranum homalobolax C. M. in Hedwig. 1897, p. 349.

Ster. — Pac. : Fret. reg. Georges.

*horridus* Welw. et Duby in Mém..... Genève, 1870, p. 8, t. 4, f. I.

— Afr. : Angola, Loanda.

*humifugus* (C. M.) Par. Ind. p. 250 (1894).

Dicranum humitugum C. M. in Linn. 1880-82, p. 400.

Lignic. — Am. merid. : Argentin. subtropic.

humilis *Mont. in Ann. sc. nat.* 1845, IV, p. 110 et *Syll. p.* 43 (1856);  
*Mitt. M. austr. am. p.* 76 (1869).

*Dicranum humile C. M. Syn. I, p.* 384 (1849).

Lignic. — *Am. merid.* : Brasil.

humoricola (*C. M.*) *Par.*

*Dicranum humoricola C. M. in Hedwig.* 1900, p. 256.

Ad rup. humid. — *Am. merid.* : Rio de Janeiro.

hygrometricus *Rehm.* — *C. purpureo-aureus.*

Imandae *Rehm. M. Afr. austr. n.* 43.

*Dicranum Imandae C. M. in Hedwig.* 1899, p. 95.

Ster. — *Afr.* : Natal.

Imerangae *Rehm. in Coll. cit. n.* 42.

*C. Incrangae Par. Ind. p.* 251 (erratum !)

*Dicranum Imerangae C. M. in op. cit. p.* 83.

Ster. — *Afr.* : Natal.

incrassatus (*Kze.*) *C. M. in Linn.* 1844, p. 686; *Mitt. M. austr. am. p.* 89 (1869).

*Dicranum incrassatum Kze. in Poeppig. Pl. chil. n.* 22; *C. M. Syn. I, p.* 408 (1849).

Saxic. — *Afr.* : Chile.

insititius *Hook. f. et Wils. Fl. Tasman. II, p.* 172, t. 172, f. 3 (1860).

*C. clavatus eor. (fid. Mitt.).*

— *Pac.* : Tasman.

intermedius *Wils.* — *Dicranodontium longirostre var. alpinum.*

interruptulus *C. M. in Besch. Fl. bryol. Réunion, etc. p.* 36 (1880-81).

Ster. — *Afr.* : Ins. Franciae.

introflexus (*Hedw.*) *Mitt. M. austr. am. p.* 81 (1869); *Duby Choix de Crypt. ex.* 1875, p. 13, t. 1, f. 5 (1).

*C. leucotrichus Sull. et Lesq. in Sull. Ic. M. p.* 28, t. 17 (1861) et *M. bor am. II ed. n.* 92.

*Dicranum introflexum Hedw. Sp. M. p.* 147, t. 29 (1801); *C. M. Syn. I, p.* 405 (1849).

Rupic. et terr. — *Afr.* : Ins. Canar., Madeir., Ascens., S. Helen, Tri-tan da Cunha. *Am. sept.* : Alabama. *Am. merid.* : Brasil. austr. or., Monte Video, Chile, ins. Maclov. *Pac.* : Austral. or., fret. reg. Georges, Nov. Seeland., Tasman., ins. Auckland, Campbell.

• *var. sublævopilus Kindb. in Rev. bryol.* 1893, p. 91.

— *Afr.* : Ins. Azor. (S. Michel).

---

(1) Cl. Mitten (l. c.) huic speciei « very variable in external appearance », synonym. sequent. ducit :

*C. grisens Hsch.*  
*C. lamellatus Mont.*  
*C. lutescens W. P. Sch.*  
*C. polytrichoides de Not.*  
*C. xanthophyllus Mont.*

*Dicranum leptoccephalum C. M.*  
*D. truncatum C. M.*  
 Et forsan :  
*C. canescens W. P. Sch.*  
*C. capensis W. P. Sch.*

*introflexus* { Brid. p.p. — *C. aureo-nitens*.  
 Mitt. — *C. eximius*.  
 Mont. — *C. Berteroanus*.  
*irregularis* Ltz. — *Leptotrichum homomallum* var. *zonatum*.  
*Jamesoni* Jaeg. — *C. concolor*.

*japonicus* Broth. in *Hedwig*. 1899, p. 207.  
 Ster. Rupic. — As. : Japon.

*Johannis Meyeri* (C. M.) Par. Ind. p. 251 (1894).  
 Dicranum *Johannis Meyeri* C. M. in *Fl.* 1888, p. 410.  
 Ster. Rupic. — Afr. : Kilima N'Djaro.

*Joinvillianus* (Hpe.) Jaeg. Ad. II, p. 646 (1877-78).  
 Dicranum *Joinvillianum* Hpe. *Symb.* 1872, p. 45.  
 — Am. merid. : Brasil. austr. or.

*Jucrangae* Par. — *C. Imerangae*.

*julaceus* (Hpe.) Jaeg. op. cit. p. 648.  
 Dicranum *julaceum* Hpe. op. cit. p. 46.  
 ♂ et fr. ign. — Am. merid. : Brasil. austr. or.

*julaceus* Rehm. — *C. pseudo-julaceus*.

*Kirkii* Mitt. in *Trans. and Proced. of the N. Zeal. Inst.* 1893, p. 230.  
 Dicranum *Kirkii* Broth. et Geh. in *Rev. bryol.* 1897, p. 67.  
 — Pac. : Nov. Seeland.

*Krauseanus* (Hpe. et Ltz.) Par. Ind. p. 252 (1894).  
 Dicranum *areodictyon* Hpe. et Ltz. in *Krause M. Ecuador*.  
*D. Krauseanum* cor. in *Bot. Zeit.* 1869, p.  
 Rupic. et sphagnic. — Am. merid. : And. Quitens.

*Kunerti* Par. *Suppl. Ind.* p. 93 (1900).  
 Dicranum *pseudo-julaceum* C. M. in *Bull. hb. Poissier*, 1898, p. 38 (nom.)  
 et in *Hedwig*. 1900, p. 259.  
 Paludic. — Am. merid. : Brasil. austr. or. (Rio Grande So Sul).

*laete-virens* (C. M.) Jaeg. Ad. I, p. 128 (1870-71).  
 Dicranum *laete virens* C. M. in *Bot. Zeit.* 1862, p. 12.  
 Ster. — Afr. : Ins. Canar.

*laetus* (Mitt.) Jaeg. op. cit. p. 121.  
 Dicranum *laetum* Mitt. M. Ind. or. p. 19 (1859).  
 Ad arb. et rup. — As. : Khasia.

*laevigatus* Brid. — *Grimmia leucophaea*.

*laevis* Tayl. in *Lond. Journ. of bot.* 1846, p. 47; Mitt. M. austr. am.  
 p. 91 (1869).  
 Dicranum *laeve* C. M. *Syn. I*, p. 409 (1849).  
*Pilopogon piliferus* Hpe. in *Ann. sc. nat.* 1866, p. 362 (ex ips. in *Bot.*  
*Zeit.* 1859, n. 52).  
 — Am. merid. : And. Nov. Granat. et Quitens.,  
 M. Chimborazo et Pichincha.

*lamellatus* Mont. in *Ann. sc. nat.* 1838, IX, p. 52 et in *d'Orbigny Voy.* p. 90 (1839); *Syll.* p. 43 (1856).

*C. introflexus* Mitt. (*ex ips.*!).

*Dicranum lamellatum* C. M. *Syn. I*, p. 411 (1849).

Terr. — *Am. merid.* : Peruv.

*lamellicosta* W. P. Sch. in *Besch. Prodr. fl. bryol. mexic.* p. 23 (1871).

*Dicranum lamellicosta* C. M. *Syn. II*, p. 60 (1851).

— *Am. sept.* : Mexic.

*lamellinervis* Mitt. { — *C. Harrisii*.  
— *C. penicillatus*.

*lamprodictyon* (Hpe.) Mitt. *M. austr. am.* p. 78 (1869).

*Dicranum lamprodictyon* Hpe. in *Ann. sc. nat.* 1866, p. 339.

— *Am. merid.* : And. Nov. Granat.

*laniger* Besch. in *Bull. Soc. bot. Fr.* 1885, p. LVI et in *Mission sc. Cap Horn.*, Bot, p. 262 (1889).

*Dicranum laniger* C. M. in *Fl.* 1885, p. 410.

Ster. Cort. et lignic. — *Am. merid.* : Chile, Patagon. occid., Fuegia, ins. Eremitae.

*lapidicola* (C. M.) Par.

*Dicranum lapidicola* C. M. in *Hedwig.* 1900, p. 258.

Ster. Saxic. — *Am. merid.* : Brasil. austr. or. (ins. S. Catharina).

*latinervis* (Mitt.) Jaeg. *Ad. I*, p. 130 (1870-71).

*Dicranum latinerve* Mitt. *M. Ind. or.* p. 17 (1859).

— *As.* : Nepal. or., Sikkim, m. Neilgherr., Ceylan.

*laxitextus* Lac. *Sp. nov. M. Archip. ind.* p. 10, t. 7 A. (1872).

Terr. — *As.* : Java.

*laxobasis* Ren. et Card. in *Bull. soc. roy. bot. Belg.* 1896, I, p. 299.

— *Afr.* : Madagasc.

*Leanus* Sull. — *Brothera*.

*Lechleri* W. P. Sch. in *Lechler Pl. Chil.*

Ad arb. et terr. — *Am. merid.* : Chile austr.

*lepidophyllus* (C. M.) Jaeg. *Ad. I*, p. 139 (1870-71).

*C. capensis* W. P. Sch. in *Breutel M. capens.*

*Dicranum lepidophyllum* C. M. *Syn. I*, p. 407 (1849).

*D. pudicum* Hsch. in *Drege M. capens.*

— *Afr.* : Pr. B. Sp.

*leptocephalus* (C. M.) Jaeg. *Ad. II*, p. 435 (1876-77).

*C. introflexus* (*fid. Mitt.*).

*Dicranum leptocephalum* C. M. in *Linn.* 1855, p. 206 (*nom.*).

Terr. — *Pac.* : Austral. or., Nov. Seeland.

*leptodrepanium* (C. M.) Broth. *Natür. Pflanzenfam.* p. 332.

*Dicranodontium leptodrepanium* C. M. in *Dusen M. Camer. n.* 348 ; *Broth. in Engler's Bot. Jahrb.* 1897, p. 233.

2. Ad ramul. — *Afr.* : Cameroon.

*leptodus* Mont. p.p. — *Pilopogon Richardi*.

- leptotrichaceus* (C. M.) Par. *Suppl. Ind.* p. 93 (1900).  
*Dicranum leptotrichaceum* C. M. in *Hedwig*. 1899, p. 84.  
 Ster. — Afr. : Pr. B. Sp.
- leucobasis* (C. M.) Par. *op. et l. cc.*  
*Dicranum leucobasis* C. M. in *Hedwig*. 1899, p. 78.  
 Ster. — Afr. : Pr. B. Sp.
- « *var. bartramiaeus* (C. M.) Par. *op. et l. cc.*  
*C. subbartramiaeus* C. M. in *Rehm. M. Afr. austr. n.* 32 ; Par. *Ind.* p. 261 (1894).  
*Dicranum leucobasis var. bartramiaeus* C. M. in *op. cit.* p. 86 et in *Coll. cit.* n. 37.  
 Ubi typ.
- « *var. longescens* (C. M.) Par. *op. et l. cc.*  
*C. Vallis Gratiae* Hpe. in *Breutel M. capens.*  
*Dicranum leucobasis var. longescens* C. M. in *op. et l. cc. et in Coll. cit.* n. 79.  
 Ubi typ.
- leucochlorus* (C. M.) Par. *Ind.* p. 253 (1894).  
*Dicranum leucochlorum* C. M. in *Fl.* 1888, p. III.  
 Ster. — Afr. : Kilima N'Djaro.
- leucogaster* (C. M.) Par. *op. et l. cc.*  
*Dicranum filiforme* Hsch. in *Deppe et Schiede M. mexic.*  
*D. leucogaster* C. M. *Syn. I*, p. 387 (1849).  
 Ster. — Am. sept. : Mexic.
- leucognodes* (C. M.) Par. *Suppl. Ind.* p. 93 (1900).  
*C. areodictyon* W. P. Sch. in *Mandon Pl. Boliv. n.* 1612.  
*Dicranum leucognodes* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 32.  
 Cortic. — Am. merid. : And. Boliv.  
 « *var. gracilescens* (C. M.) Par. *op. et l. cc.*  
*C. areodictyon var. gracilescens* W. P. Sch. in *Coll. cit.* n. 1613.  
*Dicranum leucognodes var. gracilescens* C. M. in *op. et l. cc.*  
 Ubi typ.
- leucotrichus* Sull. et Lesq. — C. *introflexus*.
- lonchochacte* C. M. in *Abhandl. d. Natur. Ver. Bremen*, 1900, p. 495.  
 — Pac. : Nov. Seeland.
- lonchoclados* C. M. in *Besch. Fl. bryol. Réunion, etc.* p. 33 (1880-81).  
 Rupic. — Afr. : Usambara, ins. Borbon., Francia.
- longescens* Rehm. M. *Afr. austr. n.* 40.  
*Dicranum longescens* C. M. in *Hedwig*. 1809, p. 85.  
 Ster. — Afr. : Pr. B. Sp. (Knysma).
- « *var. fimus* Rehm. in *Coll. cit.* n. 41.  
 Ubi typ.
- longifolius* W. P. Sch. in *Besch. Fl. bryol. Réunion, etc.* p. 35 (1880-81).  
*Cecalyphum fasciatum* P. B. *Prodr.* p. 51 ? (1805).  
*Dicranum arcuatum* Brid. *Bryol. univ. I*, p. 463 p.p. (1826) ; C. M. *Syn. I*, p. 368 p.p. (1849).  
 Ster. — Afr. : Ins. Borbon. Pac. : Ins. austr. ? (*fid. P. B.*).

*longifolius* Kindb. -- Dicranum.

*longipilus* Brid. *Bryol. univ. I*, p. 477 p.p. (1826); *Bryol. eur. Suppl. I-II*, t. 3, (1864); W. P. Sch. *Syn. II ed.* p. 104 (1876).

*C. atro-virens* de Not. *Syll. M.* p. 221 (1838) et *Epil.* p. 648 (1869), non *Bryol. eur. I*, t. 92, f. I (1847).

Dicranum *atro-virens* C. M. *Syn. I*, p. 411 (1849).

*D. flexuosum* {  $\beta$  nigro-viride H. et T. *Muscol. brit. II ed.* p. 94 p.p. (1827).  
                   {  $\gamma$  piliferum Turn. *Muscol. hib.* p. 74 p.p., t. 5, f. 2 b.c (1801).

$\sigma$  et fr. ign. Rupic. — Eur. : Norveg. occid., Britann., Ital. super., Tirol., Pyren., Armoric. Afr. : Cameroon (*fid. Mitt.*)

$\alpha$  var. *falcata* (Ferg.) Par. *Ind.* p. 254 (1894).

*C. atro-virens* var. *falcatus* Braithw. in Husn. *Muscol. gall.* p. 46 (1884).

*C. falcatus* Ferg. mss.

Ubi typ.

$\alpha$  var. *muticus* (Milde) Pur.

*C. atro-virens* var. *epilosus* Braithw. *Brit. Moossf. I, Suppl...* p. 298 (1887).

*C. atro-virens* var. *muticus* Milde in *Bot. Zeit.* 1870, p. 396.

— Eur. : Anglia, Siles.

*longipilus* Brid. p.p. — *C. polytrichoides*.

var. de Not. — *C. polytrichoides*.

Lorentzi C. M. in Linn. 1880-83, p. 404.

Fl. et fr. ign. — Am. merid. : Alp. Argentin. subtropic.

*luridus* W. P. Sch. in Besch. *Prodr. bryol. mexic.* p. 24 (1871).

*Pilopogon* ? (*fid. Broth.*).

Ster.

— Am. sept. : Mexic. (M. Orizaba).

*lutescens* W. P. Sch. in Besch. *op. et l. cc.*

Dicranum *lutescens* C. M. *Syn. II*, p. 602 (1851).

— Am. sept. : Mexic.

*luteus* (C. M.) Par. *Ind.* p. 254 (1894).

*Thysanomitrium luteum* C. M. in Linn. 1878-79, p. 470.

— Am. merid. : Venezuela.

*Mac-Leanus* Rehm. M. Afr. austr. n. 447.

— Afr. : Transvaal.

*macrogaster* (C. M.) Par.

Dicranum *macrogaster* C. M. in Hedwig. 1900, p. 252.

Rupic. — Am. merid. : Brasil. austr. or. (ins. S. Catharina).

*macrophyllus* (Kze.) Mitt. M. austr. am. p. 80 (1869).

Dicranum *filifolium* C. M. in Linn. 1844, p. 691 p.p.

*D. macrophyllum* Kze. in Poeppig Coll. III, n. 231; C. M. *Syn. I*, p. 402 (1849).

— Am. merid. : Peruv.

*macrotris* C. M. in Dusen M. Camer. n. 699; Broth. in Engler's Bot. Jahrb. 1897, p. 233.

Ster. Cortic. — Afr. : Cameroon.

? *maguiretis* (C. M.) Par. *Suppl. Ind.* p. 94 (1900).

Dicranum *maguirets* C. M. in Bull. hb. Boissier, 1897, p. 186.

Cortic. — Am. sept. : Guatemala.



- madecassus *Besch. Fl. bryol. Réunion, etc.* p. 33 (1880-81).  
— *Afr.* : Madagasc.
- Mandoni *W. P. Sch. in Mandon Pl. Boliv. n.* 1610.  
— *Am. merid.* : And. Boliv.
- Marilacii *Rehm. M. Afr. austr. n.* 65.  
Ster. — *Afr.* : Pr. B. Sp. (Montagu).
- matarensis *Besch. Fl. bryol. Réunion, etc.* p. 35 (1880-81).  
Ster. — *Afr.* : Ins. Borbon.
- megalotus *Besch. Not. M. Ins. S. Paul et Amsterdam, p.* 2 (1875).  
Ster. — *Afr.* : Ins. S. Paul.
- micans *Wulfsb. in Christian. Vid. Sellsk Foerh.* 1875 *et in Rev. bryol.* 1885, p. 15; *Hagen in Det. K. Norske Vid. dels Skrift.* 1897, n. 2.  
Rupic. — *Eur.* : Norveg.
- microcephalus (*C. M.*) *Par. Suppl. Ind. p.* 94 (1900).  
Dicranum microcephalum *C. M. in Fl.* 1896, p. 443.  
— *Pac.* : Ins. Hawai.
- microjulaceus (*C. M.*) *Par. op. et l. cc.*  
Dicranum microjulaceum *C. M. in Bull. hb. Boissier,* 1897, p. 37 *et in Ule Coll. M. brasil. n.* 1786.  
In decliv. — *Am. merid.* : Brasil. (S. do Itatiaia).
- microstomus *Mitt. — C. porphyreodictyon.*
- Mildei *Limpr. Laubm. Deutschl...* I, p. 396 (1887).  
C. brevifolius { *de Not. Epil. p.* 650 (1869) *et in Erb. critt. ital. n.* 1142.  
                          { forma Milde *in Bot. Zeit.* 1870, p. 459.  
C. polytrichoides var. { *Daldiniana de Not. mss.; Jaeg. Ad. I, p.* 145 (1870-71).  
                                  { *Mariana Amann in sched.*  
Dicranum viridissimum *Rosa in de Not. Epil. p.* 751 ?  
Ster. Terr. — *Eur.* : Ticin., lac. Major.
- Minarum *Par.*  
Dicranum laxobasis *C. M. in Hedwig.* 1900, p. 252.  
Rupic. — *Am. merid.* : Brasil. or. (Minas Geraës).
- minor *Besch. Not. M. Ins. S. Paul et Amsterdam, p.* 4 (1875).  
— *Afr.* : Ins. Amsterdam.
- Mittenii (*C. M.*) *Jaeg. Ad. I, p.* 137 (1870-71).  
Dicranum erectum *Mitt. in Lond. Journ. of bot.* 1851, p. 53.  
D. Mittenii *C. M. Syn. II, p.* 600 (1851).  
— *Am. merid.* : And. Quitens. et Boliv.
- Molkenboëri *Lac. — C. brevipilus.*
- mollis *C. M. in Rehm. M. Afr. austr. n.* 36.  
Ster. — *Afr.* : Natal.
- montanus *Welw. et Duby in Mém.... Genève,* 1870, p. 7, t. 3, f. 1.  
Ster. Lignic. et rupic. — *Afr.* : Angola.
- Morenoi (*C. M.*) *Par. Suppl. Ind. p.* 94 (1900).  
Dicranum Morenoi *C. M. in Hedwig.* 1897, p. 97.  
Ster. — *Am. merid.* : Patagon.
- Moseni *Broth. in Bihang till K. Sv. Vet.-Akad. Foerh.* 1895, n. 3.  
Ad rup. aprie. — *Am. merid.* : Brasil. or.

- Mülleri* Jur. { *Jaeg.* — *Pilopogon* Richardi.  
 { *Milde.* — *C. torfaceus* var. *Mülleri*.
- multicapsularis* W. P. Sch. in *Mandon Pl. Boliv. n.* 1611.  
*Dicranum multicapsulare* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 34.  
 — *Am. merid.* : And. Boliv.
- Nadeandianus* Besch. in *Ann. sc. nat.* 1894, p.  
 Ad terr. et rhizom. *Pteridis.* — *Pac.* : *Taiti*.
- nano-filifolius* (C. M.) *Par. Suppl. Ind.* p. 94 (1900).  
*Dicranum nano-filifolium* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 35.  
 — *Am. merid.* : And. Boliv.
- nanophyllus* C. M. in *Dusen M. Camer. n.* 305; *Broth. in Engler's Bot. Jahrb.* 1897, p. 235.  
*Fr. ign. Rupic.* — *Afr.* : *Cameroon*.
- nano-tenax* (C. M.) *Jueg. Ad. II*, p. 761 (1877-78).  
*C. natalensis* Rehm. *M. Afr. austr. n.* 51.  
*Dicranum nano-tenax* C. M. in *Hedwig.* 1899, p. 82.  
 Ster. — *Afr.* : *Zouloulund, Natal*.
- nanus* C. M. in *Bot. Zeit.* 1847, p. 804, p.p.  
*Dicranum flexuosum* var. *Hsch. in Linn.* 1841, p. 124.  
*D. nanum* C. M. *Syn. I*, p. 483 (1849) et in *Bot. Zeit.* 1859, p. 189.  
 — *Afr.* : *Pr. B. Sp.*
- nanus* Bryol. jav. — *Campylopodium euphorocladum*.  
*natalensis* Rehm. — *C. nano-tenax*.
- Naumanni* (C. M.) *Par. Ind.* p. 255 (1894).  
*Dicranum Naumanni* C. M. in *Engler's Bot. Jahrb.* 1883, p. 81 et in *Forschungsreise... Bot.* p. 41 (1889).  
 Ster. — *Afr.* : *Ins. Ascension*.
- nepalensis* (Brid.) *Jaeg. Ad. I*, p. 120 (1870-71).  
 ? *Dicranum nepalense* C. M. *Syn. I*, p. 384 (1849).  
 ? *Didymodon nepalensis* Brid. *Bryol. univ. I*, p. 82.  
 Ster. — *As.* : *Nepal*.
- Nietneri* (C. M.) *Jaeg. Ad. I*, p. 136.  
*Dicranum latinerve* Mitt. *M. Ind. or.* p. 17 p.p. ? (1859) *fid. C. M.*  
*D. Nietneri* C. M. in *Linn.* 1869-70, p. 35.  
 Ster. — *As.* : *Ceylan*.
- nigrescens* Duby — *Pilopogon* Richardi.
- nigro-flavus* (C. M.) *Par. Suppl. Ind.* p. 94 (1900).  
*Dicranum nigro-flavum* C. M. in *Hedwig.* 1897, p. 349.  
 Ster. — *Pac.* : *Austral. occ.*
- nilghiriensis* (Mitt.) *Jaeg. Ad. I*, p. 130 (1870-71).  
*Dicranum nilghiriense* Mitt. *M. Ind. or.* p. 16 (1859).  
*D. nivale* C. M. *Syn. I*, p. 393 p.p. (1849).  
*D. Perrottetii* Mont. in *Ann. sc. nat.* 1843, p. 241 p.p.  
 — *As.* : *M. Neilgherr.*

*nivalis* Brid. *Bryol. univ.* I, p. 477 (1826).

Dicranum nivale C. M. *Syn.* I, p. 393 p.p.; *Mitt. op. et l. cc.*

Weisia nivalis Brid. *Mant. M.* p. 48 (1819).

— Afr. : Cameroon, ins. Borbon.

*nodiflorus* (C. M.) Jaeg. *Ad.* I, p. 144 (1870-71).

Dicranum nodiflorum C. M. in *Bot. Zeit.* 1853, p. 38; *Mitt. M. Ind. or.* p. 21 (1859).

Ster.

— As. : M. Neilgherr.

*Novae Valesiae* Broth. in *Oefv. af Finska Vet.-Akad. Foerh.* 1898, p. 163.

— Pac. : Austral. or. (Nov. Vales.).

*nudicaulis* Besch. in *Bull. Soc. bot. Fr.* 1898, p. 55.

— Pac. : Taïti.

*nudus* (Hpe.) Jaeg. *Ad.* I, p. 125 (1870-71).

Dicranum nudum Hpe. in *Linn.* 1859-60, p. 630.

— Pac. : Austral. or.

*obscurus* Aongstr. in *Oefv. af K. Sv. Vet.-Akad. Foerh.* 1873, n. 5, p. 119.

— Pac. : Taïti.

*occultus* Mitt. *M. austr. am.* p. 86 (1869).

Lignic. — Am. merid. : Brasil. austr. or.

*ochrodictyon* Aongstr. in *op. cit.* p. 146.

— Afr. : Ins. S. Helena.

*Erstedtianus* (C. M.) Mitt. *op. cit.* p. 81.

Dicranum (Erstedtianum) C. M. *Syn.* II, p. 596 (1851).

Ster.

— Am. sept. : And. Costa Ricens.

*olivaceo-nigricans* (C. M.) Par. *Suppl. Ind.* p. 95 (1900).

Dicranum olivaceo-nigricans C. M. in *Hedwig.* 1899, p. 81.

Ster.

— Afr. : Transvaal.

*orthocomus* Besch. in *Bull. Soc. bot. Fr.* 1885, p. LVII et in *Miss. sc. Cap Horn*, V, *Bot.* p. 263, t. 2, f. 5.

Dicranum orthocomum C. M. in *Fl.* 1885, p. 409.

Ster.

— Am. merid. : Fuegia.

*orthopelma* (C. M.) Par.

Dicranum orthopelma C. M. in *Hedwig.* 1900, p. 257.

Ad rup. humid. — Am. merid. : Brasil. austr. or. (ins. S. Catharina).

*orthopodium* (C. M.) Par.

Dicranum orthopodium C. M. in *op. cit.* p. 249.

Terr. ? — Am. merid. : Brasil. austr. or. (ins. S. Catharina).

*ouro-pretensis* Par.

Dicranum flaccidum C. M. in *op. cit.* p. 251.

Ster. Rupic. — Am. merid. : Brasil. or. (Minas Geraës).

*ovalis* Wahlenb. — *Grimmia ovata*.

β Wahlenb. — *Grimmia commutata*.

*ovatus* Brid. — *Grimmia commutata*.

*pachycomus* Besch. in litt.

*C. bicolor* ej. Prodr. bryol. mexic. p. 22 (1871).

— *Am. sept.* : Mexic.

*pachyneuros* Mol. — *Dicranodontium longirostre* var. *alpinum*.

*pallescens* Besch. Fl. bryol. Réunion, etc. p. 32 (1880-81).

— *Afr.* : Ins. Bourbon.

*pallido-nitens* (C. M.) Par.

*Dicranum pallido-nitens* C. M. in Sikora Pl. Mad. n. 14297.

— *Afr.* : Madagascar.

*pallidus* H. f. et W. p.p. — *C. torquatus*.

*patagonicus* Broth. in Dusen M. Am. austr. n. 39.

Paludic. — *Am. merid.* : Patagon. occid.

*paradoxus* Wils. mss. ; Hardy in Berwick Nat. Club. hist. 1868, p. 48 ;  
*Brailho*. in Journ. bot. 1870, p. 390. t. III, f. 2 et in Brit. Moos<sup>9</sup>.  
 p. 133, t. 18 G ; W. P. Sch. Syn. II ed. p. 108.

*C. flexuosus* var. *paradoxus* Husn. Musc. gall. p. 42 (1884).

Ster. Ad terr. humos. torfac. — *Eur.* : Anglia, Belgia, Gallia.

*parvicaespitosus* (C. M.) Par.

*Dicranum parvicaespitosum* C. M. in Hedwig. 1900, p. 259.

Rupic. — *Am. merid.* : Brasil. austr. or. (ins. S. Catharina).

*patens* Brid. — *Rhacomitrium*.

*pauper* (Hpe.) Mitt. M. austr. am. p. 74 (1869).

*Dicranum pauper* Hpe. in Linn. 1863, p. 137.

Terr. — *Am. merid.* : And. Nov. Granat.

« var. *minus* (Hpe.) Jaeg. Ad. I, p. 133.

*Dicranum pauper* var. *minus* Hpe. in op. et l. cc.

Ubi typ.

*penicillatus* (Hsch.) Jaeg. Ad. I, p. 146 (1870-71).

*Bartramia recurvifolia* Duby Choix de M. ex. 1880, p. 2, t. 3 (fd. Hpe. in  
 Fl. 1880. n. 21).

*C. giganteus* Sull. in Proceed. of the Amer. Acad. August. 1861.

*C. lamellinervis* Mitt. M. austr. am. p. 82 p.p. (1869).

*Dicranum concolor* C. M. in Bot. Zeit. 1844, p. 709.

*D. lamellinerve* ej. Syn. I. p. 390 (1849).

*D. penicillatum* Hsch. Fl. brasil. p. 13 (1840) ; C. M. op. cit. p. 415.

Rupic. et terr. — *Am. sept.* : Jamaic. ? Cuba. *Am. merid.* : Brasil.  
 austr. or.

*penicillatus* Rog. — *C. fragilis*.

*perangustifolius* C. M. in Fl. 1886, p. 479.

Ster. — *Pac.* : Ins. Hawaï.

*perauriculatus* Broth. in Oefv. af Finska Vet.-Akad. Foerh. 1899, p. 93.

Ster. — *Pac.* : Austral. or. (Nov. Vales.).

*percurvatus* C. M. in Fl. 1897, p. 34.

— *Am. merid.* : And. Venezuel.

*perexilis* (C. M.) Par. Ind. p. 95 (1900).

*Dicranum perexile* C. M. in Nuov. Giorn. bot. ital. 1897, p. 34.

— *Am. merid.* : And. Boliv.

*perincanus* (C. M.) *Par. Ind.* p. 257 (1894).

*Dicranum perincanum* C. M. in *Fl.* 1885, p. 410.

Terr. — *Am. merid.* : Fuegia.

*perpusillus* Mitt. in *Journ. of the Linn. Soc.* 1886, p. 301.

*D. dicranelloides* R. C. ?

— *Afr.* : Ougogo.

*perreduncus* (C. M.) *Par. Suppl. Ind.* p. 95 (1900).

*Dicranum perreduncum* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 33.

— *Am. merid.* : And. Boliv.

*persimplex* C. M. in *Abhandl. d. Natur. Ver. zu Bremen*, 1900, p. 496.

— *Pac.* : Nov. Seeland.

*pertristis* (C. M.) *Par. op. et l. cc.*

*Dicranum pertriste* C. M. in *Fl.* 1886, p. 444.

Ster. — *Pac.* : Ins. Hawai.

*Pes Funariae* (C. M.) *Par. Ind.* p. 257 (1894).

*Thysanomitrium Pes Funariae* C. M. in *Linn.* 1874, p. 551.

— *As.* : Ins. Philippin.

*pilifer* Brid. — *C. polytrichoides*.

*pilosissimus* W. P. Sch. in *Besch. Prodr. bryol. mexic.* p. 24 (1871).

*Pilopogon* ? (*fid. Broth.*).

Ster. — *Am. sept.* : Mexic.

*piriformis* Brid. — *C. torfaceus*.

var. *Mülleri* Braithw. — *C. torfaceus* var. *Mülleri*.

*platyneuron* (Hpe.) *Par. Ind.* p. 257 (1894).

*Dicranum platyneuron* Hpe. *Symb.* 1874, p. 489.

— *Am. merid.* : Brasil. austr. or.

*pleurocarpus* (C. M.) *Par. Suppl. Ind.* p. 95 (1900).

*Dicranum pleurocarpum* C. M. in *Bull. hb. Boissier*, 1898, p. 35 et in *Ulc Coll. M. brasil.* n. 1788.

In decliv. — *Am. merid.* : Brasil. or. (S. do Itatiaia).

*poasensis* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1892, 1, p. 148.

— *Am. sept.* : And. Costa Ricens.

*poicilophyllus* (C. M.) *Par. op. et l. cc.*

*Dicranum poicilophyllum* C. M. in *Sikora Pl. Mad.* n. 14298.

— *Afr.* : Madagascar.

*polytrichoides* de Not. *Syll. M.* p. 222 (1838) et *Epil.* p. 645 (1869); *Bryol. eur. Suppl.* 1-2, t. 4, (1864); W. P. Sch. *Syn. II ed.* p. 104 (1876).

*C. ericetorum* Jaeg. *Ad. II*, p. 650 (1877-78).

*C. introflexus* Brid. *Bryol. univ. I*, p. 472 (1826) *fid. Braithw. Brit. Moosfl.* p. 135 (1882).

*C. longipilus* { *Brid. Bryol. univ. I*, p. 477 p.p. (1826); *Bryol. eur. fasc. 41 (Vol. I), Mon.* p. 6, t. 93 (1837).  
var. *de Not. Muscol. ital. Spic.* p. 14 (1837).

*C. pilifer* Brid. *Mant. M.* p. 72 (1819).

*Carpocia polytrichoides* Vent. in *Rev. bryol.* 1881, p. 19.

*Dicranum capitiflorum* P. B. *Prodr.* p. 53 (1805).

- D. flexuosum  $\gamma$  piliferum *Turn. Muscol. hib. p. 74 p.p.* (1804).  
 D. ericetorum *Mitt. M. Ind. or. p. 20* (1859).  
 D. introflexum *Hedw. Sp. M. p. 147, t. 29* (1801) *fid. Lindb.*  
 D. longipilum *C. M. Syn. I, p. 411* (1849).  
 Saxic. et torfac. — *Eur.* : Hibern., Angl., Eur. occ. et merid. (*ubi hucusq. semper sterile?*); Lusit. (c. fr. ! *fid. Venturi in Rev. bryol.* 1881, p. 19). *Afr.* : Alger., Madeir., ins. Azor. *Am. merid.* : Brasil.  
 \* *var. alte-cristatus Ren. et Card. in Bull. Soc. roy. bot. Belg.* 1896, I, p. 303.  
     — *Afr.* : Madagascar.  
 \* *var. Bessoni eor. in op. cit.* 1894, II, p. 110.  
     *C. Bessoni Ren. M. Masc. Mad. n. 58.*  
     Loc. humid. — *Afr.* : Madagascar.  
     *var. Daldinianus de Not. — C. Mildei.*  
 \* *var. elongatus (C. M.) Par. Ind. p. 257* (1894).  
     *Dicranum longipilum var. elongatum C. M. op. et l. cc.*  
     Ut typ. — *Eur.* : Gall. centr. (Andegav.).  
     *var. Mariana Amann. — C. Mildei.*  
 \* *var. vaporarius (Bolle) Jaeg. Ad. I, p. 145* (1870-71).  
     *C. vaporarius Bolle Erb. critt. n. 1210.*  
     Ut typ. — *Eur.* : Ital. (ins. Inarines).  
*porphyreocaulis (C. M.) Par. Ind. p. 258* (1894).  
     *Dicranum porphyreocaulon C. M. in Linn. 1878-79, p. 472.*  
     — *Am. merid.* : Venezuela.  
*porphyreodictyon (C. M.) Mitt. M. austr. am. p. 75* (1869).  
     *C. microstomus Mitt. in Spr. Cat.*  
     *C. tenuissimus Sull. in Proceed. of the Amer. Acad.* 1861.  
     *Dicranum arctocarpum C. M. in Linn. 1847, p. 205.*  
     *D. porphyreodictyon ej. Syn. I, p. 395* (1849).  
     *Thysanomitrium arctocarpum  $\beta$  Moritzianum Hpe. in Linn. 1847, p. 75.*  
     Lignic., saxic. et terr. — *Am. sept.* : Cuba, Jamaic. *Am. merid.* : Ins. Trinitat., Nov. Granat., Brasil. tropic., Aud. Peruv.  
*praealtus (C. M.) Par. Suppl. Ind. p. 96* (1900).  
     *Dicranum praealtum C. M. in Hedwig. 1898, p. 227.*  
     Ster. In fiss. rup. — *Am. sept.* : M. Porto-Rico.  
*praemorsus (C. M.) Jaeg. Ad. I, p. 140* (1870-71).  
     *Dicranum praemorsum C. M. in Bot. Zeit. 1862, p. 337.*  
     Ster. — *Pac.* : Ins. Hawai.  
*procerus (C. M.) Par. Ind. p. 258* (1894).  
     *Dicranum procerum C. M. in Fl. 1890, p. 472.*  
     Ster. — *Afr.* : Kilima N'Djaro.  
*procerus Broth. — C. Brotherianus.*  
*proliferus (C. M.) Mitt. M. austr. am. p. 85* (1869).  
     *Dicranum proliferum C. M. Syn. II, p. 602* (1851).  
     Ster. — *Am. merid.* : Venezuela.

*propinquus* (Hpe.) Mitt. *op. cit.* p. 75.

*Dicranum propinquum* Hpe. in *Ann. sc. nat.* 1866, p. 367.

— *Am. merid.* And. Nov. Granat.

*pseudo-bicolor* C. M. in *Bull. Soc. roy. bot. Belg.* 1895, I, p. 59.

Ster. — *Afr.* : Madagascar.

*pseudo-brachymitrium* (C. M.) Par.

*Dicranum pseudo-brachymitrium* C. M. in *Hedwig.* 1900, p. 249.

Cortic. et terr. — *Am. merid.* : Brasil. or. (Minas Geraës).

*pseudo-filifolius* (C. M.) Par. *Ind.* p. 258 (1894).

*Dicranum pseudo-filifolium* C. M. in *Linn.* 1878-79, p. 471.

— *Am. merid.* : Venezuela.

*pseudo-julaceus* Par. *op. et l. cc.*

*C. julaceus* Rehm. *M. Afr. austr.* n. 58.

Ster, — *Afr.* : Orange.

*pseudo-nanus* Jaeg. — *Campylopodium euphorocladum.*

*pterotoneuron* (C. M.) Jaeg. *Ad. I*, p. 121 (1870-71).

*Dicranum pterotoneuron* C. M. in *Linn.* 1869-70, p. 34.

Ster. — *As.* : Ceylan.

*pudicus* (Hsch.) Jaeg. *op. cit.* p. 130.

*Dicranum pudicum* Hsch. in *Sieb. M. Nov. Holl.* n. 24 p.p.; *C. M. Syn. I*, p. 407.

♀ et fr. ign. — *Pac.* : Austral. or., Tasman.

*pulvinatus* Rehm. *M. Afr. austr.* n. 62.

*Dicranum pulvinatum* C. M. in *Hedwig.* 1899, p. 80.

Ster. — *Afr.* : Pr. B. Sp. (Stinkwater).

*pulvinatus* Brid. — *Grimmia.*

var. *tennis* Wahlenb. — *Grimmia Mühlenbeckii.*

*purpurascens* Ltz. *Moosst.* p. 118 (1864).

— *Afr.* : Pr. B. Sp.

*purpureo-aureus* (C. M.) Par. *Suppl. Ind.* p. 96 (1900).

*C. hygrometricus* Rehm. *M. Afr. austr.* n. 59

*Dicranum purpureo-aureum* C. M. in *Hedwig.* 1899, p. 82.

Ster. — *Afr.* : Orange, Transvaal.

*purpureo-flavescens* Hpe. *hb.*

*Dicranum purpureo-flavescens* C. M. in *Fl.* 1896, p. 442.

♂ et fr. ign. — *Pac.* : Ins. Hawaï.

*pusillus* W. P. Sch. in *Besch. Prodr. bryol. mexic.* p. 21 (1871).

— *Am. sept.* : Mexic.

*pusillus* W. P. Sch. (Pr. B. Sp.). — *C. subpusillus.*

*Quintasii* Broth. in *Boll. da Soc. Berter.* 1890.

Ster. — *Afr.* : Ins. Sao Thoma.

*Rabenii* Ltz. *Moosst.* p. 158 (1864).

— *Am. merid.* : Brasil. austr. or.

*Rauei* Aust. — *Dicranum Mühlenbeckii.*

*reconditus* Jaeg. — *Sphaerothecium*.

*rectisetus* (Hpe.) Jaeg. *Ad. II*, p. 649 (1877-78).

*Dicranum rectisetum* Hpe. *Symb.* 1874, p. 487.

Cortic. et terr. in silv. — *Am. merid.* : Brasil. austr. or.

*recurvipilus* (C. M.) Par.

*Dicranum recurvipilum* C. M. in *Hedwig.* 1900, p. 259.

Ster. Inter *Lichenes*. — *Am. merid.* : Brasil. (Nov. Friburg.), Chile.

*recurvus* (Mitt.) Jaeg. *Ad. I*, p. 122 (1870-71).

*Dicranum recurvum* Mitt. *M. Ind. or.* p. 20 (1859).

Terr. — *As.* : M. Neilgherr., Ceylan.

*reduncus* (Hsch. et Reinw.) *Bryol. jav.* I, p. 75, t. 62 (1855-61).

*Dicranum Auberti* Bel. *Voy. Ind.* p. 104 (1835).

D. concolor Nees et Bl. in *Nor. Act. Leop.* XI, I, p. 133, t. 16.

D. reduncum H. et R. *ibid.* XIV, II, *Suppl.* p. 703 ; C. M. *Syn. I*, p. 396 (1849 et II, p. 599 (1851) ; Mitt. *M. Ind. or.* p. 18 (1859).

D. sinuosum Brid. *Bryol. univ.* I. p. 427 p.p. (1826).

Terr. — *As.* : Ceylan, Java.

*Restingae* Broth. in *Ule Bryoth. brasil.* n. 212.

In solo turfoso. — *Am. merid.* : Rio de Janeiro.

*retinervis* (C. M.) Par. *Suppl. Ind.* p. 93 (1900).

*Dicranum retinerve* C. M. in *Bull. hb. Boissier*, 1897, p. 552.

— *Am. sept.* : M. Jamaic.

*rigens* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1896, I, p. 300.

Lignic. — *Afr.* : Madagascar.

*rigidiusculus* (Hpe.) Jaeg. *Ad. I*, p. 136 (1870-71).

*Dicranum rigidiusculum* Hpe. *Symb.* 1870, p. 272.

— *Am. merid.* : Brasil. austr. or.

*rigidus* (Hsch.) Jaeg. *op. cit.* p. 147.

*Dicranum rigidum* C. M. *Syn. I*, p. 409 (1849).

*Thysanomitrium rigidum* Hsch. *Fl. Brasil.* I, p. 15 (1810).

— *Am. merid.* : Brasil. austr. or.

*ripicolus* Besch. *Fl. bryol. Réunion, etc.* p. 38 (1880-81).

Ster.

— *Afr.* : Ins. Bourbon.

*Roellii* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1899, II, p. 17.

Ster.

— *Am. sept.* : Costa Rica.

*rosulatus* (Hpe.) Mitt. *M. austr. am.* p. 79 (1863).

*Dicranum rosulatum* Hpe. in *Linn.* 1863, p. 139.

Terr. — *Am. merid.* : And. Nov. Granat.

*rufescens* Broth. in *sched.*

C. tortilisetus Ule *Coll. M. brasil.* n. 1348.

*Dicranum rufescens* C. M. in *Bull. hb. Boissier*, 1893, p. 31 et in *Coll. cit.* n. 1792, 1793.

Cortic. et terr. — *Am. merid.* : Brasil. austr. or.

• *var. lutescens* (C. M.) Par. *Suppl. Ind.* p. 96 (1900).

*Dicranum rufescens var. lutescens* C. M. in *op. et Coll. cc.* n. 1802.

In decliv. — *Am. merid.* : Brasil. or. (S. do Itatiaia).



« *var. nana* (C. M.) *Par. op. et l. cc.*

*Dicranum rufescens var. nana* C. M. *in op. et Coll. cit. n.* 1794.

Ut et ubi praeced.

*rugosus* *Besch. Fl. bryol. Nouv. Caled. p.* 198 (1873).

Cortic. in silvis. — *Pac.* : Nov. Caledon.

*Robillardi* *Besch. Fl. bryol. Réunion, etc. p.* 34 (1880-81).

Ster.

— *Afr.* : Ins. Franciae, Nossi Be.

*Saddleanus* *Besch. in Bull. Soc. bot. Fr.* 1885, *p.* LVIII et *in Miss. sc.*

*Cap Horn, V, Bot. p.* 264, *t.* 2, *f.* 6 (1889).

*Dicranum Saddleanum* C. M. *in Fl.* 1885, *p.* 407.

Ster.

— *Am. merid.* : Fuegia.

*Sanctae Helenae* W. P. Sch. — *C. helenicus.*

*Sancti-Caroli* *Besch. in Journ. de bot.* 1891, *p.*

— *Am. merid.* : Chile austr.

*sandwicensis* *Par. Suppl. Ind. p.* 97 (1900).

*Thysanomitrium hawaicum* C. M. *in Fl.* 1896, *p.* 440.

— *Pac.* : Ins. Hawaï.

« *var. robusta* *Par. op. et l. cc.*

*Thysanomitrium hawaicum var. robusta* C. M. *in op. et l. cc.*

Ubi typ.

*Sargii* *Ren. et Card. in Bull. Soc. roy. bot. Belg.* 1899, *II, p.* 216.

— *Am. sept.* : Costa Rica.

*Sauteri* *Kindb. — Dicranum.*

*savannarum* (C. M.) *Mitt. M. austr. am. p.* 85 (1869).

*Dicranum savannarum* C. M. *Syn. II, p.* 596 (1851).

Ster. Terr. — *Am. merid.* : Guian. batav.

*saxicola* *Brid. — Campylosteleum.*

*scabrellus* *Mitt. op. cit. p.* 86.

— *Am. merid.* : Brasil.

*scabricuspis* *Mitt. in Melliss's S. Helena, p.* 358 ( ).

Ster.

— *Afr.* : Ins. S. Helen.

*scabrophyllus* (C. M.) *Par. Suppl. Ind. p.* 97 (1900).

*Dicranum scabrophyllum* C. M. *in Hedwig.* 1897, *p.* 98.

— *Am. merid.* : Monte Video.

*Schenkei* C. M. *in Rehm. M. Afr. austr. n.* 39.

Ster.

— *Afr.* : Pr. B. Sp. (Montagu).

*Schimperi* *Milde in Bot. Zeit.* 1864, *Beil. p.* 13; W. P. Sch. *Syn. II ed. p.* 107 (1876).

*C. compactus* W. P. Sch. *in litt.*

*C. Sauteri* *Mol. in Fl. litterisq. (cf. Bayern's Laubm. p.* 52, 1875).

*C. subulatus var. Schimperi* *Husn. Muscol. gall. p.* 43 (1884).

*Leptotrichum tomentosum* *Kindb. Note on Canad. M.* (1893).

Ad rup. humid. — *Eur.* : Ins. Shetland, Scot., Scandin., Salisb., Stir., Carinth., Alp. Algov., Tirol., Alp., Pyren. *As.* : Caucas. *Am. sept.* : Groenland, Alaska.

Schmidii (C. M.) Jaeg. Ad. I, p. 143 (1870-71).

Dicranum Schmidii C. M. in Bot. Zeit. 1853, p. 37; Mitt. M. Ind. or. p. 21.

— As. : M. Neilgherr.

Schwaegrichenii Duby in Mém.... Genève, 1869, p. 10, t. IV, f. 2.

C. trachyblepharon C. M. ?

— Am. merid. : Brasil. austr. or.

Schwartzii W. P. Sch. Bryol. eur. Suppl. 1-2, t. 1 (1864) et Syn. II ed. p. 105 (1876).

C. auriculatus Wils. mss.

♂ et fr. ign. — Eur. : Norveg. occid., Scot., Hibern., Rhaet., Tirol., Carinth., Stir., Salisb.

sciuroides Welw. et in Duby in Mém.... Genève, 1870, p. 6, t. XI, f. 4.

Ster. Rupic. — Afr. : Angola.

scopelliformis (C. M.) Par.

Dicranum scopelliforme C. M. in Hedwig. 1900, p. 248.

Lignic. — Am. merid. : Brasil. austr. or. (ins. S. Catharina).

scottianum Brid. — Dicranum.

Sellowianus (Hpe.) Jaeg. Ad. II, p. 649 (1877-78).

Dicranum Sellowianum Hpe. Symb. 1872, p. 43.

Thysanomitrium scabrisetum ej. Ic. M. t. 23 B (1844).

— Am. merid. : Brasil. austr. or.

senex (C. M.) Par. Suppl. Ind. p. 97 (1900).

Dicranum senex C. M. in Hedwig. 1897, p. 311.

Ster. — Pac. : Austral. or. (Victoria).

serratus Lac. Sp. nov. M. Archip. ind. p. II, t. 7 B (1872).

— As. : Ins. Banca.

serridorsus (C. M.) Par. op. et l. cc.

Dicranum serridorsum C. M. in Hedwig. 1899, p. 84 et in Rehm. M. Afr. austr. n. 45.

Ster. — Afr. : Pr. B. Sp.

setaceo-rigidus (Hpe.) Jaeg. Ad. II, p. 760 (1877-78).

Dicranum setaceo-rigidum Hpe. Symb. 1877, p. 721.

Ster. — Am. merid. : Brasil. austr. or.

setaceus Card. in Ann. of the Missouri bot. Gard. 1897, p. 54.

— Afr. : Ins. Azor.

— setifolius Wils. Bryol. brit. II ed. p. 89, t. 40 (1855); Bryol. eur. Suppl. fasc. 3-4, t. 6; W. P. Sch. Syn. II ed. p. 108 (1876).

Ster. Ad rup. et in turfis. — Eur. : Hibern., ins. Skye, Scot.

Shawii Wils. in W. P. Sch. Syn. II ed. p. 851.

Fl. et fr. ign. — Eur. : Ins. Hebrid.

« var. hamata W. P. Sch. op. et l. cc.

Ubi typ.

silvaticus Rehm. M. Afr. austr. n. 50.

— Afr. : Austr.

- smaragdinus** (*Brid.*) *Jaeg. Ad. I*, p. 138 (1870-71).  
*Dicranum smaragdinum* *C. M. Syn. I*, p. 401 (1849).  
*Didymodon smaragdinus* *Brid. Bryol. univ. I*, p. 819 (1826).  
 Ster. — *Afr.* : Ins. Ascension.
- sordidus** *Jaeg.* — *Blindia*.
- Spegazzinii** (*C. M.*) *Par. Ind.* p. 261 (1894).  
*Dicranum Spegazzinii* *C. M. in Fl.* 1885, p. 110.  
 Ster. In pratis turf. — *Am. merid.* : Fuegia.
- spiralis** *Dusen M. Am. austr. n.* 617.  
 Terr. — *Am. merid.* : Patagon. occid.
- Sprucei** *Mitt. M. austr. am. p.* 81 (1869).  
 — *Am. merid.* : Amazon.
- spurio-concolor** (*C. M.*) *Par. Suppl. Ind.* p. 97 (1900).  
*Dicranum spurio-concolor* *C. M. in Nuov. Giorn. bot. ital.* 1897, p. 34.  
 — *Am. merid.* : And. Boliv.
- stenocarpus** (*Hpe.*) *Jaeg. Ad. I*, p. 138 (1870-71).  
*C. arctocarpus* *Mitt. op. cit. p.* 87 p.p.  
*Dicranum stenocarpum* *C. M. Syn. II*, p. 600 (1851).  
*D. stenopus* *ej. op. cit. I*, p. 401 (1849) *excl. syn.* *Thysanomitrium scabrisetum*.  
*Thysanomitrium stenocarpum* *Hpe. Ic. M. t.* 23 A (1844).  
 — *Am. merid.* : — Monte Video.
- stenopelma** (*C. M.*) *Rehm. M. Afr. austr. n.* 52.  
*Dicranum stenopelma* *C. M. in Hedwig.* 1899, p. 83.  
 Silv. — *Afr.* : Pr. B. Sp.
- stramineus** (*Mitt.*) *Jaeg. Ad. II*, p. 646 (1877-78).  
*Dicranum stramineum* *Mitt. in Proced. of the Linn. Soc.* 1863, p. 148.  
 — *Afr.* : Cameroon.
- stricticaulis** (*C. M.*) *Par. Suppl. Ind.* p. 97 (1900).  
*Dicranum stricticaule* *C. M. in Bull. hb. Boissier*, 1898, p. 38 *et in Ule Coll. M. brasil. n.* 1795.  
 In decliv. — *Am. merid.* : Brasil. or. (S. do Itatiaia).
- strictifolius** *Broth. in Act. Soc. sc. fenn.* 1891, n. 5.  
 Ster. Cortic. ? — *Am. merid.* : Brasil. or. (Minas Geraës).
- strictisetus** (*C. M.*) *Par.*  
*Dicranum strictisetum* *C. M. in Hedwig.* 1900, p. 253.  
 Terr. — *Am. merid.* : Brasil. or. (Minas Geraës).
- strictus** *W. P. Sch. in Besch. Prodr. bryol. mexic. p.* 23 (1871).  
*An Pilopogon* ? (*fid. Broth.*).  
 Ster. — *Am. sept.* : Mexic.
- strictus** *C. M.* — *C. erectus*.
- subarctocarpus** (*Hpe.*) *Jaeg. Ad. II*, p. 760 (1877-78).  
*Dicranum subarctocarpum* *Hpe. Symb.* 1877, p. 719.  
 — *Am. merid.* : Brasil. austr. or.
- subarenicolus** *C. M. in Bihang till K. Sv. Vet.-Akad. Foerh.* 1895, n. 3.  
 In arenos sicc. — *Am. merid.* : Brasil. or.

- subbartramiaceus* C. M. — *C. leucobasis* var. *bartramiaceus*.  
*subbrachymitius* C. M. in *Ule Bryoth. brasil. n. 50 et 52*.  
 — *Am. merid.* : Brasil. subtropic. or.  
*subbrachyphyllus* Par. Ind. p. 261 (1894).  
*C. brachyphyllus* C. M. in *Dusen M. Camer. n. 658*.  
 — *Afr.* : Cameroon.  
*subchlorophyllosus* C. M. in *Rab. Bryoth. eur. n. 1362*.  
 — *Afr.* : Prom. B. Sp.  
*subcomatus* Ren. et Card. in *Bull. Soc. roy. bot. Belg. 1896, I, p. 302*.  
 Ster. Terr. — *Afr.* : Madagascar.  
*subconcolor* (Hpe.) Mitt. *M. austr. am. p. 86* (1869).  
*Dicranum subconcolor* Hpe. in *Linn. 1863, p. 138*.  
 Terr. — *Am. merid.* : And. Nov. Granat.  
*subcuspidatus* (Hpe.) Jaeg. *Ad. I, p. 145* (1870-71).  
*Dicranum subcuspidatum* Hpe. *Symb. 1870, p. 273*.  
 Ster. — *Am. merid.* : Brasil. austr. or.  
*subfalcatus* (Hsch.) Jaeg. *op. cit. p. 146*.  
*Dicranum subfalcatum* Hsch. *Fl. brasil. I, p. 13* (1840) ; *C. M. Syn. I, p. 415* (1849).  
 Ster. — *Am. merid.* : Brasil.  
*subflexuosus* Par. Ind. p. 262 (1894).  
*C. abyssinicus* W. P. Sch. in *W. Sch. M. abyss.*  
 — *Afr.* : Abyssin.  
*subfragilis* Ren. et Card. in *Bull. Soc. roy. bot. Belg. 1895, I, p. 59*.  
 Ster. — *As.* : Sikkim.  
*subgriseus* (Hpe.) Jaeg. *Ad. II, p. 648* (1877-78).  
*Dicranum subgriseum* Hpe. *Symb. 1872, p. 47*.  
 Ster. — *Am. merid.* : Brasil. austr. or.  
*subincrassatus* (Hpe.) Jaeg. *op. et l. cc.*  
*Dicranum subincrassatum* Hpe. *op. et l. cc.*  
 Ster. Rupic. — *Am. merid.* : Brasil. austr. or.  
*subintroflexus* Kindb. in *Rev. bryol. 1898, p. 91*.  
 — *Afr.* : Ins. Azor. (ins. S. Michel).  
*subleucogaster* (C. M.) Jaeg. *op. cit. p. 645*.  
*Dicranum subleucogaster* C. M. in *Bull. Torr. bot. Club. V, p. 50* (1878).  
 Ster. Ad terr. argill. humid. — *Am. sept.* : Alabama.  
*subluteus* (Mitt.) Jaeg. *Ad. I, p. 121* (1870-71).  
*Dicranum subluteum* Mitt. *M. Ind. or. p. 19* (1859).  
 Ster. — *As.* : Ind. or., penins. Malacc.  
*subnanus* (C. M.) Jaeg. *op. cit. p. 119*.  
*C. flexuosus* Dz. et Mk. *M. frond. ined. Archip. ind. p. 138 ?* (1854).  
*Dicranum nanum* C. M. *Syn. I, p. 384 p.p.* (1819).  
*D. subnanum* ej. in *Bot. Zeit. p. 190*.  
 Terr. — *As.* : Java, ins. Sundaëns., Borneo. Pac. : Ins. Hawaï,  
 Nov. Caledon.

**subpenicillatus** (*C. M.*) *Par.*

*Dicranum subpenicillatum C. M. in Hedwig. 1900, p. 257.*

Ster. — *Am. merid.* : Brasil. or.

**subproliferus** *C. M. in Bull. Soc. roy. bot. Belg. 1892, I, p. 149.*

Silvic. — *Am. sept.* : And. Costa Ricens.

**subpusillus** *Par. Suppl. Ind. p. 98 (1900).*

*C. pusillus W. P. Sch. in Breutel M. capens.*

*Micro-Campylopus pusillus C. M. in Hedwig. 1899, p. 78.*

— *Afr.* : Pr. B. Sp.

**subreconditus** (*Geh. et Hpe.*) *Par. Ind. p. 262 (1894).*

*Dicranum subreconditum G. et H. in Fl. 1881, p. 343.*

— *Am. merid.* : Brasil. austr. or.

**subsecundus** *Lindb. mss.; Aongstr. in Oefv. af K. Vet.-Akad. Foerh. 1879, n. 4, p. 5.*

— *Am. merid.* : Brasil. austr. or.

**subtorquatus** *C. M.* — *C. Woolsii var. cylindrica.***subtricolor** *Ltz. Moosst. p. 159 (1864).*

Ster. — *As.* : India (absq. loc. spec.).

**subulatus** *W. P. Sch.* — *C. brevifolius.*

var. *Schimperi* Husn. — *C. brevifolius var. Schimperi.*

**subulifolius** *Thw. et Mitt. in Journ. of the Linn. Soc. 1872, p. 297.*

— *As.* : Ceylan.

**subvirescens** *Ren. et Card. in Bull. Soc. roy. bot. Belg. 1894, II, p. 109.*

Silvic. — *Afr.* : Madagascar.

**sulfureo-flavus** (*C. M.*) *Par. Suppl. Ind. p. 98 (1900).*

*Dicranum sulfureo-flavum C. M. in Hedwig. 1897, p. 352.*

— *Pac.* : Nov. Seeland. austr.

**sulfureo-nigritus** *Dusen M. Am. austr. n. 627.*

In turfis. — *Am. merid.* : Chile austr.

**sulfureus** *Besch. in Bull. Soc. bot. Fr. 1898, p. 55.*

— *Pac.* : Taïti.

**surinamensis** *C. M. in Linn. 1848, p. 186.*

*Dicranum surinamense ej. Syn. I, p. 386 (1849).*

♀ et fr. ign. Terr. — *Am. merid.* : Guian. batav.

**tallulensis** *Sull. et Lesq. in Sull. Ic. M. p. 27, t. 17 (1864); Lesq. et Jam. Man. p. 78 (1884).*

Ster. Ad rup. irror. — *Am. sept.* : Georgia.

**tasmanicus** *W. P. Sch. hb.*

*C. introflexus ? (fd. Broth.).*

*Dicranum tasmanicum C. M. in Hedwig. p. 351.*

— *Pac.* : Nov. Seeland., Austral. or., Tasman.

**tenax** *Rehm. M. Afr. austr. n. 54.*

*Dicranum tenax C. M. in Hedwig. 1899, p. 83.*

Ster. — *Afr.* : Pr. B. Sp.

**tenuinervis** *Fleisch. M. Archip. ind. n. 107 (1900).*

Terr. — *As.* : Java.

*tenuissimus* Sull. — *C. porphyreodictyon*.

*terebrifolius* (C. M.) Jaeg. *Ad. II*, p. 649 (1877-78).

*Dicranum terebrifolium* C. M. in *Linn.* 1874, p. 593.

Ster. — *Am. merid.* : M. Pichincha.

*tenuis* Wils. — *Campylopodium khasianum*.

*Thwaitesii* (Mitt.) Jaeg. *Ad. I*, p. 121 (1870-71).

*Dicranum Thwaitesii* Mitt. *M. Ind. or.* p. 19 (1859).

Ster. — *As.* : Ceylan, Java.

*Tijucae* Broth. in *Ule Bryoth. brasil.* n. 213.

Rupic. — *Am. merid.* : Rio de Janeiro.

*torfaceus* Bryol. *eur. fasc. 41 (Vol. I), Mon. p. 4, t. 91* (1848); *W. P. Sch. Syn. II ed. p. 193* (1876).

*Bryum flexuosum* L. *Sp. pl.* p. 1593 p.p. (1753).

*C. flexuosus* Brid. *Bryol. univ. I*, p. 469 p.p. (1826).

*C. piriformis* ej. *op. cit.* p. 471.

*Dicranum flexuosum* { *Hedw. Sp. M. t. 38* (1801).  
var. *Schultz. Fl. starg. p. 298* (1806).

*D. pinetorum* Griff. *Not. pl. asiat. P. II*, p. 419 (1849).

*D. piriforme* Schultz. *Suppl. Fl. starg. p. 73* (1819).

*D. turfaceum* C. M. *Syn. I*, p. 399 (1849).

*Thysanomitrium piriforme* Rab. *Deutsch. Kryptfl. II*, 3, p. 149 (1848).

In turfis. — *Eur.* : Fere tota. *As.* : Khasia. *Pac.* : Austral. or. (*fid. Mitt.*)

« *var. alpicola* (C. M.) *Par. Ind. p. 263* (1894).

*C. alpicola* Jaeg. *Ad. I*, p. 96 (1870-71).

*C. torfaceus* W. P. Sch. in *W. Sch. M. abyss. II Coll. n. 12*.

*Cephalogonium densum* W. P. Sch. in *Jaeg. Ad. II*, p. 650 ? (1877-78).

*Dicranum alpicolum* C. M. in *Bot. Zeit.* 1858, p. 161.

— *Afr.* : Summ. alp. Abyssin.

« *var. Mülleri* (Jur.) *Milde Bryol. siles. p. 77* (1869).

*C. Mülleri* Jur. in *Rab. Bryoth. eur. fasc. XII* (1862).

*C. piriformis* β *Mülleri Braithw. Brit. Moosfl. p. 128* (1882).

*C. torfaceus var. fragilis* Jur. *Laubm. Oesterr. Ung. p. 56* (1882).

Ut typ. — *Eur.* : Westphal.

*torfaceus* W. P. Sch. (Abyss.) — *C. torfaceus var. alpicola*.

*var. fragilis* Jur. — *C. torfaceus var. Mülleri*.

*torquatus* Mitt. *Fl. Tasman. p. 173* (1860).

*C. flexuosus* Hpe. in *Linn.* 1855, p. 206 *fid. Mitt.*

*C. pallidus* H. f. et W. *Fl. of the N. Zeal. II*, p. 68, t. 84, f. 3 p.p. (1855)  
et *Handb. p. 414* (1867).

*Dicranum flexuosum* C. M. in *Adn. ad M. Mossman. in Bot. Zeit.* 1851,  
p. 551 (*fid. Mitt.*).

*D. torfaceum* Mitt. in *Hook. Kew Misc.* 1856, p. 257.

*D. torquatum* ej. in *Journ. of the Linn. Soc.* 1859, p. 69.

Ster. — *Pac.* : Austral., Nov. Seeland., Tasman., ins.  
Campbell.

*tortilisetus* Ule. — *C. rufescens*.

*tortuosus* (Hpe.) *Par. Suppl. Ind. p. 98* (1909).

? *Dicranum tortuosum* Hpe. in *Linn.* 1852, p. 361; *Par. Ind. p. 372* (1894).

Ster. In udis inter saxa. — *Am. sept.* : Porto Rico.

- trachyblepharon (*C. M.*) *Mitt. M. austr. am.* p. 80 (1869).  
*C. Schwaegrichenii* *Duby*?  
*Dicranum concolor* *Hsch. Fl. brasil. I, p. II* (1840).  
*D. trachyblepharon* *C. M. Syn. I, p. 389* (1849).  
 Ad terr. arenac. — *Am. merid.* : Brasil. austr. or.
- trachynotus (*C. M.*) *Par. Suppl. Ind. p. 98* (1900).  
*Dicranum trachynotum* *C. M. in Bull. hb. Boissier, 1898, p. 33 et in Ule Coll. M. brasil. n. 1801.*  
 Silvic. — *Am. merid.* : Brasil. or. (S. do Itatiaia).
- trachythecius (*C. M.*) *Jaeg. Ad. I, p. 127* (1870-71).  
*Dicranum trachythecium* *C. M. in Linn. 1869-70, p. 23.*  
 — *As.* : Ceylan.
- transvaaliensis *Rehm. M. Afr. austr. n. 450.*  
 Ster. — *Afr.* : Transvaal.
- trichodes *Ltz. Moosst. p. 159* (1864).  
 Ster. — *Afr.* : Orange, pr. B. Sp.
- tricolor (*C. M.*) *Jaeg. Ad. I, p. 141* (1870-71).  
*Dicranum tricolor* *C. M. in Bot. Zeit. 1853, p. 36; Mitt. M. Ind. or. p. 21* (1859).  
 — *As.* : M. Neilgherr.
- trivialis (*C. M.*) *Par. Suppl. Ind. p. 98* (1900).  
*Dicranum triviale* *C. M. in Nuov. Giorn. bot. ital. 1896, p. 35.*  
 — *Am. merid.* : And. Boliv.
- truncatus *C. M. in Linn. 1844, p. 687.*  
*C. xanthophyllus* *Mont. in Ann. sc. nat. 1845, p. III, in Gay Hist. Chil. Bot. crypt. t. 4, f. 2, et Syll. p. 43* (1856).  
*Dicranum truncatum* *C. M. Syn. I, p. 410* (1849).  
 — *Am. merid.* : Chile austr. *Pac.* : Nov. Seeland.
- Türckheimii (*C. M.*) *Par. Suppl. Ind. p. 98* (1900).  
*Dicranum Türckheimii* *C. M. in Bull. hb. Boissier, 1897, p. 186.*  
 — *Am. sept.* : Guatemala.
- Uleanus (*C. M.*) *Par.*  
*Dicranum Uleanum* *C. M. in Hedwig. 1900, p. 258.*  
 Rupic. — *Am. merid.* : Brasil. austr. or. (ins. S. Catharina).
- uncinatus *Bryol. jav., Kindb. — Dicranodontium.*
- Valentini *Besch. Fl. bryol. Réunion, etc. p. 37* (1880-81).  
 Ster. — *Afr.* : Usambara, ins. Borbon.
- Vallis Gratiae* *Hpe. — C. leucobasis var. longescens.*
- ventri-alaris (*C. M.*) *Par.*  
*Dicranum ventri-alaris* *C. M. in Hedwig. 1900, p. 251.*  
 Cortic., rupic. et terr. — *Am. merid.* : Brasil. or. (Minas Geraës).
- verrucosus *Besch. Fl. bryol. Nouv. Caled. p. 200* (1873).  
 — *Pac.* : Nov. Caledon.
- verticillatus (*Hpe.*) *Rar. Ind. p. 264* (1894).  
*Dicranum verticillatum* *Hpe. Enum. M. brasil. p. 19* (1879).  
 — *Am. merid.* : Monte Video.

- vesticaulis* Mitt. in *Melliss's S. Helena*, p. 359.  
— *Afr.* : Ins. Tristan da Cunha.
- villicaulis* (Hpe.) Jaeg. *Ad. II*, p. 645 (1877-78).  
Dicranum villicaule Hpe. *Symb.* 1872, p. 44.  
Ster. — *Am. merid.* : Brasil. austr. or.
- virescens* Besch. *Fl. bryol. Réunion, etc.* p. 36 (1880-81).  
Ster. — *Afr.* : Ins. Borbon.
- virginicus* Lesq. et Jam. *Man.* p. 80 (1884).  
Dicranum virginicum Aust. in *Coult. Bot. Gaz. IV*, p. 150.  
Ster. — *Am. sept.* : Virgin. occid.
- viridatulus* C. M. in *Dusen M. Camer. n.* 132; *Broth. in Engler's Bot. Jahrb.* 1897, p. 234.  
Rupic. — *Afr.* : Cameroon, Guinea. gall.
- viridatus* (C. M.) Par.  
Dicranum viridatum C. M. in *Hedwig.* 1900, p. 257.  
Rupic. — *Am. merid.* : Brasil. austr. or. (ins. S. Catharina).
- viridicatus* (C. M.) Par. *Suppl. Ind.* p. 99 (1900).  
Dicranum viridicatum C. M. in *Hedwig.* 1897, p. 352.  
Ad rup. madid. — *Pac.* : Austral. or. (Nov. Vales., Queensland).
- viridis* Sull. et Lesq. — Dicranum.
- Vizliputzli* Ltz. *Moosst.* p. 158 (1864).  
*An Pilopogon ? (fid. Broth.).*  
— *Am. sept.* : Mexic.
- Walkeri* (Mitt.) Jaeg. *Ad. I*, p. 133 (1870-71).  
Dicranum Walkeri Mitt. *M. Ind. or.* p. 18 (1859).  
— *As.* : Ceylan.
- Wallisii* Sb. — Dicranodontium purpurascens.
- Wawreanus* Reichdt. *Beitr.....* p. 565.  
Terr. — *Pac.* : Ins. Hawaï.
- Weddelii* Besch. in *Journ. de Bot.* 1891, p.  
Ster. — *Am. merid.* : Peruv.
- weisiopsis* (C. M.) Par. *Suppl. Ind.* p. 99 (1900).  
Dicranum weisiopsis C. M. in *Hedwig.* 1899, p. 79.  
Ster. — *Afr.* : Pr. B. Sp.
- Wheeleri* Hpe. — Dicranoloma.
- Widgrenii* (C. M.) Mitt. *M. austr. am.* p. 88 (1869).  
Dicranum Widgrenii C. M. in *Bot. Zeit.* 1856, p. 418.  
— *Am. merid.* : Brasil.
- Woolsii* (C. M.) Par. *Suppl. Ind.* p. 99 (1900).  
Dicranum Woolsii C. M. in *Hedwig.* 1897, p. 348.  
— *Pac.* : Austral. or. (Nov. Vales., Queensland.).



« *var. cylindrica* (C. M.) *Par. op. et l. cc.*

C. subtorquatus C. M. in sched. ; *Par. Ind.* p. 262 (1894).

Dicranum Woolsii *var. theca perfecte cylindrica* C. M. in *op. cit.* p. 349.

— *Pac.* : *lisdem loc.* ; Victoria.

*xanthophyllus* Mont. — C. truncatus.

Zollingerianus (C. M.) *Bryol. jav.* I, p. 77, t. 64 (1855-61).

Dicranum Zollingerianum C. M. *Syn.* II, p. 599.

Rupic. — *As.* : Ceylan, Java, Sumbawa.

zonatus Mol. — C. flexuosus *var. zonatus.*

zygodonticarpus (C. M.) *Par. Ind.* p. 265 (1894).

Dicranum zygodonticarpum C. M. in *Linn.* 1878-79, p. 471.

— *Am. merid.* : Venezuela.

**CAMPYLOSTELEUM** *Bryol. eur. fasc. 33-36 (Vol. II) (1846).* — 3 spec.

*americanum* Solms. — C. saxicola.

saxicola (Web. et Mohr) *Bryol. eur. l. c., Mon. p. 3, t. 116*; C. M. *Syn.* I, p. 417 (1849) et II, p. 603 (1851); W. P. Sch. *Syn. II ed. p. 133* (1876).

Campylopus saxicola *Brid. Mant. M. p. 72* (1819).

Campylosteleum americanum Solms in *Jaeg. Ad. I, p. 176* (1871-72).

Dicranum saxicola W. et M. *Bot. Taschb. pp. 167 et 466* (1807).

Dryptodon saxicola *Brid. Bryol. univ. I, Suppl. p. 770* (1826).

Glyphomitrium saxicola Mitt. in *Braithw. Brit. Moosfl. II, p. 54, t. 5 D.*

Grimmia geniculata *Schwaegr. Suppl. I, P. I, p. 82, t. 22* (1811); *Brid.*

*Bryol. univ. I, p. 189.*

Gr. saxicola H. et T. *Muscol. brit. II ed., p. 87, t. 13* (1827).

Ptychomitrium geniculatum *Hpe. in Fl.* 1837.

Trichostomum saxicola *Hsch. in Linn. Syst. veg. ed. Sprengel, IV, p. 171.*

Weisia geniculata *Brid. Mant. M. p. 38* (1819).

1. Ad saxa aren. et granit. humid. — *Eur.* : Hibern., Angl. ; media valde sporadice! Gallic., Bohem., Saxon., Austr. super. et inf., Siles., Francon., vall. Rhen., Tirol., Helvet., Jurass., Gall. centr. (Arvern.), Pyren., Ital. sup. (Como); in Scand. deest. *Am. sept.* : Ins. princip. Edouard, Massachus., Nov. Angl., Kentucky.

strictum Solms Laub. *Tent. bryol. geogr. Algarv. p. 42* (1868); W. P. Sch. *Syn. II ed. p. 134* (1876).

Weisia Welwitschii W. P. Sch. *op. cit. p. 52.*

1. In fiss. rup. granit. — *Eur.* : Corsica, Lusitan. (prov. Mouchique Algarb.)

venezuelanum Dz. et Mh. *Prodr. fl. bryol. Surinam. p. 38, t. 5* (1854).

2. — *Am. merid.* : Venezuela.

**Capillimentaria** C. M. in *Hedwig.* 1898, p. 220 (*nom. sect. LEUCOBRYI*).

**Capillidium** C. M. in *Linn.* 1878-79, p. (sect. AEROBRYI).

**CARDOTIA** *Besch. in Rev. bryol.* 1899, p. 4. — 2 spec.

Boiviniana (Besch.) *Card. in op. et l. cc.*

Leucobryum Boivinianum *Besch. Fl. bryol. Réunion, etc. p. 48* (1880-81); *Par. Ind. p. 749* (1896).

Ster. Ad rup. umbr. — *Afr.* : Ins. S. Maria de Madag.

heterodictya (*Besch.*) *Card. op. et l. cc.*

*Leucobryum heterodictyum* *Besch. in Journ. de bot.* 1891, p. ; *Par. op. cit.* 750.

2. — *Afr.* : Madagasc., ins. S. Maria.

**Carpoezia** *Vent. in Rev. bryol.* 1881, p. 20 (sect. *CAMPYLOPODIS*).

**CATAGONIUM** *C. M. in Fl.* 1885, p. 425 (sect. *HYPSI*), *portea gen. propr.*  
Spec. omn. non indic. sunt *ISOPTERYGIA* ejusd. nom.

*Draytoni* *C. M.* — *Acrocladium*.

**Catenularia** *C. M. in op. cit.* p. 411 (sect. *PHILONOTIDIS*).

**CATHARINEA** *Ehrh. in Hannov. Mag.* 1780, p. 933 *et in Bêitr. I*, p. 178 (1787).

Spec. omn. non indicatae sunt *ATRICHA* ejusd. nom.

*Abriaquiae* *C. M.* — *Polytrichadelphus*.

*aequinoctialis* *C. M.* — *Psilopilum*.

*aloides* *Roehl.* — *Pogonatum*.

*alpina* *Roehl.* — *Pogonatum*.

*anomala* *Bryhn.* — *Atrichum Haussknechtii*.

*antarctica* *C. M.* — *Psilopilum*.

*aristata* *Hpe.* — *Polytrichadelphus*.

*Arnoldi* *Hpe.* — *Polytrichadelphus*.

*Australasiae* *Hpe.* — *Polytrichadelphus*.

*callibryon* *Ehrb.* — *Atrichum undulatum*.

var. *abbreviata* *C. M.* — *Atrichum undulatum* var. minor.

*canaliculata* *C. M.* — *Oligotrichum*.

*ciliata* *C. M.* — *Polytrichadelphus*.

*compressa* *C. M.* — *Psilopilum*.

*controversa* *Roehl.* — *Atrichum angustatum*.

*dendroides* *Hpe.* — *Polytrichadelphus*.

*Dixoni* *Braithw.* — *Polytrichum gracile*.

*Ehrhardti* *Voit.* — *Atrichum undulatum*.

*erosa* *Hpe.* — *Oligotrichum*.

*glabrata* *Hook.* — *Psilopilum arcticum*.

*glauc* *Hpe.* — *Polytrichadelphus*.

*grossidens* *C. M.* — *Polytrichadelphus*.

*gymnostomula* *C. M.* — *Psilopilum*.

*hercynica* *Ehrh.* — *Oligotrichum*.

β *sancta* *Brid.* — *Oligotrichum hercynicum*?

*innovans* *C. M.* — *Polytrichadelphus*.

*integrifolia* *C. M.* — *Polytrichadelphus*.

*intermedia* *Bel.* — *Pogonatum Belangeri*.

*javanica* *Hpe.* — *Oligotrichum hercynicum*.

*laevigata* *Brid.* — *Psilopilum arcticum*.

*lagenacea* *C. M.* — *Polytrichadelphus*.

*lateralis* *Valz.* — *Atrichum Haussknechtii*.

*magellanica* { *Brid.* — *Polytrichadelphus*.

{ *C. M.* (*Syn. II*, p. 560). — *Polytrichadelphus semi-angulatus*.

*microdendron* *C. M.* — *Polytrichadelphus*.

*minuta* *C. M.* — *Oligotrichum*.

*Molinae* *C. M.* — *Oligotrichum*.

*nana* *Roehl.* — *Pogonatum*.

*Nietneri* *C. M.* — *Oligotrichum*.

*ptriformis* *Hpe.* — *Psilopilum*.

*prolifcans* *C. M.* — *Polytrichadelphus*.

*pseudo-Polytrichum* *Raddi.* — *Polytrichadelphus magellanicus*.

*pulverulenta* *Roehl.* — *Pogonatum urnigerum*.

- pygmaea* C. M. — Psilopilum.  
*Riedeliana* Hpe. — Oligotrichum.  
*sancta* Brid. — Oligotrichum hercynicum.  
*semi-angulata* Hpe. — Polytrichadelphus.  
*semi-lamellata* Hpe. — Oligotrichum.  
*squamosa* C. M. — Polytrichadelphus.  
*subseriata* Lindb. — Atrichum undulatum var. subseriata.  
*sudetica* Presl. — Oligotrichum hercynicum.  
*tapes* C. M. — Psilopilum.  
*tenuirostris* Brid. — Oligotrichum.  
*tschutschica* C. M. — Psilopilum.  
*Trianae* C. M. — Polytrichadelphus.  
*trichodon* C. M. — Psilopilum.  
*undulata* { var. *abbreviata* Rab. — Atrichum undulatum var. minor.  
var. *brevilamelosa* C. M. — Atrichum polycarpum.  
var. *corsica* de Not. — Atrichum undulatum var. minor.  
var. *minor* Hüb. — Atrichum tenellum.  
*Valenciae* C. M. — Polytrichadelphus.

CATHARINELLA C. M. *Syn. I*, p. 213 (sect. POLYTRICHI = POGONATUM);  
*Kindb. Eur. and N. Am. Bryin.* p. 156 (gen. propr.)

*Dixonii* Kindb. — Polytrichum gracile.

CATHAROMNIUM Hook. f. et Wils. *Fl. of the N. Zeal. II*, p. 119 (1855).  
*ciliatum* eor. — Hypopterygium.

Catillaria C. M. in *Bull. hb. Boissier*, 1897, p. 192 (sect. BARBULÆ, nom.).

CATOSCOPIUM Brid. *Bryol. univ. I*, p. 368 (1826). — 1 spec.

*Martianum* Hüb. — Oreas.

*nigratum* (Hedw.) Brid. *op. et l. sc.*; *Bryol. eur. Vol. IV, Mon. p. 4*,  
t. 313; C. M. *Syn. I*, p. 510 (1851); W. P. Sch. *Syn. II*, p. 498 (1876).

*Bryum nigratum* Dicks. *Pl. crypt. fasc. III*, p. 9 (1793).

*Grimmia nigrita* Smith *Fl. brit. III*, p. 1195.

*Weisia nigrita* Hedw. *Descript. III*, p. 97, t. 39 (1792); *Drumm. M. bor. am. I ed. n. 73*.

2. In ericet. turf. et fiss. rup. — *Eur.*: Sporad. in zon. interm.,  
copiosiss. in Alp. reg. arct. et sept.: Spitzberg., ins. Ursor., Neer-  
land., Angl. As.: Jenissei inf. *Am. sept.*: Greenland., sin. Hud-  
son, Terra nov., lac. Huron, Ontario, Super., M. Rupestr.,  
Columb. brit.

CECALYPHUM P. B. *Prodr. p. 50* (1805).

- calycinum* ej. — Holomitrium.  
*condensatum* ej. — Dicranum spurium.  
*cylindraceum* ej. — Holomitrium vaginatum.  
*dichotomum* ej. — Dicranum.  
*fasciatum* ej. — Campylopus longifolius.  
*longirostratum* ej. — Dicranum scoparium.  
*multiflorum* ej. — Cynodontium polycarpum.  
*perichaetiale* ej. — Holomitrium.  
*sciuroides* ej. — Leucodon.  
*scoparium* ej. — Dicranum.  
*scrofulosum* ej. — Dicranum falcatum.  
? *semi-completum* ej. — Conomitrium Hedwigii.  
*spurium* ej. — Dicranum.

*strumiferum* ej. — *Cynodontium polycarpum* var. *strumiferum*.  
*tortile* ej. — *Ptychomitrium polyphyllum*.  
*undulatum* ej. — *Dicranum Schraderi*.

**CEPHALOGONIUM** W. P. Sch. in *Jaeg. Ad. II*, p. 650 (1877-78).

*densum* ej. — *Campylopus torfaceus* var. *alpicola*?

**CEPHALOTRICHUM** Br. et Sch. in W. Sch. M. *Abyss. et Bryol. eur.* vol. IV.

*oligodus* eor. — *Pogonatum*.

*simense* eor. — *Pogonatum*.

**CERATODON** Brid. *Bryol. univ.* p. 480 (1826). — 27 spec.

*amblyocalyx* C. M. — *C. purpureus* var. *amblyocalyx*.

? *antarcticus* Card. in *Rev. bryol.* 1900, p. 43.

*C. kerguelensis* C. M. in *Forschungsreise... Bot.* p. 3 (1889) nom. ?

*Ster.* — *Am. merid.* : Frét. Gerlache.

*brasilensis* Hpe. *Symb.* 1872, p. 39.

2. — *Am. merid.* : Brasil. austr. or.

*brevifolius* Milde. — *C. purpureus* var. *brevifolius*.

*calycinus* Hpe. in *Reichdt. Exped. Novara*.

*C. purpureus* Mitt. *Fl. vit.* p. 385 (1871).

— *Afr.* : Ins. S. Paul.

*capensis* W. P. Sch. in *Breutel M. capens.*

*C. corsicus* (fid. C. M.).

*Pottia ceratodontea* C. M. ?

— *Afr.* : Pr. B. Sp.

*chilensis* (Mont.) Par.

*Cheilothela chilensis* Broth. *Natür. Pflanzenfam.* p. 301.

*Tortula chilensis* Mitt. *M. austr. am.* p. 146 (1869).

*Trichostomum chilense* Mont. in *Ann. sc. nat.* 1845, IV, p. 109 et *Syll.* p. 38 (1856); *C. M. Syn. I*, p. 576 (1849); *Par. Ind. p.* 1320 (1898).

2. Ad terr. arenac. — *Am. merid.* : And. Quitens., Chile austr.

*chloropus* Brid. *Bryol. univ.* I, p. 486 (1826); W. P. Sch. *Syn. II ed.* p. 136 (1876).

*Barbula chloropus* Kindb. *Eur. and N. Am. Bryin.* p. 260 (1897).

*Cheilothela chloropus* Lindb. in *Utkast...* 1878.

*Dicranum chloropus* Brid. *Mant. M.* p. 70 (1819).

*Didymodon ictericus* de Not. *Spicil. n.* 27 (1837).

*Trichostomum strictum* Bruch in de Not. *Syll.* p. 190, n. 252 (1838); *Bryol. eur. II, Mon. p.* , t. 177 ( ) ; *C. M. Syn. I*, p. 576 ; *Rab. Bryoth. eur. n.* 1258.

2. Terr. — *Eur.* : Galloprov., Sardin., Etrur., Ital. merid., Lusit. *Afr.* : Alger.

*Columbiae* Kindb. in *Rev. bryol.* 1896, p. 20.

1. — *Am. sept.* : Columb. brit.

*condensatus* W. P. Sch. in *Breutel M. capens.*

*Diaphanophyllum Vallis Gratiae* Lindb. in *Oefc...* 1863.

*Leptotrichum Vallis Gratiae* Hpe. in sched.

— *Afr.* : Pr. B. Sp.

conicus (*Hpe.*) *Lindb. M. scand. p.* 37 (1879).

*C. purpureus* forma pallida *Boul. in M. Gall. n.* 628 et *M. Fr. p.* 466 (1884).  
*Trichostomum conicum Hpe. in C. M. Syn. I, p.* 575 (1849); *Par. Ind. p.* 1320 (1898).

2. Muric. — *Eur.* : Suec., Fenn., Germ. sept., Anglia, Gall. sept. *As.* :  
 Caucas. *Am. sept.* : Columb. brit., Minnesota.

convolutus *Reichdt. in Novara Exped. Bot. I, p.* 171.

*C. purpureus H. f. et W. Fl. of the N. Zeal. II, p.* 75.

— *Pac.* : Nov. Seeland.

*corralensis* *Jaeg. — Zygodon.*

*corsicus Bryol. eur. II, Mon. p.* , t. 191; *C. M. Syn. II, p.* 633 (1856).

*Barbula corsica Kindb. Eur. and N. Am. Bryin. p.* 260 (1897).

2. Loc. steril. — *Eur.* : Galloprov., Corsica, Sardin., Algarb. *Afr.* :  
 Prom. B. Sp.

*crassinervis Ltz. in Bot. Zeit.* 1866, p. 187.

2. Terr. — *Am. merid.* : Chile austr. *Pac.* : Tasman.

*cylindricus Bryol. eur. — Trichodon.*

*delicatus C. M. in Abhandl. d. Natur. Ver. in Bremen*, 1900, p. 498.

2. — *Pac.* : Ins. Chatham.

« var. minor *ej. in op. et l. cc.*

— *Pac.* : Nov. Seeland.

*dimorphus Philib. in Rev. bryol.* 1888, p. 28.

*C. brevifolius Milde (fid. Kindb.) non ! fid. Limpr.*

2. Muric. — *Eur.* : Alp. Vales., Simplon.

*dubius Dusen M. Am. austr. n.* 384.

Terr. — *Am. merid.* : Chile austr.

*elimatus Broth. in Abhandl. d. Natur. Ver. in Bremen*, 1900, p. 497.

2. — *Pac.* : Nov. Seeland.

*heterophyllus Kindb. in Mac. Cat. p.* 261 (1892).

1. Terr. — *Am. sept.* : Fret. Behring.

*inclinatus Hüb. — Distichium.*

*javanicus Dz. et Mk. — C. purpureus.*

*microcarpus C. M. in Fl.* 1896, p. 449.

*C. purpureus Sull. in Amer. Expl. Exped. Wilkes, p.* (1859).

— *Pac.* : Ins. Hawai.

*minor Aust. in Coult. Bot. Gaz. II, p.* 89 ( ); *Lesq. et Jam. Man. p.* 92 (1884).

2. Ad terr. humid. — *Am. sept.* : Columb. brit., ins. Vancouver.

*Novae Seelandiae (Broth.) Par. Suppl. Ind. p.* 100 (1900).

*Cheilothea Novae Seelandiae Broth. in Oefv. af Finska Vet.-Akad. Foerh.* 1898, p. 164.

2. Ad rup. et terr. sicc. — *Pac.* : Nov. Seeland.

*novo-granatensis Hpe. in Ann. sc. nat.* 1866, p.

— *Am. merid.* : And. Nov. Granat. et Boliv.

*oblungus (Lindb.) Par. — Trichodon.*

purpureus (L.) *Brid. Bryol. univ. I*, p. 480 (incl. var.  $\beta$  --  $\theta$  ?); *Bryol. eur. II, Mon. p. 5*, t. 189 et 190; *C. M. Syn. I*, p. 646; *W. P. Sch. Syn. II ed. p. 132* (1876).

*Barbula helvetica Kindb. in Bull. Soc. bot. ital. 1896*, p. 15.

*B. Saussuriana Brid. Mant. M. p. 94* (1819).

*Bryum basiflorum Gmel. Syst. nat. II*, p. 1332 (1791).

*Br. bipartitum Dicks. Pl. crypt. fasc. II*, p. 7 (1790).

*Br. Celsii L. Sp. pl. II*, p. 1120 (1753).

*Br. papillosum Dicks. Pl. crypt. fasc. IV*, p. 12, t. XI, f. 5 (1801).

*Br. papillosum Dicks. Pl. crypt. fasc. IV*, p. 12, t. XI, f. 5 (1801).  
(surculis et setis geminatis *Dill. Hist. M. p. 385*, t. 49, f. 50 (1741).

*Br. parvum* { *trichodes ramosum, erectis capitulis, subfuscis, in pediculis obscure rubris Dill. Cat. Giss. p. 224* (1718).

*Br. perangustis foliis et cauliculis, foliis crebrioribus et circa summitates magis congestis, capitulis erectis e surculis annotinis egredientibus, pediculis purpureis Dill. Cat. Giss. p. 226* (1718).

*Br. polytrichoides palustre, setis longioribus rubris sericeis Dill. Hist. M. p. 387*, t. 49, f. 51 (1741).

*Br. purpureum Huds. Fl. angl. I ed.*, p. 412 (1762).

*Br. stellare lacustre, foliis rubris capillaceis Dill. Hist. M. p. 382*, t. 48, f. 50 (1741); *id. Limpr.*

*Br. strictum Hoffm. Deutsch. Fl. II*, p. 36 (1796).

*Br. tenue Dicks. Pl. crypt. fasc. III*, p. 8 (1793).

*Br. tenue stellatum, setis purpureis Dill. Hist. M. p. 386*, t. 49, f. 51 (1741).

*Br. trichodes* { *acaulon minimum, setis et capsulis oblongis ej. op. cit. p. 388*, t. 49, f. 54.  
capitulis erectis, pediculis longioribus rectis, flavo-virentibus *ej. Cat. Giss. p. 225* (1718).

*Ceratodon javanicus Dz. et Mk. Pl. Jungh. I*, p. 337.

*Dicranum bipartitum Roth Tent. fl. germ. III*, p. 158 (1800).

*D. Celsii Sw. M. Suec. p. 36* (1799).

*D. intermedium Hedw. Sp. M. p. 138*, t. 31, f. 1-6 (1801).

*D. longisetum Brid. Mant. M. p. 66* (1819).

*D. palustre ej. Musc. rec. II, P. I*, p. 174 (1798).

*D. purpurascens Hedw. Sp. M. p. 137*, t. 35, f. 1-7 (1801).

*D. purpureum ej. Fund. II*, p. 92, t. 4, f. 17 (1782).

*D. strictum Sm. Fl. brit. III*, p. 118 (1805).

*Didymodon papillosum Brid. Sp. M. I*, p. 160 (1806).

*D. purpurascens H. et T. Muscol. brit. p. 65*, t. 20 (1818).

*Fuscina purpurea Schrank Baier. Fl. II*, p. 453 (1749).

*Mnium purpureum L. Sp. pl. I ed., II*, p. 1111 (1753).

*Tortula Saussuriana Brid. Musc. rec. II, P. I*, p. 203 (1798).

*Trichostomum basiflorum Schrank Baier. Fl. p. 449* (1789).

*Tr. Krintjingianum Dz. et Mk. in Zoll. Syst. Verz. pp. 25 et 31.*

*Tr. papillosum Sm. Fl. brit. p. 1238* (1805).

*Tr. purpureum de Not. Syll. p. 189* (1838).

*Tr. tortum Schrank op. et l.c.*

2. In ericat., agger., mur., etc. — Per totum orb. fere terr.; ab Alaska, Groenland. et Spitzberg. ad reg. antarct.

“ *var. amblyocalyx C. M. in Fl. 1885*, p. 417.

*C. amblyocalyx ej. ap. Aongstr. p. 50, in Oefv. af K. Vet.-Akad. Foerh. 1876*, n. 4.

*C. purpureus Aongstr. in op. cit. 1872*, n. 4, p. 6.

Ut typ. — *Am. merid.* : Terr. magell.

“ *subspec. arcticus Kindb. Eur. and N. Am. Bryin. p. 269* (1897).

— *Eur.* : Spitzberg.

“ *var. aristatus Aust. M. Appal. n. 17* (1870); *Lesq. et Jam. Man. p. 92* (1884).

*Arenic.* — *Am. sept.* : New Jersey.

- « *var. brevisetus* Warnst. in *Verh. d. bot. Ver. d. Prov. Brandenb.* 1889, p. 51.  
— *Eur.* : Russ.
- « *var. brevidens* W. P. Sch. in *Besch. Fl. bryol. Réunion, etc.* p. 51 (1880-81).  
— *Afr.* : Ins. Franciae.
- « *var. brevifolius* Milde in *Bot. Zeit.* 1862, p. 460.  
Ad rup. calcar. — *Eur.* : German. merid.
- « *var. brevisetus* Warnst. in *Verh. d. bot. Ver. d. Prov. Brandenb.* 1899, p. 57.  
— *Eur.* : Lusat. infer.
- « *var. cristatus* Corbiere in *Rev. bryol.* 1897, p. 54.  
— *Eur.* : Pyren.
- « *var. flavisetus* Limpr. *Laubm. Deutschl. .... II*, p. 487 (1887).  
— *Eur.* : Transilv., Stir., Tirol.
- var. gemmifera* Jaap. — *Didymodon rigidulus* f. *propagulifera*.
- « *var. Graeffii* (Schlieph.) Limpr. *op. et l. cc.*  
*C. Graeffii* Schl. mss.  
*Didymodon cuspidatus* W. P. Sch. mss. ; *Milde Bryol. siles.* p. 119 (1869).  
— *Eur.* : Alp. Helvet.
- « *var. mammosus* Warnst. in *op. cit.* 1896, p. 48.  
— *Eur.* : Russ.
- « *var. obtusifolius* Limpr. *op. et l. cc.*  
— *Eur.* : Stiria.
- « *var. rotundifolius* Berggr. in *K. Sv. Vet.-Akad. Handl.* 1875, n. 7, p. 44.  
— *Eur.* : Spitzberg.
- « *var. xanthopus* Sull. *M. bor. am. II ed.* p. 29 ( ); *Lesq. et Jam. Man.* p. 92 (1884).  
Rupic. et terr. — *Am. sept.* : Ins. Vancouver, Calif.
- purpureus* { Aongstr. — *C. purpureus* *var. amblyocalyx*.  
H. f. et W. — *C. convolutus*.  
Mitt. — *C. calycinus*.  
Sull. — *C. microcarpus*.  
Wils. — *Trichostomum tophaceum*.  
*var. palustris* Mont. — *C. stenocarpus*.
- sinensis* C. M. in *Nuov. Giorn. bot. ital.* 1896, p. 104.  
2. — *As.* : China centr. (prov. Shen-Si).
- stenocarpus* *Bryol. eur. II, Mon.* p. 4 (*nom.*); *C. M. Syn. I*, p. 647 (1849).  
*Barbula corsica* Spreng. *Syst. Veg. IV, P. II*, p. 323 (1827).  
*C. purpureus* *var. palustris* Mont. *M. Neilgherr. n.* 57.  
*Didymodon stenocarpus* Mitt. *M. Ind. or.* p. 24 (1859).
2. *Rupic. et terr.* — *As.* : Kashmir, Bhotan., M. Neilgherr., Java.  
*Afr.* : Ins. Bourbon., Madagasc. sept. *Am. sept.* : Mexic., And.  
*Costa Ric. Am. merid.* : Columb. Pac. : Tasman.
- stenocarpus* Wils. — *Trichostomum tophaceum*.
- venezuelensis* C. M. in *Linn.* 1878-79, p. 481.  
— *Am. merid.* : Venezuela.

*viridatus* (C. M.) Broth. *Natür. Pflanzenfam.* p. 459.

*Anoetangium viridatum* Par. *Ind.* p. 41, (1894).

*Zygodon viridatus* C. M. in *Fl.* 1890, p. 481.

Ster.

— *Afr.* : Kilima N'Djaro.

*vulcanicus* C. M. in *Bull. hb. Boissier*, 1897, p. 191.

— *Am. sept.* : And. vulcan. Guatemal.

*xanthocarpus* Hsch. — *Trichostomum*.

**Ceratodontium** C. M. in *Forschungsreise...* *Bot.* p. 38 (1889) (sect. MACROMITRII).

**CHÆTEPHORA** Brid. *Mant. M.* p. 148 (1819).

*cristata* Brid. — *Eriopus*.

*incurva* Hsch. — *Hookeria*.

*Langsdorffii* Hook. — *Hookeria*.

*orthotrichoides* Raddi. — *Macromitrium filiforme*.

*pendula* Hook. — *Hookeria*.

*Perrini* Spreng. — *Lepidopilum polytrichoides*.

**CHÆTEPHORA ERIOPUS** Brid. *Bryol. univ. II*, p. 339 (1827).

*cristatus* Brid. — *Eriopus*.

**Chætomitriella** C. M. in *Engler's Bot. Jahrb.* 1883, p. 85 (sect. HYPERI).

**CHÆTOMITRIUM** Dz. et Mk. *M. frond. ined. Archip. ind.* p. 117 (1854).  
(33 spec.)

*acanthocarpum* Bryol. *jav. II*, p. 53, t. 173 (1861-70).

2. — *As.* : Ins. Ceram, Andai.

*borbonicum* Besch. *Fl. bryol. Réunion, etc.* p. 145 (1880-81).

1. Lignic. — *Afr.* : Ins. Bourbon.

*bornense* Mitt. in *Voy. H. S. M. Challenger, III*, p. 213 in *adn.* (1885).

— *As.* : Borneo.

*callichroum* Besch. *Fl. bryol. Nouv. Caled.* p. 228 (1873).

— *Pac.* : Nov. Caledon.

*catractarum* Besch. *Fl. bryol. Réunion, etc.* p. 145 (1880-81).

2? — *Afr.* : Ins. : Nossi Be.

*caudatifolium* C. M. — *Taxithelium*.

*ciliatum* Bryol. *jav. II*, p. 48, t. 168 (1861-70).

2. — *As.* : Java.

*comorense* Hpe. in *Linn.* 1876, p. 253.

Lignic. — *Afr.* : Ins. Anjouan.

*confertum* Thw. et Mitt. in *Journ. of the Linn. Soc.* 1872, p. 317.

— *As.* : Ceylan.

*cygnaeum* C. M. in *Geheeb Neue Beitr. zur Moossfl. v. Neu Guin.*  
p. 6, t. 4 (1889).

— *Pac.* : Nov. Guin. merid.

*Deplanchei* Duby. — *Holoblepharum*.

*depressum* Mitt. *Sam. M.* p. 190 (1867).

Cortic. — *Pac.* : Ins. Samoa.

f See (a...



- Dusenii *C. M.* in *Dus. M. Camer. n.* 421; *Broth. in Engler's Bot. Jahrb.* 1897, p. 259.  
 2? fr. ign. Cortic. — *Afr.* : Cameroon.
- elegans *Geheeb Neue Beitr. zur Moossfl. v. Neu Guinea*, p. 6, t. 5 (1889).  
 2. — *Pac.* : Nov. Guin. merid.
- elongatum *Dz. et Mk. M. frond. ined. Archip. ind.* p. 119 (1854); *Bryol. jav. II*, p. 43 (1861-70).  
*Hookeria elongata Dz. et Mk. in Ann. sc. nat.* 1844, *II*, p. 305; *C. M. Syn. II*, p. 213 (1851).  
 2. Cortic. — *As.* : Ins. Boeroe Molucc., Borneo, Java.
- fimbriatum (*C. M.*) *Bryol. jav. II*, p. 48 (1861-70).  
*Hypnum fimbriatum C. M. Syn. II*, p. 402 (1851).  
*Leskea fimbriata Dz. et Mk. M. frond. ined. Archip. ind.* p. 177, t. 58 (1854).  
 2. — *As.* : Amboine, Borneo. *Pac.* : Nov. Guin.
- frondosum *Mitt. Sam. M. p.* 189 (1867).  
 Cortic. — *Pac.* : Ins. Samoa.
- Geheebii Broth. in Oefv. af Finska Vet.-Akad. Foerh.* 1895, p. 165.  
 2. Ad ramul. arb. — *Pac.* : Austral. or. (Queensland), Nov. Guin.
- Hookeri* Jaeg. — *Hypnum*.
- horridulum *Bryol. jav. II*, p. 40, t. 167 (1861-70).  
 2. — *As.* : Java.
- lanceolatum *Bryol. jav. II*, p. 49, t. 170.  
 — *As.* : Java.
- lancifolium *Mitt. Sam. M. p.* 212 (1867).  
 Ad fol. emort. — *As.* : Molucc. (Arrow).
- leptopoma (*Schwaegr.*) *Bryol. jav. II*, p. 52 (1861-70).  
*Holoblepharum leptopoma Dz. et Mk. M. frond. ined. Archip. ind.* p. 113, t. 36 (1854).  
*Hookeria leptopoma C. M. Syn. II*, pp. 215 et 618 (1851).  
*Hypnum leptopoma Schwaegr. Suppl. IV*, p. 323 (1842).  
*Stereodon venustus Mitt. M. Ind. or.* p. 114 (1859).  
 2. Lignic. — *As.* : Java.
- ♂ var. *Massarti Ren. et Card. in Rev. bryol.* 1896, p. 104.  
 Ubi typ.
- muricatum *Bryol. jav. II*, p. 51, t. 172 (1861-70).  
 2. — *As.* : Java.
- nematosum *Broth. in Bail. Syn. of the Queensl. Fl. IV Suppl.* p. 99 (nom., 1890) et in *Oefv. af Finska Vet.-Akad. Foerh.* 1890, p. 106.  
 2. fr. ign. Cortic. — *Pac.* : Austral. or. (Queensland).
- obscurum *Bryol. jav. II*, p. 50 (1861-70).  
 2. — *As.* : Java.
- orthorhynchum (*Dz. et Mk.*) *Bryol. jav. II*, p. 45 (1861-70).  
*Holoblepharum orthorhynchum Dz. et Mk. M. frond. ined. Archip. ind.* p. 114, t. 37 (1854).  
*Hookeria orthorhyncha eor. in Ann. sc. nat.* 1844, *II*, p. 305; *C. M. Syn. II*, p. 214 (1851).  
 2. Cortic. — *As.* : Celebes, Borneo, Java, Sumatra.

*papillifolium* *Bryol. jav. II*, p. 50, t. 171 (1861-70).

2. — *As.* : Java.

*philippinense* (*Mont.*) *Bryol. jav. II*, p. 44, t. 166 (1861-70).

*Hookeria philippinensis* *Mont. in Lond. Journ. of bot.* 1845, p. II, et *Syll.* p. 15 (1856) ; *C. M. Syn. II*, p. 213 (1851).

2. *Cortic.* — *As.* : *Ins.* Philippin., Ceram, Java.

*pseudo-elongatum* *Broth. in Warburg, Monsonia*, 1899, I, p. 47.

2. — *As.* : *Archip. ind.* (Jolo-archip., Towi Towi).

*rugifolium* *Mitt.* — *Holoblepharum*.

*speciosum* *Mitt.* — *Holoblepharum*.

*taitense* *Sull. in Amer. Expl. Exped.* 1859, p. 22, t. 23 ; *Mitt. Fl. Vit.* p. 292 (1871).

*Holoblepharum taitense* *Sull. in Am. Expl. Exped. Wilkes*, 1859, p. 22, t. 28.

*Hookeria taitensis* *ej. in Proceed. of the Amer. Acad.* 1851, n. 17.

— *Pac.* : *Ins.* Societat., Taïti.

*torquescens* *Bryol. jav. II*, p. 47, t. 169 (1861-70).

2. *Cortic.* — *As.* : *Ins.* Ceram, Java. *Pac.* : Nov. Guin. ?

*volutum* *Mitt. M. Ind. or.* p. 115 (1859).

*Cortic.* — *As.* : Ceylan.

*Vrieseanum* *Bryol. jav. II*, p. 54, t. 174 (1861-70).

2. *Silvic.* — *As.* : *Ins.* Ceram.

*Warburgii* *Broth. in Warburg, Monsonia*, 1899, I, p. 48.

2. — *As.* : Mindanao.

*Wheeleri* *Hpe. in sched.* ; *C. M. in Fl.* 1896, p. 460.

*Ster.* — *Pac.* : *Ins.* Hawai.

**Chasmodocalyx** *Lindb. M. Scand.* p. 30 (sect. *ANDREAEÆ*) (1879).

**CHEILOTHELA** *Lindb. in Ulkast....* 1878, p. 34 (sect. *CERATODONTIS*) ; *Broth.*

*in Oefv. af Finska Vet.-Akad. Foerh.* 1898 (gen. propr.)

*Spec. omn. sunt CERATODONTES ejusd. nom.*

**CHIONOSTOMUM** *C. M. in Linn.* 1870-71, p. 21. — 2 spec.

*glauco-virens* (*Mitt.*) *Jaeg. Ad. II*, p. 338. (1875-76).

*Stereodon glauco-virens* *Mitt. M. Ind. or.* p. 103 (1859).

1. — *As.* : Assam super.

*rostratum* (*Griff.*) *C. M. in op. et l. cc.*

*Neckera rostrata* *Griff. Not.* p. 456 et *l. pl. asiat. II*, t. 87, f. I (1849).

*Stereodon rostratus* *Mitt. op. cit.* p. 102.

1. *Cortic.* — *As.* : Khasia, Ceylan, penins. Malay.

**Chorisodontium** *Mitt. M. austr. am.* p. 64 (sect. *DICRANI*) (1869).

**Chrysoblastia** *C. M. in Linn.* 1878-79, p. 438 (sect. *HYPMI* = *ISOPTERYGIUM*).

**CHRYSOBRYUM** *Lindb.*

*micans* *ej.* — *Hypnum*,

**CHRYSO-HYPNUM** Hpe. *Moosbild.* p. 19 (1871).

Spec. omn. non indic. sunt *MICROTHAMNIA* ejusd. nom.

*camptorhynchum* Hpe. — M. Widgrenii.

**Chylophyllum** Sb. in *Jaeg. Ad. II*, p. 38 (sect. *PTERYGOPHYLLI vel gen. propr.*) (1874-75).

**CINCLIDIUM** Sw. in *Schrad. Bot. Journ.* 1801, p. 25, t. 2. — 6 spec.

arcticum W. P. Sch. *Coroll.* p. 80 (1860) et *Syn. II ed.* p. 494 (1876); C. M. *Syn. I*, p. 154 (1851).

Mnium *Cinclidium arcticum* *Bryol. eur. fasc. 31 (Vol. IV), Mon. Suppl.* p. I, t. 386 (1846).

2. Paludic. — *Eur.* : Spitzberg., ins. Ursorum, Norveg. (Dovre).

• *subsp. polare* Kindb. *Eur. and N. Am. Bryin.* p. 332 (1897).  
Ut typ. — *Eur.* : Spitzberg.

« *subsp. virescens* Kindb. *op. et l. cc.*  
Ut typ. — *Am. sept.* : M. Rupestr.

*Hymenophyllum* Lindb. — Mnium.

latifolium Lindb. in *Hedwig.* 1877, p. 77.

2. — *As.* : Jenissei inf.

Macounii Kindb. in *Rev. bryol.* 1896, p. 21.

*Syn.* — *Am. sept.* : Alberta, Saskatchewan, M. Rupestr.

stygium Sw. in *op. et l. cc.*; *Brid. Bryol. univ. I*, p. 714 (1827); C. M. *Syn. I*, p. 753 (1849); W. P. Sch. *Syn. II ed.* p. 492 (1876).

*Amblyodon stygius* P. B. *Prodr.* p. 41 (1805).

C. subrotundum *Mac. Canad. M. n.* 206.

*Meesea stygia* *Brid. Muscol. rec. II, P. III*, p. 172, n. 403 (1803).

Mnium *Cinclidium stygium* *Bryol. eur. fasc. 5 (Vol. IV), Mon.* p. 17, t. 385 (1838).

*Syn.* Paludic. — *Eur.* : Scand., Angl., German. sept., Belgia, Würtemb., Rhaet., Salisb., Stir., Carinth., Alp. Bavar., Helvet. et Gall. *As.* : Kamschatka (an C. arcticum?). *Am. sept.* : Groenland., Labrador, Anticosti, Nov. Scot.

subrotundum Lindb. in *Not. ur Saellsk. pr. Faun. et Fl. fenn.* 1867, pp. 72-74, W. P. Sch. *op. cit.* p. 494.

*Syn.* Paludic. — *Eur.* : Spitzberg., mar. Alb., Lapon., Norveg. *As.* : Jenissei infer. *Am. sept.* : Groenland., Labrador; Terr. nov. ?

**CINCLIDOTUS** P. B. *Prodr.* p. 28 p.p. (1805), *Mém. de la Soc. de Paris*, 1822. — 5 spec.

acutifolius *Broth. in Act. Soc. sc. Fenn.* 1898, n. 2, p.

2. — *As.* : Kashmir.

aquaticus (Jacq.) *Bryol. eur. fasc. 16 (Vol. III), Mon.* p. 8, t. 276 (1842); W. P. Sch. *Syn. II ed.* p. 237 (1876).

*Anictangium aquaticum* *Hedw. Sp. M.* p. 41 (1801).

A. falcatum P. B. *Prodr.* p. 42 (1805).

*Bryum aquaticum* *Gmel. in L. Syst. nat. XIII ed., 2, P. II*, p. 1331 (1791).

C. falcatus Kindb. in *Rev. bryol.* 1887, p. 43 et in *Malpighia*, 1891, p. 560.

*Fontinalis capillacea* *Scop. Fl. Carn. II ed. n.* 1287 (1769).

*Gumbelia aquatica* C. M. *Syn. II*, p. 652 (1851).

*Gymnostomum aquaticum* *Hoffm. Deutsch. Fl. II*, p. 29 (1796).

*Harrisonia aquatica* Spreng. in *L. Syst. veg. XVI et l., IV, P. I*, p. 145 (1827).

*Hedwigia aquatica* Hedw. *Descr. III*, p. 29, t. II (1792).

*Hypnum aquaticum* Jacq. *Fl. austr. icon. III*, p. 48, t. 298 (1775).

*H. aquaticum nigricans*, foliis angustis falcatis Dill. *Hist. M.* p. 331, t. 43, f. 70 (1741).

*H. nigricans* Vill. *Pl. Dauph. II*, p. 904 (1786).

*Muscus squamosus aquaticus* Mich. *Nov. gen.* p. 115, n. 100 (1729).

*Sekra aquatica* Lindb. *M. Scand.* p. 23 (1879).

2. Ad rup. et sax. calcar. in rivul. — *Eur.* : Med. et merid.; deest in Scandin., Britann., German. sept. *As.* : Syria, Kurdistan. *Afr.* : Alger.

*Brebissoni* Husn. — *Dialytrichia*.

*falcatus* Kindb. — *C. aquaticus*.

fontinaloides (*Hedw.*) *P. B. Prodr.* pp. 28 et 52 (1805); *Brid. Bryol. univ. I*, p. 229 (1826); *Bryol. eur. fasc.* 16 (*Vol. III*), *Mon.* p. 9, t. 277 (1842); *W. P. Sch. Syn. II ed.* p. 236 (1876).

*C. minor* Lindb. in *Oefv. af K. Vet.-Akad. Foerh.* 1864, p. 240 in nota.

*Fontinalis alpina* Dicks. *Pl. crypt. fasc. II*, p. 2, t. 4, f. I (1790).

*F. erecta* Vill. *Pl. Dauph. III*, p. 919 (1786).

*F. minor* L. *Sp. pl. I ed., II*, p. 1107 (1753).

*F. triangularis minor*, carinata e cymis fructifera Dill. *Hist. M.* p. 257, t. 33, f. 2 (1741) et *Hb.*

*Gumbelia fontinaloides* C. M. *Syn. II*, p. 652 (1851).

*Hypnum antipyreticum*  $\beta$  minus Neck. *Meth. M.* p. 152 (1771).

*H. fontinaloides* Hoffm. *Deutsch. Fl. II*, p. 79 (1796).

*Rhacomitrium fontinaloides* Brid. *Mant. M.* p. 80 (1819).

*Sekra minor* Lindb. *M. Scand.* p. 23 (1879).

*Trematodon fontinaloides* Hoffm. *Deutsch. Fl. II ed. III*, p. 65 (1813).

*Trichostomum fontinaloides* Hedw. *Descr. III*, p. 36, t. 14 (1792).

2. Ut praeced. — *Eur.* : Fere tota. *As.* : Tibet occ., Luristan, Caucas. *Afr.* : Alger.

" var. *Lorentzianus* Mol. Bayer. *Laubm.* p. 115 (1875); *Ltz. in Bot. Zeit.* 1869, p. 553.

*Scouleria* ? Gehebbi de Not. *Epil.* p. 755 (1869).

Ut typ. — *Eur.* : Os Elbae, Rhoengeb., Bade, Helvet.

*minor* Lindb. — *C. fontinaloides*.

*pachyloma* E. S. Salmon in *Rev. bryol.* 1900, p. 59, f. 1-10.

*Ster.* — *As.* : Coele Syria.

*riparius* (Host) W. Arn. in *Mém. Soc. hist. nat. Paris, II*, p. 271 (1825); *Bryol. eur. l. c. Mon.* p. 10, t. 278; *W. P. Sch. op. cit.* p. 235.

*Gumbelia riparia* C. M. *Syn. II*, p. 651 (1851).

*Gymnostomum riparium* Host *Syn. pl. Austr.* p. 563 (1797).

*Rhacomitrium riparium* Brid. *Mant. M.* p. 80 (1819).

*Trichostomum aquaticum* Schleich. *Cent. pl. Helvet. n.* 20.

*Tr. nigricans* Brid. *Sp. M. I*, p. 242 (1806).

*Tr. riparium* Web. et M. *Bot. Taschb.* p. 120 (1807).

2. Ad lign. et sax. in rivul. reg. campestr. et subalp. — *Eur.* : Vall. Rhen., Gallic., Austr. inf., Salisb., Stir., Bavar., Tirol., Istria, Helvet., Gall., Hispan. sept. (Astur.).

var. *terrestris* Bryol. eur. — *Dialytrichia Brebissoni*.

CIRRIPHYLLUM Grout in *Bull. Torr. bot. Club.* 1898, p. 222.

Spec. omn. non indic. sunt EURHYNCHIA ejusd. nom.

*Brandegei* Gr. — *Rhynchostegium*.

*cirrosum* var. *coloradense* Gr. — *E. Boscii* var. *coloradense*.

**CLADASTOMUM** C. M. in Bull. hb. Boissier, 1898, p. 21. — 2 spec.

subulatum C. M. in Hedwig. 1900, p. 235.

2. Terr. — *Am. merid.* : Brasil. or. (S. do Itatiaia).

Ulei C. M. in Bull. hb. Boissier, l. c. et in Ule Bryoth. brasil. n. 201.

Terr. — *Am. merid.* : Brasil. or. (S. do Itatiaia).

**CLADODIUM** Brid. Bryol. univ. I, p. 620 (1826).

Sp. omn. non indic. sunt BRYA ejusd. nom.

Zierii Hpe. — Plagiobryum.

**CLADOMNION** Hook. f. et Wils. Fl. of the N. Zeal. II, p. 96 (1855). — 9 spec.

crenato-obtusum Dusen M. Am. austr. n. A. G.

Saxic. — *Am. merid.* : Patagon. occid.

ericoides (Hook.) H. f. et W. op. cit. p. 99 et Handb. p. 458 (1867).

Hypnum ericoides C. M. Syn. II, p. 458 (1851).

Leskea ericoides Hook. M. ex. t. 28 (1818-20); Brid. Bryol. univ. II, p. 293 (1827).

1. — *Pac.* : Nov. Seeland.

gracile (Hpe.) Mitt. M. austr. am. p. 535 (1869).

Leskea distans Mont. in Ann. sc. nat. 1845, p. 95 et Syll. p. 18 (1856).

Leucodon gracilis Hpe. Ic. M. t. 18 (1844).

Neckera Poeppigiana C. M. Syn. II, p. 115 (1851).

2. Cortic. — *Am. merid.* : Chile austr.

montevidense C. M. in Hedwig. 1897, p. 108.

Cortic. — *Am. merid.* : Monte Video.

Müllerianum (Hpe.) Jaeg. Ad. II, p. 130 (1874-75).

Glyphthecium Müllerianum Hpe. in Linn. 1859, p. 637.

Filicic. — *Pac.* : Austral. or., Tasman.

octangulum (C. M.) Bryol. jav. II, pp. 105 et 228, t. 216 (1861-70).

Leskea sciuroides Dz. et Mh. M. frond. ined. Archip. ind. p. 178 (1854).

Neckera octangula C. M. Syn. II, p. 668 (1851).

2. Lignic. — *As.* : Java.

pallens Lac. Sp. nov. Arch. ind. p. 12 (1872).

2. — *As.* : Java.

sciuroides (Hook.) H. f. et W. op. cit. p. 100.

Garovaglia sciuroides Mitt. in Trans. and Proceed. of the roy. Soc. of Victoria. 1883, p. 80.

Glyphthecium sciuroides Hpe. in Linn. 1859, p. 637.

Leskea sciuroides Hook. M. ex. t. 175 (1818-20); Brid. Bryol. univ. II, p. 286 (1827).

Neckera glyphtheca C. M. Syn. II, p. 669 (1851).

Stereodon sciuroides Mitt. in Journ. of the Linn. Soc. 1859, p. 89.

2. Cortic. — *Pac.* : Nov. Guin., Austral. or., Nov. Seeland., Tasman.

setosum H. f. et W. — Cyrtopus.

Valviviae C. M. in Hedwig. 1898, p. 108.

— *Am. merid.* : Chile austr.

**CLADOPANTHUS** Dz. et Mk. M. frond. ined. Archip. ind. p. 80, t. 28 (1854); Bryol. jav. I, p. 18, t. 13 (1855-61). — 1 spec.

pilifer Dz. et Mk. op. et l. cc.

Leucobryum Cladopanthus C. M. Syn. I, p. 78 (1849).

L. piliferum Jaeg. Ad. I, p. 163 (1871-72).

2. Ad arbor. — As. : Java.

**CLAOPODIUM** Lesq. et Jam. Man. p. 327 (subgen. *HYPSI*) = *THUIDIUM* (1884); Ren. et Card. M. Afr. sept. p. 50 ( ), gen. propr.

*laxifolium* Card. — *Brachythecium reflexum*.

**CLASMATODON** Hook. f. et Wils. in Journ. of bot. IV, p. 421, t. 25 (1842). 2 spec.

*Bertrami* Lindb. — Cl. parvulus.

parvulus (*Hpe.*) Sull. in A. Gray, Man. bot. U. St. II ed. p. 660, n. 1, t. 5 (1856) et Ic. M. p. 126, t. 79 (1864); Lesq. et Jam. Man. p. 297 (1884).

Anisodon acutirostris Bryol. eur. fasc. 59-61 (Vol. V), Mon. p. 4 (1852).

A. Bertrami W. P. Sch. Syn. I ed. p. 506, t. 5 (1860) et II ed. p. 588 (1876).

A. perpusillus Bryol. eur. l. c. t. 452.

Cl. Bertrami Lindb. in Journ. of the Linn. Soc. 1871, p. 71.

Cl. parvulus  $\beta$  julaceus C. M. in Linn. 1875, p. 437.

Cl. pusillus H. f. et W. in Drumm. M. bor. am. I ed. n. 40 (1828); Wils. in Hook. Journ. bot. IV, p. 421, t. 25 A (1842).

Hypnum parvulum C. M. Syn. II, p. 409 (1851).

Leskea parvula Hpe. in Linn. 1839, p. 46.

Neckera perpusilla C. M. Syn. II, p. 666, n. 79 b, excl. syn. (1851) et

Deutsch. M. p. 383 c. ic. excl. syn. (1853).

? Pterogonium marginatum Schwein. mss. in Bryol. eur. l. c.

Rhegmatoxon parvulus Hpe. Ic. M. II, n. 14, t. 14 excl. f. 3 (1814).

1. Ad Abietes. — Eur. : Saxon.

var. *julaceus* C. M. — Cl. parvulus.

" var. *rupestris* Sull. et Lesq. in Lesq. et Jam. op. cit. p. 298 et M. bor. am. II ed. n. 246 b.

Cl. rupestris Kindb. in Rev. bryol. 1895.

Ad rup. irror. — Am. sept. : Tennessee, reg. merid.

*parvulus* Sull. (M. Wright.) — *Austinia tenuinervis*.

*pellucidus* Hpe. Symb. 1874, p. 152

1. — Am. merid. : Brasil. austr. or.

*pusillus* H. f. et W. — Cl. parvulus.

*rupestris* Kindb. — Cl. parvulus var. *rupestris*.

*trichelyma* Sull. — *Austinia tenuinervis*.

**CLASTOBRYUM** Dz. et Mk. M. frond. ined. Archip. ind. p. 43 (1854). 1 spec.

*indicum* Dz. et Mk. op. et l. cc. t. 16; Bryol. jav. II, p. 129 (1861-70).

*Astrodonium indicum* Dz. et Mk. in Ann. sc. nat. 1844, II, p. 299.

*Neckera Clastobryum* C. M. Syn. II, p. 91 (1851).

1. Cortic. et terr. — As. : Borneo, Java.

**CLEISTOSTOMA** Brid. Bryol. univ. I, p. 153 (1826). — 1 spec.

Spec. omn. non indic. sunt SYKROPODONTES ejusd. nom.

ambiguum Brid. op. cit. p. 154; Mill. M. Ind. or. p. 125 (1859).

Neckera ambigua ' '. M. Syn. II, p. 107 (1851).

Pterogonium ambiguum Hook. in Trans. of the Linn. Soc. IX, p. 310, t. 26, f. 4.

2? Cortic. — As. : Bhotan, Sikkim, Nepal.

*elegans* Br. — Dendropogon rufescens.

**CLIDOSTOMA** v. Martius (fide W. P. Sch.).

*rufulum* ej. — Dendropogon rufescens.

**Climacio-pterobryum** C. M. in Bull. Soc. bot. Fr. 1878, p. 65 (sect. PTEROBRYELLÆ).

**CLIMACODONTIUM** Hpe. in Jaeg. Ad. II, p. 713 (1877-78).

*cygnicollum* Hpe. — Anomobryum.

**CLIMACIUM** Web. et Mohr, Reise durch Schweden, p. 96 (1804). — 7 spec.

*americanum* Brid. Bryol. univ. II, p. 273 (1827); Sull. Ic. M. p. 151, t. 97 (1864); Lesq. et Jam. Man. p. 314 (1884).

Neckera dendroides  $\beta$  americana C. M. Syn. II, p. 122.

2. Lignic. — Am. sept. : Terr. nov., Canada merid. or., M. Rupestr., reg. sept., centr., or. et merid.

« var. *Kindbergii* Ren. et Card. in Bot. Gaz. 1890, p. 59.

— Am. sept. : Indiana, Massachus., Ludovic.

*dendroides* (Dill., L.) W. et M. op. et l. cc., et Bot. Taschb. p. 252, t. 10, f. 2 (1807); Brid. Bryol. univ. II, p. 271 (1827); Bryol. eur. V, t. 437; W. P. Sch. Syn. II ed. p. 627 (1876).

Cl. europaeum P. B. Mém. Soc. Linn. Paris, I, t. 9, f. 8.

Hypnum dendroides L. Sp. pl. ed. p. 1128 (1753); et Hb.

Leskia dendroides Hedw. Sp. M. p. 228 (1801).

Neckera dendroides Timm. Prodr. fl. Megap. p. 221, n. 800 (1788); C. M. Syn. II, p. 121 (1851).

2. In prat. udis, ad marg. foss. et ped. mur. — Eur. : E reg. campestr. usq. in montana. As. : Japon., China, Jenissei, Obi, Caucas. occ. et centr. Am. sept. : Groenland., Terra nov., ins. Miquelon, Canada merid. or.; Alaska, Columb. brit., Washington, Kansas. Calif.; New York, M. Alb., Wisconsin, Minnesota, Idaho.

« var. *fluitans* Hüb. Muscol. germ. p. 595 (1833).

Cl. dendroides  $\gamma$   $\beta$  *inundatum* Mol. in Fl. 1881, n. 19.

† f. *limosa* vel *inundata* ej. in Ltz. Moost. p. 105 (1864).

Ad rivul. silvat. — Eur. : Norveg. bor., Vall. Rhen., Pinzgov., Bavar., etc.

var. *inundatum* Mol. — Cl. dendroides var. *fluitans*.

« var. *oregonense* Ren. et Card. in Bot. Gaz. 1890, p. 59.

— Am. sept. : Oregon.

*europaeum* P. B. — Cl. dendroides.

*japonicum* Lindb. in Act. Soc. sc. Fenn. 1872, p. 2.

Hypnum dendroides Thunb. Fl. Jap. p. 363 ? (1784).

Pterobryum imbricatum Duby in Mém... Genève, 1876, p. 6, t. I, f. I.

2. — As. : Tibet. or., China centr. (prov. Schen-Si), Japon.

*lutescens* Voit. — Camptothecium.

*neckeroides* Brid. — Porotrichum dendroides.

*Novae Seelandiae* C. M. in Trans. and Proceed. of the N. Zeal. Inst. 1892, p. 292.

— Pac. : Nov. Seeland.

*ramulosum* (Mitt.) Jaeg. Ad. II, p. 364 (1875-76).

Neckera ramulosa Mitt. in Journ. of the Linn. Soc. 1863, p. 160.

— Afr. : Cameroon.

*ruthenicum* (Weinm.) Lindb. in Act. Soc. sc. Fenn. 1872, p. 248.

Hypnodendron Savatieri W. P. Sch. in Savat. M. Jap. n. 828.

Hypnum ruthenicum Weinm. in Pull. Soc. imp. Moscou, 1845, II, p. 485, n. 34 ; C. M. Syn. II, p. 503 (1851).

H. spectabile H. f. et W. in Wils Bryol. brit. p. 382, n. 58 in obs. (1855).

Pleuroziopsis ruthenica Kindb. Check List. of Eur. and N. Am. M. p. 19. (1894).

2. — As. : Ins. Kuril., Japon., Saghalien. Am. sept. : Fret. Behring, Alaska, Sitka, ins. Vancouver.

*sulcatum* (Hook.) Brid. Bryol. univ. II, p. 274 (1827).

Braithwaitea nematosa Jaeg. Ad. II, p. 365 (1875-76)

Br. sulcata Lindb. in op. cit. p. 250 in adn.

Dendro-Leskea sulcata Hpe. in sched.

Isoetecium sulcatum H. f. et W. Fl. of the N. Zeal. II, p. 104 (1855) et Handb. p. 464 (1867).

Leskea sulcata Hook. M. ex. t. 164 (1818-20).

Neckera sulcata C. M. Syn. II, p. 122 (1851).

Pilotrichum nematosum ej. in Bot. Zeit. 1864, p. 273 et in Linn. 1867-68, p. 614 (cf. ips. in Linn. 1871-73, p. 157).

Pterobryum sulcatum Mitt. in Trans. and Proceed. of the roy. Soc. Victoria, 1883, p. 82.

2 ? — Pac. : Austral. or., Nov. Seeland. centr. et merid.

**Climacocaulon** C. M. in Linn. 1878-78, p. 329 (sect. BARBULÆ).

**CODONOBLEPHARUM** Schwaegr. Suppl. II, p. 142, t. 37 (1823) = ZYGODON ;

Dz. et Mk. M. frond. ined. Archip. p. 95 (1854) = SYRRHOPODON.

Spec. omn. non indicatae sunt ZYGODONTES ejusd. nom.

*crassinerve* C. M. — Syrrhopodon.

*fasciculatum* Bryol. jav. — Syrrhopodon.

*subfasciculatum* Hpe. — Syrrhopodon.

*subluteum* C. M. — Syrrhopodon.

*undulatum* Dz. et Mk. — Syrrhopodon Codonoblepharum.

**CÆLIDIUM** Hook. f. et Wils. Fl. of the N. Zeal. p. 110 (subgen. HYPNI) (1855) ; Reichdt. in Novara Exp. Bot. I, p. 391 (gen. propr.) (1870).

Spec. omn. non indicatae sunt LEMBOPHYLLA ejusd. nom.

*apertum* Jaeg. — Ectropothecium inflectens.

var. *condensatum* ej. — Ectropothecium inflectens var. *condensatum*.

**Cœlophyllum** Kindb. Eur. and N. Am. Bryin. p. 298 (sect. ORTHOTRICH) (1897).



**COLEOCHAETIUM** Besch. Fl. bryol. Réunion, etc. p. 66 (sect. **MACROMITRIUM**) (1880-81); Ren. et Card. in Bull. Soc. roy. bot. Belg. 1894, II, p. 120 (gen. propr.). — 4 spec.

**appendiculatum** Ren. et Card. in op. et l. cc.; Ren. *M. Masc. Mad.* n. 218. Ster. — *Afr.* : Madagascar (Diego-Suarez).

**plicatum** (P. B.) Besch. in Ren. *Prodr. fl. bryol. Madag.* ... etc. p. 143 (1897).

*Leiomitrium plicatum* Mitt. in *Phil. Trans. of the roy. Soc.* 1879 (*extr. vol.*) p. 390.

*Orthotrichum plicatum* P. B. *Prodr.* p. 81 (1805); *C. M. Syn. I*, p. 690 (1849).

*Ulota plicata* Brid. *Bryol. univ. I*, p. 303 (1826).

l. Cortic. — *Afr.* : Madagasc., ins. Borbon, Franciae, Rodriguez.

**rugifolium** (C. M.) Broth. *Natür. Pflanzenfam.* p. 475.

*Macromitrium rugifolium* C. M. in *Hedwig.* 1899, p. 115.

*M. schlotheimiaeforme* Par. *Suppl. Ind.* p. 241 (1900).

*Zygodon rugifolius* C. M. in *Rehm. M. Afr. austr.* n. 151.

Ster. — *Afr.* : Prom. B. Sp.

**secundum** (C. M.) Broth. op. et l. cc.

*Macromitrium secundum* C. M. in *Bot. Zeit.* 1856, p. 420.

Ster. — *Afr.* : Prom. B. Sp.

**Comatosphagnum** C. M. in *Fl.* 1887, p. 404 (*Sphagna subsecunda*).

**Comatulina** C. M. *Syn. II* ed. p. 503 (subgen. **HYPNI** = **MNIODENDRON**).

**Cometium** Thw. et Mitt. in *Journ. of the Linn. Soc.* 1872, p. 302 (sect. **MACROMITRIUM**).

**Complanaria** C. M. in *Linn.* 1878-79, p. 409 (sect. **POROTRICHI**).

**COMPLANATO-HYPNUM** Hpe. *Symb.* 1877, p. 732.

*leptostegium* ej. — *Stereophyllum*.

**CONOMITRIUM** Mont. in *Ann. sc. nat.* 1837, p. 250. — 33 spec.

Spec. omn. non indicatae sunt **FISSIDENTES** ejusd. nom.

**amphibium** C. M. in *Linn.* 1875, p. 356.

*Octodiceras amphibium* Jaeg. *Ad. II*, p. 52 (1874-75).

Ster. — *Afr.* : Niam-Niam.

**assimile** Hpe. — *Fissidens commutatus*.

**atratum** C. M. in *Hedwig.* 1897, p. 89.

*Aquat.* — *Am. merid.* : Argentin. Patagon.

**Beccarii** Hpe. in *Nuov. Giorn. bot. ital.* 1872, p. 290.

*Octodiceras Beccarii* Jaeg. *Ad. II*, p. 52 (1874-75).

l. — *As.* : Borneo.

**bengalense** Hpe. — *F. xiphioides*.

**Berterii** Mont. in *Ann. sc. nat.* 1838, p. 250; *C. M. Syn. II*, p. 524 (1851).

*Fissidens Berterii* C. M. *Syn. II*, p. 45 (1849); *Mitt. M. austr. am.* p. 581 (1869).

*Octodiceras Berterii* Jaeg. *Ad. II*, p. 51 (1874-75).

l. *Ad saxa in scaturig.* — *Am. merid.* : Chile.

- bi-areolatum *C. M. in Linn.* 1878-79, p. 466.  
2. — *Am. merid.* : Venezuela.
- brevicaule (*Broth.*) *Par.*  
*Fissidens brevicaulis Broth. in Bihang till K. Sv. Vet.-Akad. Föerh.*  
1900 p.  
Ster. Submers. — *Am. merid.* : Brasil. austr. (Rio Grande do Sul).
- Bryum *C. M. in Dusen M. Camer. n.* 834.  
— *Afr.* : Cameroon.
- capense *C. M. Syn. II*, p. 524 (1851).  
*Fissidens Berterii ej. Syn. I*, p. 45 p.p. (1819).  
*Octodiceras capense Jaeg. op. et l. cc.*  
Fl. et fr. ign. — *Afr.* : Pr. B. Sp.
- ♂ var. *nigrescens Rehm. M. Afr. austr. n.* 278 et 582.  
— *Afr.* : Orange, Transvaal.
- carnosum (*Broth.*) *Par.*  
*Fissidens carnosus Broth. in Records of the bot. Survey of India*, 1899,  
p. 314; *Par. Suppl. Ind.* p. 157.  
2. Ad rup. granitic. rivul. — *As.* : M. Ghates (Coorg).
- Dillenii (*la Pyl.*) *Mont. et C. M. op. et l. cc.*  
*Fissidens Dillenii C. M. Syn. I*, p. 45 (1849).  
*F. parvus Mitt. M. austr. am.* p. 584 (1869).  
*F. semi-completus Hedw. Descript. III*, p. 34, t. 13 (1792) p.p.  
*Fontinalis parva, foliis lanceolatis Dill. Hist. M.* p. 259, t. 33, f. 45 (1741).  
*Octodiceras Dillenii Brid. Bryol. univ. II*, p. 667 (1827).  
*Skitophyllum Dillenii la Pyl. in Desv. Journ. bot.* 1813, V, p. 52, t. 34.  
— *Am. merid.* : Chile, Patagon., ins. Provident.  
*Pac.* : Austral., Nov. Seeland.
- Duseni *C. M. in Dus. M. Camer. n.* 834.  
— *Afr.* : Cameroon.
- Goebelii C. M. — Fissidens subprosenchymaticus.*
- Hallianum *Sull. et Lesq. in Aust. M. Appal. n.* 108 (1870); *Sull. Ic. M.*  
*Suppl.* p. 43, t. 38 (1874); *Lesq. et Jam. Man.* p. 90 (1884).  
*Octodiceras Hallianum Jaeg. op. cit.* p. 52.  
1. Lignic. in puteis et rupic. — *Am. sept.* : New York, New Jersey  
Illinois, Idaho, Florid.
- Hallianum Mac. — C. Julianum.*
- Hedwigii *Mont. in op. et l. cc.*  
*Cecalyphum ? semi-completum P. B. Prodr.* p. 57 (1805).  
*Fissidens Hedwigii C. M. Syn. I*, p. 45 (1819).  
*F. semi-completus Hedw. op. et l. cc. t. 13 p.p.* ; *Mitt. op. cit.* p. 583.  
*Octodiceras fissidentoides Brid. Bryol. univ. II*, p. 676 (1827).  
*O. Hedwigii Lindb. in Oefv. af K. Vet.-Akad. Föerh.* 1863, p. 405.  
*Skitophyllum semi-completum la Pyl. in op. et l. cc.*  
1. — *Am. merid.* : Chile.
- Hydropogon (*Spr.*) *Par.*  
*Fissidens Hydropogon Spr. in Mitt. M. austr. am.* p. 585 (1869).  
*Octodiceras Hydropogon Jaeg. op. cit.* p. 51.  
1. — *Am. merid.* : And. Quitens.

immersum (Mitt.) *Par. Ind.* p. 281 (1894).

*Fissidens immersus* Mitt. *op. cit.* p. 584.

*Octodiceras immersum* Jaeg. *op. et l. cc.*

— *Am. merid.* : Brasil. tropic.

*Julianum* (Savi) *Mont. in op. cit.* 1837, p. 246, t. 4; *C. M. Syn.* II, p. 524 (1849); *W. P. Sch. Syn.* II ed. p. 122 (1876).

*C. Hallianum* Mac. *Canad. M. n.* 54.

*Fissidens debilis* Schwaegr. *Suppl. 1, P. II, p. II* (1816); *Brid. Bryol. univ.* II, p. 706 (1827).

*F. Julianus* W. P. Sch. in *Fl.* 1838, *P. I, p. 271.*

*F. semi-completus* Drumm. *M. bor. am. II ed., n. 44.*

*Fontinalis Juliana* Savi *Bot. etrusc. III, p. 107* (1818), *et in Pollin. Fl. Veron. III, p. 385* (1824).

*Muscus pinnatus, aquaticus, ramosissimus, Linariae foliis, capitulis...* Mich. *Gen. pl. p. 114* (1728).

*Octodiceras fontanum* Lindb. *Bidrag. Moss. Syn. p. 23* (1863).

*O. Julianus* *Brid. Bryol. univ. II, p. 678* (1827); *Bryol. eur. I, t. 108.*

*Skitophyllum fontanum* laPyl. in *Journ. bot. Desvieux*, 1813, *n. V, p. 153, t. 34, f. 2.*

*Schistophyllum Julianum* Lindb. *M. Scand. p. 13* (1879).

1. Ad saxa in rivul. et ad pariet. lapid. font. — *Eur.* : Temper. et calidior passim : deest in Britann. *Afr.* : Alger; *Pr. B. Sp. ? Am. sept.* : Ontario, M. Rupestr., Calif.; *reg. or., centr. et merid.*

« *var. mexicanum* (*C. M.*) *W. P. Sch. in Besch. Prodr. bryol. mexic. p. 27* (1871).

*C. mexicanum* *C. M. in Bot. Zeit.* 1860.

*Fissidens mexicanus* Mitt. *M. austr. am. p. 584* (1869).

*Octodiceras mexicanum* Jaeg. *op. cit. p. 50.*

Ster.

— *Am. sept.* : Mexic.

*Lorentzia* *C. M. in Hedwig.* 1897, p. 89.

In alveo exsicc. — *Am. merid.* : Argentin. Buenos Ayr.

*Mariei* *Besch. Fl. bryol. Réunion, etc. p. 47* (1880-81).

2. Humic. — *Afr.* : Ins. Nossi Be.

*mexicanum* *C. M.* — *C. Julianum var. mexicanum.*

*molle* *C. M. in Linn.* 1878-79, p. 247.

*Fissidens mollis* Broth. in *Bihang till K. Sv. Vet.-Akad. Foerh.* 1900.

*Fl. et fr. ign.*

— *Am. merid.* : Paraguay, Uruguay.

*Mülleri* *Hpe. in Linn.* 1856, p. 214 et 1859, p. 644.

*Fissidens Dillenii* Mitt. in *Kew Journ. bot. VIII, p. 262* (1858).

*F. Mülleri ej. in Trans. and Proceed. of the roy. Soc. of Victoria*, 1883, p. 91.

*Octodiceras Mülleri* Jaeg. *op. cit. p. 51.*

1. Submers. — *Pac.* : Austral. or. (Nov. Vales., Victoria).

*nigritellum* *C. M. in Hedwig.* 1897, p. 89.

— *Am. merid.* : Argentin. Buenos Ayr.

*palmifolium* (*P. B.*) *Besch. Fl. bryol. Réunion, etc. p. 47* (1880-81).

? *Bryum palmifolium* *Brid. Bryol. univ. I, p. 687* (1826); *C. M. Syn. I, p. 341* (1849).

*C. borbonicum* *C. M. in Bot. Zeit.* 1864.

*Mnium palmifolium* *P. B. Prodr. p. 76* (1805).

*Octodiceras borbonicum* Jaeg. *op. c. p. 50.*

Ster. ? *Aquat.* — *Afr.* : Ins. Borbon.

- perpusillum* C. M. — *Fissidens australiensis*.  
*polycarpum* Besch. in *Journ. de bot.* 1891, p.  
 — *Am. merid.* : Paraguay.  
*platybryoides* C. M. — *Fissidens catharinensis*.  
*Regnellii* (Broth.) Par. *Suppl. Ind.* p. 103 (1900).  
*Fissidens Regnellii* Broth. in *Bihang till K. Sv. Vet.-Akad. Foerh.* 1895,  
 n. 3.  
 1. *Ad saxa irror.* — *Am. merid.* : Brasil. or.  
*scleromitrium* Besch. in *Journ. de bot.* 1891.  
 1. — *Afr.* : Madagascar.  
*semi-limbatum* Schlieph. — *Fissidens Pabstii*.  
 — *Fissidens*.  
*serratum* C. M. } — *Fissidens pseudo-serratus*.  
 — *Fissidens Wilsoni*.  
*sinense* Rab. *Bryoth. eur.* n. 1212.  
 — *As.* : Cochinch.  
*smaragdinum* Ltz. et C. M. in *Linn.* 1878-79, p. 248.  
*Fl. et fr. ign.* — *Am. merid.* : Argentin. Cordob.  
*stissotheca* (Hpe.) C. M. *Syn. II*, p. 625 (1851).  
*Fissidens stissotheca* Hpe. *mss. in C. M. l. c.*  
 1. *Ad saxa irror.* — *Am. merid.* : Brasil. austr.  
*subpalmatum* } Hpe. (Jaeg. p. 762). — *Fissidens evadiinervis*.  
 } Jaeg. p. 41. — *Fissidens*.  
*substissotheca* (Broth.) Par. *Suppl. Ind.* p. 103 (1900).  
*Fissidens substissotheca* Broth. in *Bihang till K. Sv. Vet.-Akad. Foerh.*  
 1895, n. 3.  
 1. *Ad saxa irror.* — *Am. merid.* : Brasil. or.  
*tenerrimum* C. M. in *Nuov. Giorn. bot. ital.* 1896, p. 90.  
 Ster. — *As.* : China centr. (prov. Schen-Si).  
*trachelyma* (Sull.) C. M. in *Linn.* 1878-79, p. 466.  
*Fissidens trachelyma* Sull. *mss.*  
 2. — *Am. merid.* : Venezuela.  
*Türkheimii* C. M. in *Bull. hb. Boissier*, 1897, p. 173.  
 Ster. Aquat. — *Am. sept.* : Guatemala.  
*Ulei* C. M. in *Ule Bryoth. brasil.* n. 121.  
 — *Am. merid.* : Brasil. subtropic. or.

**CONOSTOMUM** Sw. in *Schrad. Neu Journ. f. Bot.* I, P. III, p. 26  
 (1806.). — 9 spec.

- aequinoctiale* W. P. Sch. in *Mandon Fl. Boliv.* n. 1681; C. M. in *Nuov.*  
*Giorn. bot. ital.* 1897, p. 47.  
 2. — *Am. merid.* : And. Boliv.  
 var. *piliferum* W. P. Sch. (n. 1682.) — C. *aequinoctiale*.  
*arcticum* Sw. — C. boreale.  
*australe* So. in *op. et l. cc.*; *Brid. Bryol. univ.* I, p. 152 (1826); C. M.  
*Syn.* I, p. 470 (1849); *Fl. antarct.* II, p. 132 (1847); *Fl. of the N.*

- Zeal. II*, p. 87 (1855); *Fl. Tasman, II*, p. 196 (1860); *Handb.* p. 411 (1867).
- Bartramia australis* Mitt. *M. austr.* p. 287 (1869).
- B. pentasticha* Brid. *Musc. rec. II*, P. III, p. 134, t. I, f. 8 (1803).
- Bryum angulosum* Menz. (fid. *Brid. Bryol. univ. l. c.*).
- C. pentastichum* Lindb. in *Oefv. af K. Vet.-Akad. Foerh.*.... 1863, p. 392 et in *Bidrag till...* p. 8 (1863).
- Philonotis australis* Mitt. in *Journ. of the Linn. Soc.* 1859, p. 81.
- Weisia pentasticha* Hedw. in *Brid. Bryol. univ. l. c.*
1. *Rupic.* — *Afr.* : Ins. Kerguelen (1)? *Am. merid.* : And. Quitens., terr. Magell., Fuegia, ins. Maclov., Eremitae. *Pac.* : Nov. Seeland. sept. et med., *Tasman.*, ins. Auckland et Campbell.
- boreale *Sw.* in *op. cit.* p. 24, t. 5; *Brid. Bryol. univ. I*, p. 150 (1826); *C. M. Syn. I*, p. 469 (1849) et *II*, p. 615 (1851); *W. P. Sch. Syn. II ed.* p. 514 (1876).
- Bartramia conostoma* *Bryol. eur. fasc. 12 (Vol. IV., Mon. p. 16, t. 322 (1842).*
- Bryum tetragonum* (haud Villars 1786) *Dicks. Pl. crypt. fasc. II*, p. 8, t. 4, f. 9 (1790).
- C. arcticum* *Sw.* in *Brid. Sp. M. I*, p. 127 (1806).
- C. tetragonum* Lindb. in *Oefv. af K. Vet.-Akad. Foerh.* 1863, p. 8.
- Grimmia conostoma* Sm. *Fl. brit. III*, p. 1169 (1804).
2. Ad terr. humid. — *Eur.* : Summ. Alp. praecip. reg. sept. : Spitzberg., ins. Ursor., Salisb., Stir., Carinth., Tirol., Helvet. As. : Jenissei inf., penins. Samojed. et Tschutsch., Kamschatka. *Am. sept.* : Praecip. artic. : Groenland., Labrador, ins. Sabine, fret. Kotzebue et Smith, Alaska, ins. Vancouver, Columb. brit., M. Rupestr., Gaspes., M. Alb. et Adirondack, Terr. nov.
- curvirostre (Mitt.) *Par. Ind.* p. 283 (1894.)
- Bartramia curvirostris* Mitt. in *Journ. of the Linn. Soc.* 1859, p. 81.
- *Pac.* : Austral. (Victoria).
- Lorentzi *C. M.* in *Linn.* 1880-82, p. 404.
2. — *Am. merid.* : Alp. Argentin. subtropic.
- magellanicum *Sull.* in *Hook. Journ. of bot.* 1850, p. 316; *C. M. Syn. II*, p. 615 (1851).
- Bartramia magellanica* Mitt. *M. austr. am.* p. 268 (1869).
1. fr. ign. Ad rup. madid. — *Am. merid.* : Fuegia.
- parvulum* Hpe. — *C. pusillum.*
- pentastichum* Lindb. — *C. australe.*
- pusillum* *Hook. f. et Wils. Fl. of the N. Zeal. II*, p. 88, t. 86, f. 2 (1855) et *Handb.* p. 445 (1867).
- C. parvulum* Hpe. in *Linn.* 1856, p. 207.
- Philonotis pusilla* Mitt. in *Journ. of the Linn. Soc.* 1859, p. 81.
- 1 (Mitt.) 2 (Hpe.). Ad rup. madid. — *Pac.* : Alp. Austral. or., Nov. Seeland. sept. et med., *Tasman.*
- rhynchostegium *C. M. Bryol. austr. Georg. p.* 27 (1890).
2. Paludic. — *Am. merid.* : Austr. Georg.
- speirostichum *C. M.* in *Linn.* 1874, p. 594.
- Ster. — *Am. merid.* : M. Pichincha.

1) Cf. *Philonotis anisothecioides*.

*tetragonum* Lindb. — *C. boreale*.  
*variegatum* Brid. — *Weisia zonata*.

**COSCINODON** Spreng. Einleit. in d. Stud. d. Krypt. Gew. p. 281 (1804). — 5 spec.

*aciphyllus* Brid. — *Pottia lanceolata* var. *aciphylla*.  
*comatus* Brid. — *Pottia lanceolata*.

*cribrosus* (Hedw.) Spruce in *Ann. a. Mag. nat. hist.* 2, ser. III, p. 491 (1849); *C. M. Syn.* I, p. 765 (1849).

*Bryum cribrosum* Hoffm. *Deutsch. Fl.* II, p. 31 (1796).

*C. Persoonii* Hpe. in *Fl.* 1837, p. 280.

*C. pulvinatus* Spreng. op. cit. p. p. 281 et 372, t. 8, f. 75 (1804); *Bryol. eur.* III, Mon. p. 3, t. 230; *W. P. Sch. Syn.* II ed. p. 287 (1876; *Par. Ind.* p. 284 (1891)).

*Grimmia cribrosa* Hedw. *Descr.* III, p. 73, l. 31 A (1792).

2. Ad rup. aren. et granit. humid. — *Eur.* : Per totam sporadice, except. *Britann. As.* : *Caucas occ. et centr. Am. sept.* : *Fret. Scoresby, Alaska, Columb. brit, Idaho.*

« *f. propagulifera* Jens. *Mossor fra Ost. Groenland*, p. 422 (1898).

— *Am. sept.* : *Fret. Scoresby.*

« *var. subperforatus* Philib. in *Rev. bryol.* 1880, p. 43.

Ut typ. — *Eur.* : *Gall. centr. (Ardèche).*

*Hartzii* C. Jens. in *Saertryk af Meedd. om Groenland*, 1898, p. 422 c. ic.

— *Am. sept.* : *Fret. Scoresby.*

*Raui* (Aust.) Lesq. et *Jam. Man.* p. 155 (1884).

*Grimmia Raui* Aust. in *Bull. Torr. bot. Club*, VI, p. 46 (1879).

1.

— *Am. sept.* : *Colorado, Minnesota.*

*Renaaldi* Card. in *Bot. Gaz.* 1890, p. 41, t. 6.

*C. Raui* (fid. D<sup>na</sup> E. G. Britt. in *Bull. Torr. bot. Club*, 1895, pp. 447-458 et t. 248-249).

1.

— *Am. sept.* : *Colorado, Kansas.*

*verticillatus* Brid. — *Eucladium*.

*Wrightii* (Aust.) Sull. *M. of U. St.* p. 38, t. 4 (1856) et *Id. M.* p. 71, t. 45 (1864); *Lesq. et Jam. Man.* p. 155 (1884).

*Grimmia Wrightii* Aust. in *Bull. Torr. bot. Club*, VI, p. 46 (1879).

1. *Rupic.* — *Am. sept.* : *Colorado, Kansas, Texas, Nov. Mexic.*

**Grassidicranum** Limpr. *Laubm. Deutschl.*.... I, p. 370 (sect. *DICRANI*) (1886).

**CRATONEURON** Sull. *M. a. Hep. of U. St.* p. 73 (subgen. *HYPNI*) (1856);

*C. M. gen. propr.*

*Spec. omn. sunt HYPNA ejusd. nom.*

**Crenularia** C. M. in *Bull. hb. Boissier*, 1897, p. 173 (sect. *FISSIDENTIS*, nom.).

**Crispidium** C. M. in *Fl.* 1896, p. 485 (sect. *FISSIDENTIS*).

**CROSSIDIUM** Jur. Laubm. v. Oesterr.-Ung. p. 127 (1882). — 6 spec.

chloronotos (*Bruch*) *Limpr. Laubm. Deutschl...* I, p. 645 (1888).

*Barbula chloronotos* Br. in Fr. Müller M. Sardin. (1829) et in Fl. 1829. II, p. 407; *Bryol. eur. fasc.* 13-15 (Vol. II), Mon. p. 18, t. 141 (1842); *C. M. Syn.* I, p. 598 (1849); *W. P. Sch. Syn.* II ed. p. 193 (1876); *Par. Ind.* p. 67 (1894).

*Crossidium crassinerve* Jur. op. cit. p. 128.

*Tortula chloronotos* Broth. Enum. M. Cauc. p. 42 (1892).

*T. crassinervis* de Not. Spec. de Tortul. ital. n. 15 et Syll. p. 176 (1838).

2. *Muric.* — *Eur.*: Rarior: Gall. austr., Sardin., ins. Melit. As.: Reg. transcauc., Afghan. Afr.: Alger., Canar., Madeir. Am. sept.: Colorado. Pac.: Nov. Seeland. sept.?

α *f. niveum* (*Besch.*) *Par.*

*Barbula chloronotos f. nivea* Besch. Cat. M. Aly. p. 12 (1882); *Par. op. et l. cc.*

Ut typ. — Afr.: Alger.

*Geheebii* Broth. Natur. Pflanzenfam. p. 426.

*Tortula Geheebii* ej. in Oefv. af Finska Vet.-Akad. Foerh. 1899, p. 114.

2. — Pac.: Nov. Seeland.

*griseum* Jur. op. et l. cc.

*Barbula grisea* Boul. M. Fr. I, p. 434 (1884); *Par. Ind.* p. 76 (1894).

*B. membranifolia* { *Garov. Bryol. austr.* p. 14 (1840).  
var. *grisea* Vent. in Comm. sc. Faun., Fl. etc. Vene-  
zia, n. 3 (1868).

*Desmatodon griseus* Jur. in Verh. d. K. K. zool. bot. Gesells. in Wien, 1864, p. 399.

*Tortula grisea* Del. Fl. crypt. Belg. M. p. 105 (1883).

*T. squamigera* var. *pottioidea* de Not. Musc. ital. I, p. 22 (1862).

1. Ad rup. praecip. dolomit. — *Eur.*: Helvet occ., Tirol., Istria, Austr. inf. As.: Transcauc., Syria.

*Henrici* (*Rau*) Broth. op. et l. cc.

*Barbula Henrici* Rau in Bull. of the Wasb. Coll. Lab. I, p. 172 (1886); *Par. Ind.* p. 77 (1894).

Ster. Rupic. — Am. sept.: Kansas.

*Manniae* (*C. M.*) Broth. op. et l. cc.

*Barbula Manniae* C. M. in Fl. 1887, p. 222; *Par. op. cit.* p. 82.

— Am. sept.: Colorado.

*squamigerum* (*Viv.*) Jur. op. cit. p. 127.

*Barbula arenaria* Griff. Not. p. 409, et Ic. pl. asiat. II, t. 103, f. I.

*B. chloronotos* Brid. Mant. M. p. 90 (1819) p.p. et *Bryol. univ.* I, p. 539 (1826) excl. syn. *Schultz.*

*B. membranifolia* *Schultz Rec. de Barbula* p. 226, t. 34, f. 35 (1823); *Bryol. eur. fasc.* 13-15 (Vol. II), Mon. p. 17, t. 140 (1842); *C. M. Syn.* I, p. 597 (1849); *W. P. Sch. Syn.* II, p. 192 (1876); *Par. Ind.* p. 83 (1894).

*B. squamigera* Viv. Ann. bot. I, P. II, p. 191 (1804).

*Desmatodon chloronotos* Mitt. M. Ind. or. p. 38 (1859).

*Tortula chloronotos* Brid. Sp. M. I, p. 253 (1806).

*T. membranifolia* Hook. M. ex. I, t. 26 (1818).

*T. squamigera* de Not. Musc. Ital. I, p. 20, t. 5 (1862).

1. Ad mur., rup. et coll. apric. — *Eur.*: Med. occ. et merid., doest in Scand. et Britann.; vall. Rhen., Lutet., Helvet. occ., Tirol., Herzegov., Ital., Galloprov., Hispan. mer., Lusit. As.: Minor, Caucas., Persia. Afr.: Penins. Sinait., Alger. Am. sept.: M. Rupestr., Colorado, Arizona occ., Calif.

« *var. brevisetum* (Besch.) Par.

*Barbula membranifolia var. breviseta* Besch. *Cat. M. Alg.* p. 12 (1882);  
*Par. Ind.* p. 83 (1894).

Ut typ. — *Afr.* : Alger.

**CROSSOMITRIUM** C. M. in Linn. 1874, p. 611. — 22 spec.

*Crügeri* C. M. in *op. et l. cc.*

*Hookeria Patrisiae* ej. *Syn. II*, p. 138 p.p. (1851).

2. *Cortic.* — *Am. sept.* : Antill. *Am. merid.* : Ins. Trinitat.

*epiphyllum* (Mitt.) C. M. in Linn. 1874, p. 613.

*Lepidopilum epiphyllum* Mitt. *M. austr. am.* p. 370 (1869).

2. *Ad fol. emort.* — *Am. merid.* : And. Quitens.

*fontanum* (Mitt.) Jaeg. *Ad. II*, p. 242 (1874-75).

*Lepidopilum fontanum* Mitt. in *Journ. of the Linn. Soc.* 1863, p. 4.

Ster. — *Afr.* : Madeir., ins. Azor.

*Goebelii* C. M. in *Fl.* 1897, p. 335.

*Ad fol. emort.* — *Am. merid.* : And. Venezuel.

*Herminieri* (W. P. Sch.) Jaeg. *op. cit.* p. 241.

*Lepidopilum Herminieri* W. P. Sch. in Besch. *Fl. bryol. Antill. fr.*,  
p. 54 (1876).

Fr. ign. — *Am. sept.* : Guadalup.

*heterodontium* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1893, p. 195.

Ster. *Filicic.* — *Am. sept.* : Costa Rica.

*jamaicense* C. M. in Hedwig. 1898, p. 245.

Ster. — *Am. sept.* : Jamaica.

*Erstedtianum* C. M. in *Fl.* 1875, n. 34-35.

Ster. — *Am. sept.* : Costa Rica.

*orbiculatum* C. M. in Hedwig. 1898, p. 244.

Ster. — *Am. sept.* : Porto Rico.

*Patrisiae* (Brid.) C. M. in Linn. 1874, p. 612 p.p.

*Hypnum Patrisiae* Brid. *Bryol. univ. II*, p. 539 (1827).

2. *Ad fol. emort.* — *Am. merid.* : Brasil. austr. or.

*phragmidiaceum* C. M. in *Fl.* 1897, p. 335.

Ster. — *Am. merid.* : And. Venezuel.

*porto-ricense* C. M. in Hedwig. 1898, p. 244.

Ster. *Cortic.* — *Am. sept.* : Porto Rico.

*radulaeforme* C. M. in Malpighia, 1896, p. 513.

Ster. — *Pac.* : Nov. Guin. brit.

*ramulicolum* C. M. in *op. cit.* p. 514.

Ster. — *Pac.* : Nov. Guin. brit.

*Sellowii* C. M. in Linn. 1874, p. 612.

*Hookeria Patrisiae* Hpe.  *Ic. M. t. I* (1814); *C. M. Syn. II*, p. 198 p.p. (1851).

2. *Ad fol. emort.* — *Am. merid.* : Brasil. austr. or.

*Sintenesii* C. M. in *Fl.* 1898, p. 244.

Ster. — *Am. sept.* : Porto Rico.



- Splitgerberi (*Mont.*) *C. M. in Linn.* 1874, p. 613.  
*Hookeria Splitgerberi Mont. Syll.* p. 14 (1856).  
*Lepidopilum Splitgerberi Mitt. M. austr. am.* p. 370 (1869).  
 Fl. et fr. ign. Ad fol. emort. — *Am. merid.* : Guian. batav.  
*Spruceanum C. M. in op. et l. cc.*  
*Lepidopilum Patrisiae Mitt. op. cit.* pp. 369-370 p.p.  
 2. — *Am. merid.* : And. Quitens.  
*subepiphyllum (Besch.) Jaeg. Ad. II,* p. 241 (1874-75).  
*Lepidopilum subepiphyllum Besch. Fl. bryol. Antill. fr.* p. 54 (1876).  
 Fr. ign. — *Am. sept.* : Guadalup.  
*tenellum C. M. in Fl.* 1897, p. 335.  
 Ster. Ad fol. emort. — *Am. merid.* : And. Venezuel.  
*Ulei C. M. in Ule Bryoth. brasil. n.* 156.  
 Ad ramul. fruticet. — *Am. merid.* : Brasil. austr. or.  
*Wallisii C. M. in Fl.* 1875, n. 34-35.  
 Ster. — *Am. merid.* : And. Nov. Granat.

**CRYPTHÆA** Mohr. in Web. Tab. syn. M. (1803). — 76 spec. (1).

- acuminata Hook. f. et Wils. Fl. of the N. Zeal.,* p. 102, t. 28, f. 4 (1855)  
*et Handb.* p. 461 (1867).  
 — *Pac.* : Nov. Seeland. sept.  
*affinis Mitt.* — *Cr. ramosa.*  
*apiculata Bryol. eur. V, Mon.* p. 5 (1850); *C. M. Syn. II,* p. 675 (1851).  
*Cr. cuspidata W. P. Sch. in Besch. Prodr. bryol. mexic.* p. 69 (1871).  
 — *Am. sept.* : Mexic.  
*apiculata Mitt.* — *Cr. Mittenii.*  
*arborea Lindb.* — *Cr. heteromalla.*  
*attenuata Bryol. eur. l. c.; C. M. op. cit.* p. 674; *W. P. Sch. in op. cit.*  
 p. 72.  
*Cr. patens Hpe. in hb. Schlechtendalano! C. M. in Linn.* 1814, p. 679 p.p.  
*Cr. protensa C. M. in hb. Schlechtendalano!*  
 — *Am. sept.* : Mexic., Orizaba.  
*Aurantiorum C. M. in Linn.* 1880-82, p. 469.  
 — *Am. merid.* : Argentin. Buenos Ayr.  
*Blumeniana Hpe. Enum. M. Brasil.* p. 38 (1879).  
 — *Am. merid.* : Brasil. austr. or.  
*boliviana W. P. Sch. in Mandon Pl. Boliv. n.* 1668; *C. M. in Nuov.*  
*Giorn. bot. ital.* 1897, p. 140.  
 — *Am. merid.* : And. Boliv.  
*brachycarpa C. M. in op. cit.* p. 141.  
 — *Am. merid.* : And. Boliv.  
*brasilensis Hpe.* — *Cr. filiformis.*  
*Breuteliana W. P. Sch. in Breut. M. capens.*  
 — *Afr.* : Pr. B. Sp. (Montagne-Pass).

---

(1) Spec. omn. monoicae et, ut videtur, corticolae.

- brevipila* Mitt. *M. austr. am.* p. 413 (1869).  
— *Am. merid.* : M. Chimborazo.
- caldensis* Aongstr. in *Oefv. af K. Vet.-Akad. Foerh.* 1876, n. 4, p. 31.  
— *Am. merid.* : Brasil. austr. or.
- calomira* Broth. in litt.  
    *C. gracilis* ej. in *Mosen Pl. brasil.* n. 88.  
— *Am. merid.* : Brasil.
- caripensis* C. M. — *Acrocryphaea squarrulosa*.  
*Coffeae* C. M. — *Acrocryphaea*.  
*concaifolia* Mitt. — *Acrocryphaea*.
- consimilis* Mont. in *Ann. sc. nat.* 1845, p. 100 et *Syll. p.* 25 (1856).  
    *Pilotrichum consimile* C. M. *Syn. II*, p. 166 (1851).  
— *Am. merid.* : Chile austr., Patagon.
- consimilis* H. f. et W. — *Cr. parvula*.  
*crenulata* Mitt. — *Cyrtodon*?
- cuspidata* Sull. in *U. St. Expl. Exped. t.* 23 A.  
— *Am. merid.* : Chile.
- cuspidata* { Hpe. — *Cr. Lindigiana*.  
              { W. P. Sch. — *Cr. apiculata*.  
*decurrens* Mitt. — *Cr. patens* var. *decurrens*.
- dentata* Mitt. in *Journ. of the Linn. Soc.* 1886, p. 311.  
— *Afr.* : M. Usagara.
- dilatata* H. f. et W. — *Cyrtodon*.
- Duseni* C. M. in *Dusen M. Camer.* n. 675.  
— *Afr.* : Cameroon.
- evanescens* C. M. in *Ule Coll. M. brasil.* n. 736 et 903.  
— *Am. merid.* : Brasil. austr. or.
- exigua* (C. M.) Jaeg. *Ad. II*, p. 95 (1874-75).  
    *Pilotrichum exiguum* C. M. *Syn. II*, p. 166 (1851).  
— *Afr.* : Pr. B. Sp., Transvaal.
- fasciculata* Duby in *Besch. Fl. bryol. Nouv. Caled.* p. 221 (1873).  
    *Dendropogon fasciculatus* Jaeg. *Ad. II*, p. 103 (1874-75).  
— *Pac.* : Nov. Caledon.
- fasciculosa* Mitt. *M. austr. am.* p. 411 (1869).  
— *Am. merid.* : And. Quitens.
- ferruginea* { Lindb. — *Acrocryphaea*.  
              { Mitt. — *Dendropogon*.
- filiformis* (Sw.) Brid. *Bryol. univ. II*, p. 251 (1827).  
    *Cr. brasiliensis* Hpe. in sched.  
    *Daltonia filiformis* W. Arn. *Disp.* p. 51 (1825).  
    *Fontinalis filiformis* Sw. *Prodr. fl. Ind. occ.* p. 138 ( ).  
    *Hypnum filiforme* Gmel. *Syst. nat. II*, p. 1340 (1791).  
    *Neckera filiformis* Hedw. *M. frond. III*, p. 41, t. 16 (1787-94).  
    *Pilotrichum filiforme* P. B. *Prodr.* p. 83 (1805); C. M. *Syn. II*, p. 169 (1851).  
— *Am. sept.* : Mexic., Guatemala, Cuba, S. Doming.,  
    Porto Rico. *Am. merid.* : Venezuela, Columb., Brasil.

- filiformis* { Hsch. — Cr. Schiedeana.  
 { Sull. } (M. Cub.). — *Acrocryphaea Coffeae* (fide C. M.).  
 { (M. Allegh.). — Cr. *glomerata*.
- funalis* C. M. in *Hedwig*. 1898, p. 241.  
 Ad *Retam.* — *Am. sept.* : Porto Rico.
- furcinervis* C. M. in *Linn.* 1880-82, p. 467.  
 — *Am. merid.* : Argentin. subtropic.
- Gardneri* Mitt. { (pl. Brasil.) — *Acrocryphaea ferruginea*.  
 { (pl. Cub.) — *Acrocryphaea Coffeae*.
- glomerata* *Bryol. eur.* V, *Mon.* p. 5 (1850); *Lesq. et Jam. Man.* p. 276 (1884).  
 Cr. *filiformis* *Sull. M. Allegh. n.* 81 (1846).  
*Pilotrichum heteromallum* γ *glomeratum* C. M. *Syn. II*, p. 168 (1851).  
 — *Am. sept.* : Ohio, Georg., Carol. ambo, Ludovic., etc.; Columb. brit.
- “ *var. pendula* (*Lesq. et Jam.*) *Par. Ind.* p. 287 (1894).  
 Cr. *pendula* *Lesq. et Jam. in Proceed. of the Amer. Acad. XIV*, p. 138 et *Man. l. c.*  
*Daltonia heteromalla* var. *Drumm.* M. *bor. am. II ed.*, p. 99.  
 Ad *Tillandsiam.* — *Am. sept.* : Georg., Florid., Ludovic., Alabama.
- Gorveana* { Mont. — *Dendrocryphaea*.  
 { W. P. Sch. (pl. Lechl.). — *Dendrocryphaea Lechleri*.
- gracilis* Mitt. *Fl. Vit.* p. 393 (1871).  
 — *Pac.* : Ins. Fidji, Samoa.
- gracilis* Broth. — Cr. *calomira*.
- helictophylla* { Aongstr. — *Papillaria Aongstroemii*.  
 { Mont. — *Meteorium*.  
 { Wils. — *Lindigia imponderosa*.
- Heuscheni* C. M. *mss.*; Aongstr. in *Oefv. af K. Vet.-Akad. Foerh.* 1876, n. 4, p. 29.  
 — *Am. merid.* : Brasil. austr. or.
- heteromalla* (*Dill.*) *Mohr l. c.*; *Brid. Mant. M.* p. 139 (1819) et *Bryol. univ. II*, p. 250 (1827); *Bryol. eur. V*, t. 438 (1850); *W. P. Sch. Syn. II ed.* p. 561.  
 Cr. *arborescens* *Lindb. Bidrag till. Moss. Syn.* p. 10 (1863).  
 Cr. *vermicularis* *Brid. Bryol. univ. II*, n. 253 ? (1827).  
*Daltonia heteromalla* H. et T. *Muscol. brit.* p. 81, t. 22 (1818).  
*Fontinalis secunda* *Linn. j. Meth. M.* p. 368 (1787).  
 ? *Grimmia arborescens* *Schrank, Baier. Fl. II*, p. 445 (1789).  
*Hypnum heteromallum* *Gmel. Syst. Nat.* p. 1340 (1791).  
 H. *polyccephalum* *Neck. Meth. M.* p. 190 (1771).  
*Neckera heteromalla* *Hedw. Descr. III*, p. 38, t. 15 (1792).  
 ? *Phascum repens* *With. Bot. Arrang.* p. 785 (1776).  
*Pilotrichum arborescens* P. B. *Prodr.* p. 37 (1805).  
 P. *heteromallum* C. M. *Syn. II*, p. 167 (1851).  
*Sphagnum arborescens* *Huds. Fl. angl. I ed.* p. 396 (1762).  
 Sph. *heteromallum, polyccephalum* *Dill. Hist. M.* p. 248, t. 42, f. 6 (1741).  
 Ad *Oleas, Popul., Ulmos*, etc. — *Eur.* : Britann., Neerland., vall. Rhen.; occid. et merid. *Afr.* : Alger., ins. Azor., Canar.
- “ *var. Lamyana* (*Mont.*) *Husn. Muscol. gall.* p. 289 (1892).  
 Cr. *arborescens* β *Lamyana* *Boulay M. Fr. I*, p. 188 (1884).  
 C. *heteromalla* var. { *aquatica* de Not. *Epil.* p. 218 (1869).  
 { *aquatilis* *Wils. Bryol. brit.* p. 420 (1855); *W. P. Sch. Syn. I ed. l. c.* (1876); *Par. Ind.* p. 288 (1894).

- Cr. *Lamyi* C. M. in *Linn.* 1844, p. 680; *Lindb. in Medd. p. Faun. et Fl. fenn.* 1881, n. VI, pp. 71-75.  
*Daltonia Lamyana* Mont. in *Ann. sc. nat. II, VI*, p. 327, t. 18 (1837) et *Syll.* p. 22 (1856).  
*Pilotrichum heteromallum* var. *aquatile* C. M. *Syn. II*, p. 167 (1851).  
 Ad arb., rup. et sax. ab aqua fluit. superfusa. — *Eur.* : *Angl. merid.*, *Gall. occ.*, *Etrur.*, *Lusit.*  
*heterophylla* Brid. — *Neckera*.  
*hygrophila* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 142.  
     — *Am. merid.* : *And. Boliv.*  
     var. *nitens* C. M. — *Cr. nitens*.  
*imbricata* Mitt. — *Leskea*.  
*indica* Mitt. — *Forsstroemia*.  
*intermedia* C. M. in *Linn.* 1847, p. 212.  
     *Pilotrichum intermedium* ej. *Syn. II*, p. 171 (1851).  
     — *Am. sept.* : *Mexic.*  
 ? *inundata* Nees in *Pl. Maxim. v. Wied*, p. 27 (     ); *Sull. M. of U. St.* p. 56 (1856) (1).  
     *Aquat.* — *Am. sept.* : *Illinois*.  
*Jamesoni* Tayl. in *Lond. Journ. of bot.* 1848, p. 192.  
     *Pilotrichum Jamesoni* C. M. *Syn. II*, p. 172 (1851).  
     — *Am. merid.* : *And. Quitens.*  
*Lamyi* C. M. — *Cr. heteromalla* var. *Lamyana*.  
*latifolia* Mitt. *M. austr. am.* p. 411 (1869).  
     — *Am. merid.* : *And. Quitens.*  
*laxifolia* Mitt. in *Journ. of the Linn. Soc.* 1886, p. 310.  
     — *Afr.* : *M. Usagara*.  
*Lechleri* C. M. — *Dendrocryphaea*.  
*Leiboldi* C. M. — *Acrocryphaea mexicana*.  
*leiophylla* *Bryol. eur. V, Mon.* p. 5 (1850); *W. P. Sch. et Par. Mon. Cryph. ined.* ; (2) *C. M. Syn. II ed.* p. 674 (1851).  
     — *Am. sept.* : *Mexic.*  
*leptoclada* Sull. — *Leucodon domingensis*.  
*leptophylla* *W. P. Sch. Mon. Cryph. ined.* ; *Besch. Pr. bryol. mexic.* p. 70 (1871) *excl. syn.* *Cr. leiophylla*.  
     — *Am. sept.* : *Mexic.*  
*leptopteris* C. M. in *Ule Bryoth. brasil. n.* 71.  
     — *Am. merid.* : *Brasil. austr. or.*  
*Lindigiana* *W. P. Sch. et Par. Mon. Cryph. ined.* ; *Par. Ind.* p. 982 (1894).  
     *C. cuspidata* Hpe. in *Ann. sc. nat.* 1866, IV, p. 353.  
     — *Am. merid.* : *And. Nov. Granat.*  
*longifolia* *W. P. Sch.* — *Cr. nitidula* var. *longifolia*.

---

(1) \* E descriptione pauperrima nil aliud nisi *Dichelyma capillaceum* vel *subulatum* videtur \*. *Lesq. et Jam. Man.* p. 413.

(2) *Peristomium simplex* !

- Lorentziana* C. M. in *Linn.* 1880-82, p. 468.  
— *Am. merid.* : Argentin. subtropic.
- lutescens* W. P. Sch. *Mon. Cryph. ined.* ; *Par. Ind.* p. 289 (1894).  
— *Am. sept.* : Mexic.
- madagassa* C. M. in *Abhandl. Bremen*, VII, p. 210 (1881).  
— *Afr.* : Madagascar.
- Malmiei* Broth. in *Bihang till K. Sv. Vet.-Akad. Foerh.* 1900, p.  
— *Am. merid.* : Brasil. austr. or. (Rio Grande do Sul).
- manoclada* (1) Aongstr. in *Oefv. af K. Vet.-Akad. Foerh.* 1876, n. 4, p. 30.  
*Cr. nanoclada* (erratum!) *Par. Ind.* p. 289.  
— *Am. merid.* : Brasil. austr. or.
- maynensis* Tayl. — *Acrocryphaea julacea*.
- microcarpa* W. P. Sch. — *Cr. nitidula*.
- microcyathea* Jaeg. — *Hedwigia*.
- Mittenii* Jaeg. *Ad. II*, p. 96 (1874-75).  
*Cr. apiculata* Mitt. *M. austr. am.* p. 411 (1869).  
— *Am. merid.* : Ins. Gorgon.
- mollis* Dusen M. *Am. austr. n.* 537.  
— *Am. merid.* : Patagon. occid.
- Moseni* Broth. in *Mosen Pl. brasil. n.* 87 a et in *Bihang till K. Sv. Vet.-Akad. Foerh.* 1895, n. 3.  
— *Am. merid.* : Brasil. or. (Minas Geraës).
- Mülleriana* W. P. Sch. — *Cr. Sartorii*.
- nanoclada* Par. (erratum!) — *Cr. manoclada*.
- nervosa* (Hook. f. et Wils.) *Bryol. eur. V, Mon.* p. 5 (1850) ; *Lesq. et Jam. Man.* p. 277 (1884).  
*Daltonia nervosa* H. f. et W. in *Lond. Journ. of bot.* 1842, II, p. 420, t. 24 et in *Drumm. M. bor. am. II ed. n.* 100 ; *Sull. Ic. M.* p. 109, t. 68 (1864).  
*Pilotrichum nervosum* C. M. *Syn. II*, p. 169 (1851).  
— *Am. sept.* : Alabama, Ludovic.
- nitens* W. P. Sch. in *Mandon Pl. Boliv. n.* 1689 et *Mon. Cryph. ined.*  
*Cr. hygrophila* var. *nitens* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 142.  
— *Am. merid.* : And. Boliv.
- nitidula* W. P. Sch. *Mon. Cryph. ined. et in Besch. Prodr. bryol. mexic.* p. 69 (1871).  
*Cr. microcarpa* ej. in sched.  
— *Am. sept.* : Mexic., And. Costa Ricens.
- « var. *longifolia* W. P. Sch. *Mon. Cryph. ined.* ; *Par. Ind.* p. 290 (1894).  
*Cr. longifolia* ej. in sched.  
— *Am. sept.* : Mexic.
- novo-granatensis* Hpe. in *Ann. sc. nat.* 1866, IV, p. 352.  
— *Am. merid.* : And. Nov. Granat.
- orbifolia* Besch. in *Journ. de bot.* 1891, p.  
— *Am. merid.* : Brasil. austr. or., Monte Video.
- orizabensis* W. P. Sch. — *Cr. subsimplex*.

---

(1) « Nonne rectius *monoclada* ? » (Hagen in litt.).

- ovalifolia* (C. M.) Jaeg. *Ad. II*, p. 98 (1874-75).  
*Philudora ovalifolia* Mitt. in F. v. Müller's *Fragm. phyt. austr. XI, Suppl. p. 14*.  
*Pilotrichum ovalifolium* C. M. in *bot. Zeit.* 1851, p. 564.  
 — Pac. : Austral. or. (Queensland, Nov. Vales., Victoria).
- pachycarpa* W. P. Sch. *Mon. Cryph. ined. et in Besch. Prodr. bryol. mexic. p. 71* (1871).  
 — Am. sept. : Mexic.
- paraguensis* Besch. — *Acrocryphaea*.
- parvula* Mitt. in *Handb. of the N. Zeal. Fl. p. 460* (1867).  
*Cr. consimilis* H. f. et W. *Fl. of the N. Zeal. II*, p. 101 (1855) et *Fl. Tasman. II*, p. 204 (1860).  
 — Pac. : Austral., Nov. Seeland., Tasman.
- patens* Hsch. in *Deppe et Schiede M. mexic. (excl. syn. Hypnum patens Hook.)*; W. P. Sch. et Par. *Mon. Cryph. ined.*; Mitt. *M. austr. am. p. 413* (1860).  
*Pilotrichum Deppii* (nomen mutatum) C. M. *Syn. II*, p. 674 (1851).  
*P. decurrens* ej. *op. cit. p. 172*.  
 — Am. sept. : Mexic. Am. merid. : And. Quitens.
- \* *var. decurrens* (C. M.) W. P. Sch. et Par. *Mon. Cryph. ined.*; Par. *Ind. p. 290* (1894).  
*Cr. decurrens* Mitt. *M. austr. am. p. 414* (1869).  
*Pilotrichum decurrens* C. M. *Syn. II*, p. 172 (1851).  
 — Am. sept. : Mexic.
- patens* Hpe. — *Cr. attenuata*.
- pendula* Lesq. et Jam. — *Cr. glomerata var. pendula*.
- pilifera* Tayl. in *Spr. M. Amaz. et And. n. 1287*; Mitt. *op. cit. p. 412*.  
 — Am. merid. : And. Nov. Granat., Quitens. et Boliv.
- pinnata* Bryol. eur. *V, Mon. p. 5 (nom.)* (1850); C. M. *Syn. II*, p. 675 (1851); W. P. Sch. in *Besch. Prodr. bryol. mexic. p. 70* (1871).  
 — Am. sept. : Mexic.
- plicata* Nees. — *Garovaglia*.
- polycarpa* W. P. Sch. *Mon. Cryph. ined. et in Besch. op. cit. p. 71*.  
 — Am. sept. : Mexic.
- protensa* Br. et W. P. Sch. in W. Schimp. *M. Abyss. II*, n. 87.  
*Pilotrichum protensum* C. M. *Syn. II*, p. 170 ? (1)  
 — Afr. : Abyssin.
- pulchella* Par. et W. P. Sch. *Mon. Cryph. ined.*; Par. *Ind. p. 291* (1894).  
*Cr. tenella* Mitt. *M. austr. am. p. 414*.  
 — Am. merid. : Argentin., Chile.
- Raddiana* (Brid.) Hpe. *Enum. M. brasil p. 39* (1879), nom.  
*Hypnum Raddianum* Brid. *Bryol. univ. II*, p. 511 (1827); C. M. *Syn. II*, p. 510 (1851).  
 — Am. merid. : Brasil. austr. or.

---

(1) *Peristomium simplex* ! nec duplex, ut ait C. M. l. c.

*ramosa* Wils. in Spr. M. Amaz. et And. n. 1284 b.

*C. affinis* Mitt. in sched.

— *Am. merid.* : And. Quitens. et Boliv., Brasil.

austr. or.

*Ravenelii* Aust. in Coult. Bot. Gaz. II, p. 89; Lesq. et Jam. Man. p. 277 (1).

— *Am. sept.* : Georg.

*reticulata* Besch. Prodr. bryol. mexic. p. 69.

— *Am. sept.* : Mexic.

*rhacomitrioides* C. M. in Linn. 1880-82, p. 466.

— *Am. merid.* : Argentin. Tucuman.

« *var. gracilis* ej. in op. et l. cc.

Ut et ubi typ.

*rubricaulis* Mitt. — *Acrocryphaea*.

*rufescens* C. M. — *Dendropogon*.

*Rutenbergii* C. M. in Abhandl. Brem. VII, p. 210 (1881).

— *Afr.* : Madagascar.

*Sartorii* W. P. Sch. Mon. Cryph. ined. et in Besch. op. cit. p. 72.

*Cr. Mülleriana* ej. in sched.

— *Am. sept.* : Mexic. (Orizaba).

*scariosa* C. M. in Fl. 1890, p. 486.

Ster.

— *Afr.* : Kilima N'Djaro.

*Schiedeana* (C. M.) Mitt. op. cit. p. 413.

*C. filiformis* Hsch. in Deppe et Schiede M. mexic.

*Pilotrichum Schiedeana* C. M. Syn. II, p. 167.

— *Am. sept.* : Mexic.

*Schleinitziana* C. M. — *Dendropogon*.

*sphaerocarpa* (Hook.) Brid. Bryol. univ. II, p. 251 (1827); Mitt. M. Ind. or. p. 125 (1859).

*Neckera sphaerocarpa* Hook. in Trans. of the Linn. Soc. IX, p. 312, t. 27.

*Pilotrichum sphaerocarpum* C. M. op. cit. p. 170.

— *As.* : Nepal., Himalaya.

*squarrulosa* Hpe. — *Cyrtodon*.

*subintegra* Ren. et Card. in Bull. Soc. roy. bot. Belg. 1892, II, p. 105.

— *Afr.* : Madagascar.

*subsimplex* W. P. Sch. Mon. Cryph. ined. ; Par. Ind. p. 292 (1894).

*C. orizabensis* ej. in sched. et in Besch. (Cr. Orizabae) op. cit. p. 70.

— *Am. sept.* : Mexic. (Orizaba).

« *var. congesta* W. P. Sch. et Par. Mon. Cryph. ined. ; Par. op. et l. cc.

Ut et ubi typ.

*tahitica* Besch. in Bull. Soc. bot. Fr. 1898, p.

— *Pac.* : Taïti.

---

(1) « Probabiliter *Leptodon* » L. et J. l. c.

*tasmanica* Mitt. in *Journ. of the Linn. Soc.* 1859, p. 90; *Hook. f. et Wils. Fl. Tasman. II*, p. 204, t. 175, f. 9 (ic. *pessima*!) (1860) et *Handb. of the N. Zeal. Fl.* p. 462 (1867).  
*Rupic.* — *Pac.* : Nov. Seeland., Tasman.

*tenella* Hsch. in *Sieb. Pl. Nouv. Holl.*

*Neckera tenella* Schwaegr. *Suppl. II, P. II*, t. 198 (1826).

*Pilotrichum tenellum* C. M. *Syn. II*, p. 166 (1851).

— *Pac.* : Austral. or. (Nov. Vales.).

*tenella* Mitt. — Cr. *pulchella*.

*tenuicaulis* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 141.

— *Am. merid.* : And. Boliv.

*tenuissima* H. f. et W. — *Meteorium*.

*vermicularis* Brid. — Cr. *heteromalla*?

*viridissima* C. M. in *Wild. Pl. Austral. n.* 465.

— *Pac.* : Austral. or. (Queensland).

*vitiana* C. M. — *Dendropogon Schleinitzianus*.

*Wolwitschii* Duby in *Mém.... Genève*, 1871, p. 3, t. 2, f. 1.

*Ad Coffeam arab.* et *Diospyr.* — *Afr.* : Angola, M. Usagara, Kilima N'Djaro, Madagascar.

**Cryphæadelphus** C. M. *Syn. II*, p. 144 (subsect. NECKERÆ) (1851).

**Cryphæopsis** C. M. in *Bull. hb. Boissier*, 1898, p. 116 (*nom.*, sect. METEORII).

4 **CRYPHIDIUM** Mitt. *M. austr. am.* p. 457 (sect. NECKERÆ) (1869); *Jaeg. Ad. II*, p. 178 *gen. propr.* (1874-75).

*leucocoleum* Jaeg. — *Neecka*.

**CRYPHIUM** P. B. *mss.*

*vaginans* ej. — *Calymperes Palisoti*.

**CRYPTANGIUM** C. M. in *Linn.* 1843, p. 599.

*gymnostomum* Card. }  
*Schomburgkii* C. M. } *Hydropogonella gymnostoma*.

**CRYPTOCARPON** Dz. et Mk. in *Ann. sc. nat.* 1844, *II*, p. 302 }  
**CRYPTOCARPUS** eor. *M. frond. ined. Arch. ind.* p. 37 (1854) } **DESMOTHECA**.

**CRYPTOGONIUM** Lindb. in *Oefv. af K. Sv. Vet.-Akad. Foerh.* 1873, p. 603.

*cylindricum* Lindb. — *Phyllogonium*.

**CRYPTOLEPTODON** Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1899, *II*, p. 238. — 1 spec.

*flexuosus* (Harv.) R. C. in *op. et l. cc.*

*Leptodon filiformis* W. P. Sch. in *hb. Hpe.*

*L. flexuosus* Jaeg. *Ad. II*, p. 106 (1874-75); *Par. Ind.* p. 728 (1896).

*Neckera camptoclada* R. C. in *op. cit.* 1895, *II*, p. 73.

*N. flexuosa* Harv. in *Hook. Journ. of bot.* 1840, p. 14 et in *Hook. Ic. pl. rar.* t. 21 (1841); *C. M. Syn. II*, p. 119 (1851).

2. — *As.* : Bhotan, Nepal, Himalaya, Calcutta.

**CRYPTOPODIA** Roehl. *Deutsch. Fl. III*, p. 82 (1813).

*pennata* ej. — *Neckera*.



**CRYPTOPODIUM** Brid. Bryol. univ. II, p. 30 (1827). — 7 spec.

*Bartramia* Schwaegr. — Cr. *bartramioides*.

*bartramioides* (Hook.) Brid. op. cit. p. 31; *H. f. et W. Fl. of the N. Zeal.* p. 88 (1855) et *Handb.* p. 446 (1867).

*Bartramia* Cryptopodium C. M. Syn. II, p. 507 (1851).

*Bryum* *Bartramia* Schwaegr. Suppl. II, P. II, p. 25, t. 160 (1826).

Br. *bartramioides* Hook. M. ex. t. 18 (1818-20).

Cr. *Bartramia* Schwaegr. Sp. M. p. 90.

2. — Pac. : Ins. Hawaiï, Nov. Seeland.

*bartramioides* Aongstr. — *Rhizogonium strictum*.

*brachyphyllum* (C. M.) Par. Suppl. Ind. p. 106 (1900).

*Bartramia* *brachyphylla* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 47.

— Am. merid. : And. Boliv.

« var. *viridissima* (C. M.) Par. op. et l. cc.

*Bartramia* *brachyphylla* var. *viridissima* C. M. in op. et l. cc.

Ubi typ.

*capillare* Hpe. Symb. 1874, in adn. n. 47 (nom.) et hb. ? (nunc in hb. British Museum).

2. — Am. merid. : Brasil. austr. or.

*Hookeri* Hpe. in Linn. 1847, p. 75.

*Bartramia* *Jamesoni* Tayl. in *Lond. Journ. of bot.* 1847, p. 334; C. M. Syn. I, p. 500 (1849) et II, p. 619 (1851); Mitt. M. austr. am. p. 269 (1869).

B. *viridissima* C. M. op. cit. I, p. 496.

*Leucodon* *bartramioides* Hook. Ic. pl. rar. I, t. 71 (1841).

Syn. Cortic. — Am. sept. : Jamaic. Am. merid. : And. Nov. Granat., Quitens. et Boliv.

« var. *borbonicum* Besch. Fl. bryol. Réunion, etc. p. 102 (1880-81).

— Afr. : Ins. Borbon. ?

*javanicum* Ren. et Card. in *Rev. bryol.* 1896, p. 100.

*Bartramia* *Wichurae* Broth. in sched.

*Leiomela* *Hookeri* Fleisch. M. Archip. ind. n. 33.

Syn. Filicic. — As. : Java.

*lutescens* (Hpe.) Jaeg. Ad. I, p. 538 (1873-74).

*Bartramia* *lutescens* Hpe. in *Ann. sc. nat.* 1863, p. 339; Mitt. op. et l. cc.

— Am. merid. : And. Nov. Granat.

*piligerum* Hpe. Symb. 1874, p. 47.

2. fr. ign. Cortic. et rupic. — Am. merid. : Brasil. austr. or.

**CRYPTOTHECA** Hsch. in *Deppe et Schiede M. mexic.*

Spec. omn. non indicatae sunt *PTEROBRYA* ejusd. nom.

*mexicana* W. P. Sch. — Garovaglia.

**CTENIDIUM** W. P. Sch. Syn. I ed. p. 631 (1860) (subgen. *HYPMI*); Mitt.

M. austr. am. p. 509 (1869), gen. propr.

Spec. omn. non indicatae sunt *HYPMNA* ejusd. nom.

*Berthelotianum* Mitt. — *Hylocomium*.

*molluscum* var. *procerum* Bryhn. — *Hypnum molluscum* var. *subplumiferum*.

CTENIUM W. P. Sch. op. et l. cc. (subgen. HYPNI).

CTENODONTIUM W. P. Sch. hb. (fid. Jaeg. Ad. II, p. 269).

*squarrosulum* ej. — Hookeria.

CUPRESSINA C. M. Syn. II, p. 229 (1851, sect. HYPNI), *postea* gen. propr.

Spec. omn. non indicatae sunt ECTROPTHECIA ejusd. nom.

*afro-cupressiformis* C. M. — Hypnum subafro-cupressiforme.

*alaris* ej. — Hypnum.

*anacamptopteris* ej. — Rhaphidostegium.

*arcuatipes* ej. (1897). — E. arcuatipes.

*entodonticarpa* ej. — Hypnum.

*flaris* C. M. — Hypnum.

*leptothalla* ej. — Hypnum.

*luridissima* ej. — Sematophyllum.

*minuta* ej. — Hypnum.

*producta* ej. — Rhaphidostegium.

*sanguiseta* ej. — Rhaphidostegium.

*sinensi-molluscum* ej. — Hypnum.

*subscabra* ej. — Rhaphidostegium.

*tereticautis* ej. — Hypnum.

**Cupressinadelphus** C. M. Syn. II, p. 265 (subsect. HOOKERIA) (1851).

CUSPIDARIA C. M. in Fl. 1896, p. 478 (sect. HYPNI, *postea* gen. propr.).

Spec. omn. sunt HYPNA ejusd. nom.

**Cyatophoridium** C. M. in Hedwig. 1898, p. 246 (sect. LEPIDOPILI, nom.).

**CYATOPHORUM** P. B. Prodr. p. 52 (1805). — 12 spec.

*Adiantum* (Griff.) Mitt. M. Ind. or. p. 147 (1859); *Bryol. jav. II*, p. 4, t. 134 (1861-70).

Neckera *Adiantum* Griff. Ic. pl. asiat. II, t. 85, f. 2? (1849).

2. — As. : Sikkim, Khasia, Bhotan, M. Java, ins. Batjan.

*bulbosum* (Hedw.) C. M. Syn. II, p. 54 (1851).

*Anoetangium bulbosum* Hedw. Sp. M. p. 43, t. 6 (1801).

*C. pennatum* Brid. Bryol. univ. II, p. 722 (1827).

*C. pteridioides* P. B. Prodr. p. 53 (1805).

*Hookeria pennata* Sm. in Trans. of the Linn. Soc. IX, p. 277; *Hook. M. ex. t.* 163 (1820).

*Leskea pennata* la Billard. Pl. Nov. Holl. II, p. 106, t. 253.

*Pterigophyllum pennatum* Brid. Mant. M. p. 154 (1819).

— Pac. : Austral. or., Nov. Seeland., ins. Auckland, Tasman.

« *var. apiculatum* Hook. f. et Wils. Handb. of the N. Zeal. Fl. p. 490 (1867).

— Pac. : Nov. Seeland. med.

« *var. minus eor.* Fl. Antarct. I, p. 143, t. 62, f. 3 (1844).

— Pac. : Nov. Seeland., ins. Auckland, Tasman.

*densirete* Broth. in Oesf. of K. Sv. Vet.-Akad. Foerh. 1893.

2. Cortic. — Pac. : Tasman.

*Dupuisii* R. C. — Rhacopilopsis.

*Hookerianum* (Griff.) Mitt. op. et l. cc.

*Neckera Hookeriana* Griff. op. cit. t. 84, f. 2 A.

— As. : Sikkim, Khasia.

*intermedium* Mitt. op. cit. p. 148.

Ster.

— As. : Himalaya occid.

*limbatulum* R. C. — *Epipterygium*.

*limbatum* R. C. — *Eriopus remotifolius*.

*Novae Zelandiae* Col. in *Trans. and Proced. of the N. Zeal. Inst.* 1886, p. 226.

— Pac. : Nov. Seeland.

*parvifolium* Bryol. jav. II, p. 5, t. 135.

2. Ster?

— As. : M. Java.

*penicillatum* C. M. in *Fl.* 1896, p. 457.

Ster.

— Pac. : Nov. Guinea.

*pennatum* Brid. — C. bulbosum.

*pteridioides* P. B. — C. bulbosum.

*splendidissimum* (Mont.) Hpe. et Ltz. in *Bot. Zeit.* 1866, p. 188.

*Hookeria speciosissima* Schwaegr. t. 328 (fid. Mitt.).

*H. splendidissima* Mont. in *Ann. sc. nat. II ser., IV*, p. 96 et in d'Orbigny, *Voy. Amer. VII, Crypt.* p. 110 (1839); *C. M. Syn. II*, p. 203 (1851).

*Lamprophyllum splendidissimum* W. P. Sch. *Coroll.* p. 101 (1856).

*Lepidopilum splendidissimum* Mitt. *M. austr. am.* p. 378 (1869).

2. Cortic. — *Am. merid.* : Chile, ins. Juan Fernandez, Patagon.

*sublimbatum* Thw. et Mitt. in *Journ. of the Linn. Soc.* 1872, p. 309.

Ster.

— As. : Ceylan.

*taitense* Besch. in *Ann. sc. nat.* 1894.

Ster. Cortic. ? — Pac. : Taiti.

*tenerum* Bryol. jav. II, p. 6, t. 136 (1831-70).

*Hookeria spinosa* Dz. et Mh. in *Zoll. Syst. Verz.* p. 26.

2. Ster. ?

— As. : Ins. Bima, Sumatra.

**CYCLODICTYON** Mitt. in *Journ. of the Linn. Soc.* 1863, p. 163 (*postea* sect. *HOOKERIA*).

*laete-virens* ej. — *Hookeria*.

**CYCNEA** Berk *Handb. Brit. M.* p. 301 (1863).

*curvicolla* ej. — *Phascum*.

**CYLICOCARPUS** Lindb. in *Oefv. af K. Sv. Vet.-Akad. Foerh.* 1862, n. 10, p. 605.

*lapponicus* ej. — *Amphoridium*.

*Mougeotii* ej. — *Amphoridium*.

**CYLINDROTHECIUM** Bryol. eur. fasc. 46-47 (1851).

Spec. omn. non indicatae sunt ENTODONTES ej. nom.

*abbreviatum* W. P. Sch. — *E. cyrtocladus*.

*angustifolium* Besch. — *E. Hahnianus*.

*cladorhizans* W. P. Sch. p.p. — *E. Schleicheri*.

*Drummondii* Mac. p.p. — *E. Macounii*.

*europaeum* Bryol. eur. — *E. cladorhizans*.

*gracilescens* Bryol. eur. — *E. Sullivantii*.

*Montagnei* Bryol. eur. — *E. concinnus*.  
*Mihlenbergii* Bryol. eur. — *E. seductrix*.  
*Regelianum* Bryol. eur. — *E. Drummondii*.

CYMBARIA *Tayl. in Journ. of bot.* VII, p. 190.

*Jamesoni* ej. — *Eustichia longirostris*.

CYNODON *Brid. Mant. M. p. 99* (1819) *et Bryol. univ. I, p. 500* (1826).

*cernuus* ej. — *Bryum pendulum*.  
*inclinatus* ej. — *Distichium*.  
*latifolius* ej. — *Desmatodon cernuus*.  
*luridus* ej. — *Didymodon*.  
*proscriptus* Hsch. — *Ditrichum*.

CYNODONTIELLA *Limpr. Laubm. Deutschl.... I, p. 281* (sect. CYNODONTII)  
 (1886); *Bryhn in N. Mag. f. Naturv. p. 116* (gen. propr.).  
 Spec. omn. sunt CYNODONTIA ejusd. nom.

→ CYNODONTIUM (Bryol. eur.) *W. P. Sch. Coroll. p. 12* (1855). — 13 spec.

alpestre (*Wahlenb.*) *Lindb. mss. (non Milde)*.

*Cynodontiella alpestris Bryhn op. et l. cc.*  
*Dicranum alpestre Wahlenb. Fl. lapp. p. 339* (excl. var.  $\beta$ ) t. 21 (1812).  
*Oncophorus alpestris Lindb. M. scand. p. 27* (1879).  
*O. (Oreweisia) obtusatus ej. in Medd. III, p. 192* (1878).

1. — *Eur.* : Lappon., Fennia.

\* *f. acutifolia Limpr. Nachtraeg. p. 649.*  
*Rupic.* — *Eur.* : Fennia, Helvet. (Gemmi).

alpestre } *Jur.* — *C. fallax*.  
 } *Milde.* — *C. torquescens*.

asperifolium *Lindb. in sched.* ; *Arn. in Rev. bryol. 1898, p. 8.*  
 1. — *As.* : Ala Tau.

*bicolor* (C. M.) *Par.* — *Oncophorus*.  
*bogotense* *Hpe.* — *Ditrichum pallidum var. strictum*.  
*boreale* *Kindb.* — *Philonotis*.  
*brevipes* (C. M.) *Par.* — *Cynodontium schisti var. brevipes*.  
*Bruntoni* *Bryol. eur.* — *Dicranoweisia*.  
*canadense* *Mitt.* — *Dichodontium*.  
*capillaceum* *Brid.* — *Distichium*.

*cernuum* } *Schleich.* — *Bryum pendulum var. arcuatum*.  
 } *Schwaegr.* — *Br. pendulum*.  
*cirratum* *Kindb.* — *C. gracilescens*.  
*conicum* *Mitt.* (1876). — *Blindia pulvinata*.

*crassirete Aongstr. in Oesv. af K. Vet.-Akad. Foerh. 1876, n. 4, p. 4.*  
 — *Am. merid.* : Brasil. austr. or.

*crispifolium* (*Mitt.*) *Jaeg.* — *Oncophorus*.  
*curvicaule* (C. M.) *Par.* — *Oncophorus*.  
*decumbens* (*Thw. et Mitt.*) *Jaeg.* — *Oncophorus*.

*debile* *Jaeg. p.p.* } — *Dicranella Heyrichiana*.  
 } — *Microdus exiguus*.  
 } — *Microdus Martianus*.

*fallax Limpr. Laubm. Deutschl.... I, p. 287* (1886).

*C. alpestre Jur. Laubmfl v. Oesterr. Ung. p. 26 p.p.* (1882).  
*C. gracilescens Un. itin. crypt. 1863, n. 6.*

*C. gracilescens* γ *alpestre* W. P. Sch. *Syn. II* ed. p. 62 (1876).

*Dicranum alpestre* β *majus* Wahlenb. *Fl. lapp* p. 309 (1812).

*Diobelon alpestre* Hpe. *Fl. Hercyn.* (1873).

1. Ad semit. cav. et in decliv. saxos. — *Eur.* : Riesengeb., Stir., Carinth., Helvet.

*flexicaule* Schwaegr. — *Leptotrichum*.

*glaucescens* (L. et A.) Par. — *Oncophorus*.

*gracilescens* (Web. et Mohr) W. P. Sch. *Coroll.* p. 12 et *Syn. II* ed. p. 62 *excl.* var. *alpestre* (1876).

*Campylopus cirratus* Brid. *Bryol. univ. I.* p. 479 (1826).

*Cynodontium cirratum* Kindb. *Laubm. Schwed. u. Norw.* p. 102 (1883).

*C. gracilescens* β *inflexum* W. P. Sch. *Syn. I* ed. p. 61 (1860).

*Dicranum gracilescens* W. et M. *Bot. Taschb.* p. 467 (1807); *Bryol. eur.*, fasc. 37-40 (Vol. I) p. 13, t. 45 (1847) *excl.* var. γ; W. P. Sch. *Syn. II* ed. p. 62.

*D. gracilescens* β *curvisetum* *Bryol. eur. I. c.*

*D. mixtum* de Not. *Mant. M. n.* 52 (1835).

*D. polycarpum* var. *curvisetum* C. M. *Syn. II*, p. 591 (1851).

*D. polycarpum* β *gracilescens* *ej. op. et l. cc.*

*Diobelon gracilescens* Hpe. *Fl. Hercyn.* (1873).

*Dryptodon Campylopus* Brid. *Bryol. univ. I.* p. 773 (1826).

*Oncophorus cirratus* Lindb. *M. scand.* p. 27 (1879).

*O. gracilescens* *ej. op. et l. cc.*

1. In fiss. rup. et in decliv. herbos. — *Eur.* : Reg. subalp., rarius alp. As. : Amour, Sikkim, Caucas, occ. et centr. *Am. sept.* : Groenland., Nov. Angl., M. Alb., Adirondack et Rupestr.

*gracilescens* Un. itin. 1863, n. 6. — *C. fallax*.

var. *alpestre* *Bryol. eur.* — *C. fallax*.

var. *curvisetum* } *Bryol. eur.* — *C. gracilescens*.

var. *inflexum* }

var. *microcarpum* Ltz. }

var. *pusillum* Pfeff. }

var. *tenellum* W. P. Sch. } — *C. torquesens*.

*Hookeri* Mitt. (ins. Kerg.) — *Blindia aschistodontoides* var. *nigrescens* ?

*inclinatum* Brid. — *Distichium*.

*latifolium* Schwaegr. — *Desmatodon cernuus*.

*laxirete* Greb. — *C. polycarpum* var. *laxirete*.

*Limprichtianum* Grebe in *Hedwig*. 1897, *Beibl.* n. 45, p. 103, t. X.

Ster. Ad rup. porphyrit. — *Eur.* : Westphal.

*longirostre* vel *longirostrum* { Schwaegr. — *Dicranodontium*.  
Zett. — *Leptotrichum flexicaule* var.  
longifolia.

*obtusatum* Kindb. — *Oreoweisia serrulata* var. *tenuior*.

*polycarpum* (Ehrh.) W. P. Sch. *Coroll.* p. 12 (1855) *excl.* var. β; de Not. *Epil.* p. 280 (1869).

*Bryum cirratum* setis et capsulis brevioribus et pluribus *Dill. Hist. M.* p. 278, t. 48, f. 41 (1741) (fid. C. M.).

Br. medium Vill. *Pl. Dauph. III*, p. 878 (1786).

Br. polycarpum Hoffm. *Deutschl. Fl. II*, p. 37 (1796).

*Cecalyphum multiflorum* P. B. *Prodr.* p. 51 (1805).

*Dicranum polycarpum* Ehrh. *Pl. crypt. exs.* n. 84 (1786); *Bryol. eur.* fasc. 37-40 (Vol. I), Mon. p. 14, t. 47 (1847); C. M. *Syn. II*, p. 591 (1851).

*D. scottianum* Drumm. *M. bcr. am. I* ed. n. 107.

*Didymodon Jenneri* W. P. Sch. mss.; *Howell in Trans. of the Edinb. bot. Soc.* IX, 1868, p. 198.

*Diobelon polycarpum* Hpe. *Fl. Hercyn.* (1873).

*Fissidens polycarpus* Hedw. *Descr.* II, p. 85, t. 31 (1788).

*Oncophorus polycarpus* Brid. *Bryol. univ.* I, p. 397 (1826).

*O. strumifer*  $\beta$  *substrumosus* ej. op. cit. p. 396.

1. In rup. silic. fiss. humid. — *Eur.*: Spitzberg., ins. Ursor.; e reg. campestr. usq. in alp. *As.*: Japon., Ainour, Caucas. or. *Am. sept.*: Groenland., Terr. nov.; Canada merid. or.; reg. sept. or., Alaska, ins. Vancouver, Columb. brit., Washington, Idaho, Wyoming.

« *var. brevifolium* C. Jens. *Consp. fl. Groenland.* p. 396 (1887).

— *Am. sept.*: Groenland.

*var. laevifolia* Hagen. — *C. polycarpum var. laxirete*.

« *var. laxirete* Dix. *Handb. brit. M.* p. 73 (1896).

*C. laxirete* Greb. in *Hedwig*. 1901, p. 106.

*C. polycarpum var. laevifolia* Hagen in *Tromsø Mus. Aarsh.* 1898-99, p. 8.

— *Eur.*: Anglia (Cheshire), Norveg.

*robustum* Kindb. — *Dicranoweisia*.

*schisti* (Wahlenb.) Lindb. in *Oesfv. af K. Vet.-Akad. Foerh.* 1864, p. 230 in not.; W. P. Sch. *Syn.* II ed. p. 61 (1876).

*Bryum foliis capillaribus subfasciculatis, seta bilineari, capsulis sphaeroides, operculo conico arcuato* Stroem. in *Skr. af Nat.-hist. Selsk.* I, 2, p. 32 et t. XI, n. 2 (1791) p.p. ?

*Br. hyperboreum* ej. in D. Vid. *Selsk. Skr., Ny Saml.* III, p. 358 (1788) p.p. ?

? *Br. schisti* Gunn. *Fl. norveg.* 21, p. 138 (1776).

*Cynodontiella schisti* Bryhn op. et l. cc.

*Dicranum splachnoides* Brid. *Musc. rec.* II, I, p. 81 (1798).

*Grimmia schisti* *Fl. brit.* III, p. 1185 1804-1805).

*Oncophorus schisti* Lindb. *M. scand.* p. 27 (1879).

*Rhabdoweisia schisti* *Bryol. eur. fasc.* 33-36 (*Vol. I*), *Mon.* p. 5, t. 43 (1846).

*Weisia schisti* Wahlenb. *Fl. lapp.* p. 325 (1812).

*W. striata*  $\zeta$  *minor* H. et T. *Musc. brit.* p. 45, t. 15 p.p. (1818).

1. In rup. praepr. schist. fiss. — *Eur.*: Anglia (*fid. C. M.*), Scand., Harz? Siles. infer., Bohem., Transilv., Salisb., Stir. merid., Tirol., Engadin., Helvet. *As.*: Jénissei super. *Am. sept.*: Groenland., Tefra nov., m. Rupestr.

« *var. brevipes* Lindb. mss.; *Linpr. Laubm. Deutschl.... I*, p. 283 (1886).

*Cynodontium brevipes* Par. *Ind.* p. 306 (1894).

*Oncophorus brevipes* Lindb. *M. scand.* p. 27 (1879).

— *Eur.*: Scand., Fennia.

*serrulatum* Jur. — *Oreoweisia*.

*sinense* (C. M.) Par. — *Oncophorus*.

*strumiferum* (Ehrh.) R. Hartm. *Bryac. scand. exs. fasc.* 10, n. 271 (1863) *fid. Lindb.*; *de Not. Epil.* p. 280 (1869).

*Bryum inclinans* Dicks. *Pl. crypt. fasc.* 4, p. II, t. II, f. 9 (1801).

*Br. strumiferum* ej. op. c. *fasc.* 3, p. 8 (1793).

*Cecalyphum strumiferum* P. B. *Prodr.* p. 52 (1805).

*Cynodontium polycarpon var. strumiferum* W. P. Sch. *Coroll.* p. 15 (1855).

*Dicranum gibbosum* Brid. *Sp. M. I*, p. 225 (1806).

*D. polycarpum*  $\zeta$  *strumiferum* *de Not. Syll.* p. 210 (1838); *Bryol. eur. fasc.* 37-40 (*Vol. I*), *Mon.* p. 11, t. 47  $\beta$  (1846).

- D. strumiferum* Ehrh. *Pl. crypt. ers. n.* 74 (1786; *Sic. Disp. M. Succ.* p. 33 (1799); *C. M. Syn. II*, p. 592 (1851).  
*Diobelon strumiferum* Hpe. *Fl. Hercyn.* (1837).  
*Fissidens strumifer* Hedw. *Descr. II*, p. 88, t. 32 (1789).  
*Hypnum strumosum* Gmel. *Syst. Nat. II*, p. 1339 (1791).  
*Onchophorus strumifer* Brid. *Bryol. univ. I*, p. 395 (1826).

1. Ad rup. silic. vel granit. umbr. — *Eur.* : Hic illic cum praeced. *As.* : Fret. Sinjawin, Amour, Jenissei med. et infer., Caucas. occ. *Am. sept.* : Groenland., Labrador, Ontario, lac. Super., Minnesota, M. Rupestr., Columb. brit., ins. Vancouver.

« *var. humile* C. Jens. *Consp. fl. Groenland.* p. 396 (1887).  
 — *Am. sept.* : Groenland.

« *var. scabrior* Hagen in Tromsø Mus. *Aarsh.* 1898-99, p. 8.  
 — *Eur.* : Norveg.

strumulosum C. M. et Kindb. in *Mac. Cat.* p. 16 (1892).  
 Rupic. secin torrent. — *Am. sept.* : M. Rupestr.

subalpestre Kindb. in *Mac. Cat.* p. 17 et 257 (1892).

*C. gracilescens* Mac. *Canad. M. n.* 20.

Dicranum *gracilescens* Aust. *M. Appal. n.* 467.

*D. polycarpum* Drumm. *M. bor. am. I ed. n.* 103 p.p.

Rupic. — *Am. sept.* : Labrador, Gaspes., lac. Huron, Super.

suecicum (Lindb. et Arn.) Pur. *Suppl. Ind.* p. 111 (1900).

Dicranum *Starkei* Kindb. *Eur. a. N. Amer. Bryin.* p. 191 p.p. (1897).

*Onchophorus suecicus* Lindb. et Arn. in *Rev. bryol.* 1895, p. 75 c. ic.

1. Rupic. — *Eur.* : Scandin.

*tenellum* Limpr. — *C. torquescens*.

*torquescens* (Bruch) Limpr. *Laubm. Deutschl.... I*, p. 288 (1886).

*C. alpestre* Milde *Bryol. siles.* p. 51 (1869).

*C. gracilescens* } *var. microcarpum* Ltz. in sched.  
 } ♂ pusillum Pfeffer, *Bryog. Stud.* p. 17 (1869).  
 } γ *tenellum* W. P. Sch. *Syn. I ed.* p. 61 (1860).

*C. tenellum* Limpr. *Kryptfl. v. Schles. I*, p. 425, n. 417 (1877).

Dicranum *gracilescens* γ *alpestre* Hüb. *Musc. germ.* p. 255 p.p. (1833).

*D. gracilescens* γ *tenellum* *Bryol. eur. fasc.* 37-40 (Vol. I), *Mon.* p. 13, t. 47 p.p. (1846).

*D. torquescens* Bruch in *Un. itin.* 1828.

*Onchophorus nigricans* Kindb. in *Bot. Not.* 1882, p. 146 et *Laubm. Schwed. u. Norw.* p. 143 (1883).

*O. torquescens* Arn. et Jens. in *Bihang t. K. Sv. Vet.-Akad. Handl.* n. 10 p. 53 (1896).

1. In fiss. mur. et rup. — *Eur.* : Lappon., Scandin., Siles., Salisb., Carinth., Bavar., Helvet., etc. *As.* : Caucas. *Am. sept.* : (sine loc. indic., fid. Broth.).

*trifarium* Brid. — *Didymodon luridus*.

*virens* W. P. Sch. (c. var.) — *Onchophorus* (c. var.).

*Wahlenbergii* Hartm. — *Onchophorus* (c. var.).

CYNONTODIUM Hedw. *Sp. M.* (1801).

Spec. omn. non indicatae sunt DITRICHIA ejusd. nom.

*australe* Mitt. — *Lophiodon strictus*.

*bogotense* Hpe. — *Ditrichum pallidum* var. *strictum*.

*capillaceum* Hedw. — *Distichium*.

*capillaceum* Mitt. — *Distichium setifolium*.

*cernuum* Hedw. — *Bryum pendulum*.  
*elongatum* Mitt. — *Trichostomum*.  
*inclinatum* Hedw. — *Distichium*.  
*strictum* Mitt. — *Ditrichum pallidum* var. *strictum*.  
*Trematodon* P. B. — *Trematodon longicollis*.  
*trifarium* Hedw. — *Didymodon luridus*.

**Cyphoma** Hook. f. et Wils. *Fl. Tasman. II*, p. 200 (sect. POLYTRICHI) (1860).

**CYPTODON** (1) Par. et W. P. Sch. Mon. Cryph. ined.; Par. Ind. p. 310 (1894). — 4 spec.

? *crenulatus* (Mitt.) Par. op. et l. cc.

*Cryphaea crenulata* Mitt. in *Journ. of the Linn. Soc.* 1859, p. 90.

*Dendropogon crenulatus* Jaeg. *Ad. II*, p. 103 (1874-75).

— Pac.: Austral. or. (Victoria), Tasman.

*dilatatus* (Hook. f. et Wils.) Par. et W. P. Sch. op. et ll. cc.

*Cryphaea dilatata* H. f. et W. *Fl. of the N. Zeal.* p. 102, t. 88, f. 2 (1855).

Rupic. — Pac.: Austral. or. (Victoria), Nov. Seeland.

*Mulleri* (Hpe.) Par. et W. P. Sch. op. et ll. cc.

*Dendropogon Mulleri* Hpe. in *Linn.* 1856, p. 212.

— Pac.: Austral. or.

*squarrosulus* (Hpe.) Par. et W. P. Sch. op. et ll. cc.

*Cryphaea squarrosula* Hpe. in *Linn.* 1859-60, p. 636; F. v. Müll. *Anal. draw. austr. M. t. XI* (1864).

*Dendropogon squarrosulus* Jaeg. op. et ll. cc.

— Pac.: Austral. or.

**CYRTODON** R. Br. *Suppl. ad Iter Parray.* (1824).

*splachnoides* ej. — *Dissodon*.

**CYRTO-HYPNUM** Hpe. *Moosb.* p. 20 (1871).

Spec. omn. sunt THUIDIA ejusd. nom.

**CYRTO-LESKEA** Hpe. (*fid. Sh. in Jaeg. Ad. II*, p. 788 (1877-78), spec. ANOMODONTIS vel LESKEÆ).

**CYRTOPOUS** Brid. *Bryol. univ. II*, p. 235 (sect. NECKERÆ) (1827); Hook. f. et Wils. *Fl. of the N. Zeal.* p. 1 (1855). — 3 spec.

*Cameruniae* Broth. in *Engler's Bot. Jahrb.* 1897, p. 254.

*Garovaglia simplex* C. M. in *Dusen M. Camer. n.* 444.

2. Ad arbor. — Afr.: Guinea gall., Cameroon.

*cryphaeoides* C. M. *M. Polyn.* p. 21 (1874).

Ster. Ad arbor. — Pac.: Ins. Viti.

*curtipendulus* Spr. — *Antitrichia*.

*setosus* (Hedw.) *Handb. of the N. Zeal. Fl.* p. 461 (1867).

*Anoetangium setosum* Hedw. *Sp. M.* p. 43, t. I, f. 4-6 (1801).

*Cladomnium setosum* H. f. et W. *Fl. of the N. Zeal. II*, p. 100 (1855).

---

(1) De κυρτω (procumbo) et οδους (dens, propter dentes (coriaceos) vivos madefactosve peristomii orificium c'upsulae fere horizontaliter obtegentes. Spec. omn. monoicae et, ut videtur, subaquaticae.



*Neckera setosa* Hook. *M. exot. t.* 7 (1818); *Brid. Bryol. univ. II*, p. 225 (1827).

*Pilotrichum setosum* C. M. *Syn. II*, p. 162 (1851).

*Spiridens setosus* Mitt. in *Journ. of the Linn. Soc. IV*, p. 89.

2. — *Am. merid.* : (*fid. Handb., sine loc. indic.*

*Pac.* : Ins. Hawaï, Nov. Seeland., Tasman.

*tailensis* W. P. Sch. — Garovaglia.

**DALTONIA** Hook. et Tayl. *Muscol. brit. I*, p. 80 (1818). — 58 spec.

*affinis* W. Arn. — *Pilotrichum*.

*androgyne* Geh. et Hpe. in *Fl.* 1881, p. 405.

1 vel syn.? — *Am. merid.* : Brasil. austr. or.

*angustifolia* Dz. et Mk. *M. frond. ined. Archip. ind. p.* 31, *t.* 13 (1854);

*C. M. Syn. II pp.* 18 et 660 (1851); *Bryol. jav. II*, p. 30 (1861-70).

1. Ad arb. et terr. — *As.* : Borneo, Java.

*apiculata* Mitt. *M. Ind. or. p.* 145 (1859).

1 ? — *As.* : Nepal.

*aristata* Geh. et Hpe. *Enum. M. brasil. p.* 49 (1879).

1. Lignic. — *Am. merid.* : Brasil. austr. or.

*aristifolia* Ren. et Card. in *Rev. bryol.* 1893, p. 105.

Ad *Trichoman.* — *As.* : Java.

*bilimbata* Hpe. in *Linn.* 1863, p. 151; *Mitt. M. austr. am. p.* 400 (1879).

1. Ad arbor. — *Am. merid.* : And. Nov. Granat.

*binervis* Hpe. — *Lepidopilum daltoniaceum*.

*bipinnata* W. Arn. — *Pilotrichum*.

*brasiliensis* Mitt. *M. austr. am. p.* 399 (1869).

1 vel syn.? — *Am. merid.* : Brasil. or.

*brevipedunculata* Mitt. *M. Ind. or. p.* 146 (1859).

1 vel syn.? — *As.* : Bombay.

*composita* W. Arn. — *Pilotrichum*.

*compressa* Mitt. *M. austr. am. p.* 400 (1869).

1 vel. syn. Cortic. — *Am. merid.* : And. Quitens.

*contorta* C. M. *Syn. II*, p. 660 (1851); *Bryol. jav. II*, p. 32, *t.* 155 (1861-70).

1. — *As.* : Ceylan, Java.

*crispata* *Bryol. eur. V (nom.)*; W. P. Sch. in *Besch. Prodr. bryol. mexic. p.* 84 (1871).

— *Am. merid.* : Mexic.

*curvuspis* C. M. in *Hedwig.* 1900, p. 263.

1. Ad ramul. — *Am. merid.* : Brasil. or. (Minas Geraës).

*disticha* W. Arn. — *Neckera*.

*Duseni* C. M. in *Dus. M. Camer. n.* 776; *Broth. in Engler's Bot. Jahrb.* 1897, p. 259.

*Syn. Cortic.* — *Afr.* : Cameroon.

*elegantula* W. P. Sch. *hb. (nunc in hb. Kew.)*; C. H. Wright. *M. Mad. p.* 6 (nom.)

— *Afr.* : Madagascar.

- Fendleri C. M. in *Linn.* 1878-79, p. 491.  
 l. — *Am. merid.* : Venezuela.
- filamentosa* W. Arn. — *Meteorium*.  
*filiformis* W. Arn. — *Cryphaea*.
- flexifolia* Mitt. *M. Ind. or.* p. 146 (1859).  
 l. — *As.* : Nepal., Ceylan.
- fuscescens* W. Arn. — *Papillaria*.
- gracilis* Mitt. *M. austr. am.* p. 402 (1869).  
 l vel syn. ? Cortic. et foliic. — *Am. merid.* : And. Quitens.
- Hampeana* Geh. in *Hpe. Enum. M. brasil.* p. 50 (1879) et in *Rev. bryol.* 1879, p. 67.  
 l. — *Am. merid.* : Brasil. austr. or.
- heteromalla* H. et T. — *Cryphaea*.  
 var. eor. — *Cryphaea glomerata* var. *pendula*.  
*heterophylla* W. Arn. — *Neckera*.  
*hypnoides* W. Arn. — *Pilotrichum*.  
*imbricata* Spreng. — *Pilotrichella versicolor*.
- intermedia* Ren. et Card. in *Bull. Soc. roy. bot. Belg.* 1894, II, p. 127.  
 Syn ? — *Afr.* : Madagasc., Comor. magn.
- intermedia* W. Arn. — *Neckera*.
- irrorata* Mitt. *M. austr. am.* p. 399 (1869).  
 l vel syn. Lignic. et ad sax. irror. — *Am. merid.* : And. Nov. Granat.
- Jamesoni* Tayl. in *Lond. Journ. of bot.* 1848, p. 283; C. M. *Syn. II*, p. 19 (1851); Mitt. *op. cit.* p. 401.  
 l. — *Am. merid.* : Summ. m. Pichincha.
- Lamyana* Mont. — *Cryphaea heteromalla* var. *Lamyana*.
- latimarginata* Besch. *Fl. bryol. Réunion, etc.* p. 136 (1880-81).  
 Syn. — *Afr.* : Ins. Borbon.  
 « var. *madagassa* Ren. *Podr. fl. bryol. Madag. etc.* p. 292 (1897).  
 — *Afr.* : Madagascar.
- Leucoloma* Hpe. *Symb.* 1874, p. 502.  
 l. — *Am. merid.* : Brasil. austr. or.
- Lindigiana* Hpe. in *Ann. Sc. nat.* 1866, p. 363.  
 l. — *Am. merid.* : And. Nov. Granat.
- longecuspidata* C. M. in *Bull. hb. Boissier*, 1897, p. 201.  
 — *Am. sept.* : Guatemala.
- longifolia* Tayl. in *Lond. Journ. of bot.* 1848, p. 283; C. M. *Syn. II ed.* p. 19 (1851); Mitt. *op. cit.* p. 401.  
 l. — *Am. merid.* : And. Quitens.
- longinervis* Mitt. in *Journ. of the Linn. Soc.* 1863, p. 163.  
 l. — *Afr.* : Ins. Fernando Po.
- longipedunculata* C. M. *Syn. II*, p. 18 (1851); *Bryol. jav. II*, p. 31, t. 154 (1861-70).  
 l. — *As.* : Borneo, Java.
- lorifolia* C. M. in *Fl.* 1875, n. 34-35.  
 Ster. Cortic. — *Am. sept.* : Costa Rica.

- Mac-Gregorii* Broth. in *Oefv. af Finska Vet.-Akad. Foerh.* 1898, p. 183.  
 l. — *Pac.* : M. Nov. Guin.
- macrotheca* Mitt. *M. austr. am.* p. 402 (1869).  
 l vel syn. ? — *Am. merid.* : M. Chimborazo.
- madagassa* W. P. Sch. *hb. (nunc in hb. Kew.)*; C. H. Wright *M. Mad.*  
 p. 6 (nom.). — *Afr.* : Madagascar.
- marginata* Mitt. *M. Ind. or.* p. 145 (1859).  
 l. Cortic. — *As.* : Khasia.
- minor* Besch. *Fl. bryol. Réunion, etc.* p. 136 (1880-81).  
 — *Afr.* : Ins. Bourbon.
- minutifolia* C. M. in *Nuov. Giorn. bot. ital.* 1897, p. 148.  
 — *Am. merid.* : And. Boliv.
- mucronata* *Bryol. jav. II*, p. 32, t. 156 (1861-70).  
 Syn. — *As.* : Java.
- nervosa* (Hook. f. et Wils.) Hpe. in C. M. *Syn. II*, pp. 17 et 660 (1851);  
*Handb.* p. 496 (1867).  
 D. Novae Zeelandiae Mitt. in *Journ. of the Linn. Soc.* 1859, p. 95.  
*Hookeria nervosa* H. f. et W. in *Lond. Journ. of bot.* 1844, p. 519 et *Fl.*  
*antarct. I*, p. 142, t. 61 (1844).  
 l. Ad ramulos arb. — *Pac.* : Nov. Seeland. sept., ins. Auckland.
- nervosa* H. f. et W. (1842). — *Cryphaea*.  
*Novae Zeelandiae* Mitt. — *Daltonia nervosa*.
- occania* C. M. in *Fl.* 1875, n. 34-35.  
 Syn. — *Am. merid.* : And. Nov. Granat.
- ovalis* Tayl. in *Lond. Journ. of bot.* 1846, p. 66; C. M. *Syn. II*, p. 19  
 (1851); Mitt. *M. austr. am.* p. 401 (1869).  
 l. Ad arbor. — *Am. merid.* : And. Quitens.
- patula* Mitt. in *Journ. of the Linn. Soc.* 1863, p. 163.  
 l. — *Afr.* : Ins. Fernando Po, M. Usagara, Kilima  
 N'Djaro.
- pennata* W. Arn. — *Neckera*.
- peruviana* Mitt. *M. austr. am.* p. 401 (1869).  
 l vel syn. — *Am. merid.* : And. Peruv.
- polytrichoides* W. Arn. — *Lepidopilum*.
- pulvinata* Mitt. *op. cit.* p. 398.  
 l vel syn. — *Am. merid.* : And. Nov. Granat.
- pusilla* Hook. f. et Wils. *Fl. Tasman. II*, p. 221, t. 177, f. 5 (1860).  
 l. Lignic. et ad font. — *Pac.* : Tasman.
- reticulata* C. M. in *Linn.* 1869-70, p. 20.  
 l. — *As.* : Ceylan.
- robusta* Aongstr. in *Oefv. af K. Vet.-Akad. Foerh.* 1873, n. 5, p. 177.  
 l. — *Am. merid.* : Ins. Gallapagos.
- scaberula* Mitt. *M. Ind. or.* p. 146 (1859).  
 l. — *As.* : Ceylan.
- scabriseta* W. Arn. — *Lepidopilum*.

- semitorta* Mitt. *op. et l. cc.*  
 1. — *As.* : Sikkim.
- sphaerica* Besch. in *Bull. Soc. bot. Fr.* 1898, p. 119.  
 1. *Ad Cyrtandram pendulam.* — *Pac.* : Taïti.
- splachnoides* (Sm.) Hook. et Tayl. *Muscol. brit. I ed.* p. 90, *tt.* 3 et 22 (1818); *Brid Bryol. univ. II*, p. 255 (1827); *C. M. Syn. II*, pp. 17 et 660 (1851); *Bryol. eur. V*, t. 449; *W. P. Sch. Syn. II ed.* p. 579 (1876).  
*Hookeria splachnoides* Tayl. in *Mackay's Fl. hibern. II*, p. 37 (1836).  
*Neckera splachnoides* Engl. *Bot. t.* 2564 (*excl. syn. Schwaegr.*).  
 1 et syn. *Ad rup. humid. umbros., rar. ad arb.* — *Eur.* : Hibern. *Afr.* : Madeir, ins. Fernando Po. *Am. sept.* : Mexic.; *Antill.* (*fid. W. P. Sch.*).
- stenoloma* Besch. *Fl. bryol. Réunion, etc.* p. 136 (1880-81).  
 1 vel syn. — *Afr.* : Ins. Bourbon.
- stenophylla* Mitt. *M. austr. am.* p. 402 (1869).  
 1 vel syn. — *Am. merid.* : And. Quitens.
- straminea* C. M. in *Trans. and Proceed. of the N. Zeal. Inst.* 1893, p. 276.  
 — *Pac.* : Nov. Seeland.
- strictifolia* Mitt. *M. Ind. or.* p. 146 (1859).  
 1 vel syn. — *As.* : Ceylan, Java.
- tenella* Broth. in *Act. Soc. sc. fenn.* 1891, n. 5.  
 1. — *Am. merid.* : Brasil. or. (Minas Geraës).
- tenuifolia* Mitt. *M. austr. am.* p. 402 (1869).  
 1 vel syn. *Ad arb.* — *Am. merid.* : And. Quitens.
- trachyodontea* Mitt. *op. cit.* p. 400.  
 1 vel syn. *Cortic.* — *Am. merid.* : And. Quitens.
- Uleana* C. M. in *Bull. hb. Boissier*, 1898, p. 111 et in *Ule Coll. M. brasili.* n. 1868.  
 1. *Ad ramul. arb.* — *Am. merid.* : Brasil. or.
- undulata* W. Arn. — *Neckera.*
- Wallisii* C. M. in *Fl.* 1875, n. 34-35.  
 1. — *Am. merid.* : And. Nov. Granat.

**DASYMITRIUM** Lindb. in *Oefv. af K. Sv. Vet.-Akad. Föerh.* 1864, XXI, pp. 421 et 423. — 6 spec.

- gymnostomum* (Sull. et Lesq.) Lindb. *Contrib. ad fl. crypt. As. bor. or.* p. 229 (1872).  
*Macromitrium gymnostomum* S. et L. in *Proceed. of the Amer. Acad...* 1859, p. 3, n. II; *Par. Ind.* p. 777 (1896).  
 2. *Cortic. et rupic.* — *As.* : China, Japon.; Tonkin?
- incurvum* Lindb. in *Seeman's Journ. of bot.* 1864, p. 335.  
*Macromitrium incurvum* *Par. Ind.* p. 778 (1896).  
*M. japonicum* C. M. *Syn. I*, p. 740 (1849).  
 — *As.* : China, Tonkin., Japon.
- Mac Leai* Rehm. *M. Afr. austr. n.* 508.  
*Macromitrium Mac Leai* *Par. op. cit.* p. 780.  
 — *Afr.* : Transvaal.

*Makinoi* Broth. in *Hedwig*. 1899, p. 215.

*Macromitrium Makinoi* Par. *Suppl. Ind. p.* 239 (1900).

1. Cortic. — *As.* : Japon.

*Nadeandii* Besch. in *Bull. Soc. bot. Fr.* 1898, p. 65.

2. — *Pac.* : Taïti.

*Rehmanni* C. M. — *Macromitrium serpens*.

*rupestre* (Mitt.) Lindb. *Contrib. ad fl. bryol. As. bor. or.* p. 229 (1872).

*Macromitrium rupestre* Mitt. in *Journ. of the Linn. Soc.* 1864, p. 150 ;  
*Par. Ind. p.* 786 (1896).

2. Rupic. — *As.* : Japon.

**DAWSONIA** R. Br. in *Trans. of the Linn. Soc. X*, p. 316, t. 23. — 11 spec.

? *altissima* Geh. in *Fl.* 1886, n. 23.

— *As.* : Borneo.

*appressa* Hpe. in *Linn.* 1859-60, p. 635; *F. v. Müll. Anal. draw. af Austr. M. t. IX* (1864).

Rupic. — *Pac.* : Austral. or.

*Beccarii* Broth. et Geh. in *Rev. bryol.* 1896, p. 78 et in *Bibl. bot.* 1898, p. 13, t. XII.

2. — *Pac.* : Nov. Guin. (m. Arfak.)

*gigantea* C. M. in *Hedwig*. 1897, p. 336; Geh. in *Bibl. bot. l. c. t. XI*.

2. — *Pac.* : Nov. Guin. (m. Arfak.)

*grandis* Schlieph. et Geh. in *Rev. bryol.* 1896, p. 76.

— *Pac.* : Nov. Guinea.

*intermedia* C. M. in *op. cit.* p. 335.

2. — *Pac.* : Austral. (Upper Yarra River, Nov. Vales.).

*longisetata* Hpe. in *Linn.* 1859-60, p. 634.

*D. longisetacea* F. v. Müll. *op. cit. t. IX*.

— *Pac.* : Austral. or.

*longisetacea* F. v. Müll. — *D. longisetata*.

*papuana* F. v. Müll. *mss.*; Geh. in *Rev. bryol. l. c.*

— *Pac.* : Nov. Guin.

*polytrichoides* R. Br. in *op. et l. cc.*; *Brid. Bryol. univ. II*, p. 156 (1827);  
*C. M. Syn. I*, p. 226 (1849).

*Triplocoma polytrichoides la Pyl.* in *Desv. Journ. bot.* 1813, p. 7.

2. — *Pac.* : Austral. or., Nov. Seeland.

*superba* Grev. in *Ann. and Mag. of nat. Hist.* 1847, p. 226; *C. M. op. et l. cc.*

*Polytrichum longifolium* *Bryol. eur. IV (Mon. p. 2, nom.)* ?

2. — *Pac.* : Austral., Nov. Seeland. sept. et med.,  
Tasman.

*Victoriarum* C. M. in *Hedwig*. 1897, p. 335.

2. Terr. — *Pac.* : Austral. or. (Victoria).

DECODON (1) *C. M. in sched.*

Spec. omn. sunt HYPNODONTES ejusd. nom.

DENDIA *Rob. Br. in Trans. of the N. Zeal. Inst.* 1897, p. 411.

*maritima* ej. — Pottia.

**DENDROCRYPHÆA** *Par. et W. P. Sch. Mon. Cryph. ined.* — 2 spec.

*Gorveana (Mont.) Par. et W. P. Sch. op. cit.*

*Cryphaea Gorveana Mont. in Ann. sc. nat.* 1845, p. 99, *in Gay, Hist.*

*Chil. Bot. crypt. t.* 5, f. 3 (1850) *et Syll. p.* 24 (1856).

*Dendropogon Gorveanus Jaeg. Ad. II, p.* 103 (1874-75).

*Pilotrichum Gorveanum C. M. Syn. II, p.* 175 (1851).

1. — *Am. merid.* : Chile.

*Lechleri (C. M.) Par. et W. P. Sch. op. cit.*

*Cryphaea Gorveana W. P. Sch. in Lechler Pl. Chil. n.* 474.

*Cr. Lechleri C. M. mss.*

*D. ramosissima Par. et W. P. Sch. prius.*

1. — *Am. sept.* : Guadalup. (???) *Am. merid.* : Valdiv.

*ramosissima Par. et W. P. Sch.* — *D. Lechleri.*

**Dendrobium** *C. M. in Dusen M. Camer. (sect. POROTRICH).*

**Dendroglossophyllum** *C. M. Syn. II, p.* 223 (sect. *HOOKERIE*) (1851).

DENDRO-HYPNUM *Hpe. Moossb. p.* 21 (1871).

*Beccarii* ej. — *Hypnodendron.*

*Leichardti* ej. — ? *Ptychomium.*

*praenitens* ej. — *Pterobryella.*

DENDRO-LESKEA *Hpe. in Bot. Zeit.* 1869, n. 28 ?

*Krauseana* ej. — *Porotrichum.*

**Dendroligotrichum** *C. M. Syn. I, p.* 99 (1849) *et II, p.* 559 (1851) (sect. *POLYTRICH).*

**DENDROPOGON** *W. P. Sch. in Bot. Zeit.* 1843, p. 377, t. 1. — 3 spec.

*capensis* *W. P. Sch.* — *Papillaria africana.*

*crenulatus* *Jaeg.* — *Cyptodon?*

*dentatus* *Mitt.* — *Pilotrichopsis.*

var. *filiformis* *Besch.* — *Pilotrichopsis dentata var. filiformis.*

*fasciculatus* *Jaeg.* — *Cryphaea.*

? *ferrugineus (Mitt.) Jaeg. Ad. II, p.* 188 (1874-75).

*Cryphaea ferruginea Mitt. M. Ind. or. p.* 125 (1859).

*Ster.* — *As.* : *M. Neilgherr.*

*Gorveanus* *Jaeg.* — *Dendrocryphaea.*

*helictophyllus* *Jaeg.* — *Meteorium.*

*Mülleri* *Hpe.* — *Cyptodon.*

---

(1) Nom. jam abusum : *Decodon* gen. *Lythriacearum* *Gmel. Syst. p.* 677 (1791); *DC. Prodr. III, p.* 90).

*rufescens* W. P. Sch. *op. et ll. cc.*

*Cleistostoma elegans* Br. *in litt.*

*Clidostoma rufulum* v. Mart. (*fid.* W. P. Sch.).

*Cryphaea rufescens* C. M. *in Linn.* 1844, p. 682; *Mitt. M. austr. am.* p. 416 (1869).

*Pilotrichum rufescens* C. M. *Syn. II*, p. 176 (1851).

l. Ad arbor. — *Am. sept.* : Mexic., ins. S. Thomas.

? *Schleinitzianus* (C. M.) Par. *Ind.* p. 315 (1894).

*Cryphaea* (Dendropogon) *Schleinitziana* C. M. *in Engler's Bot. Jahrb.* 1883, p. 84.

Ster? — *Pac.* : Ins. Fidji, Tonga.

*squarrulosus* Jaeg. — *Cyptodon*.

**DERMATODON** Hüb. *Muscol. germ.* p. 109 (1833).

Spec. omn. non indicatae sunt *POTTIAE* ejusd. nom.

*affinis* Hüb. — *P. Starkeana* var. *brachyodus*.

*cernuus* ej. — *Desmatodon*.

*Funchii* ej. — *P. latifolia* var. *pilifera*.

**DESMATODON** Brit. Mant. M. p. 86 (1819) et *Bryol. univ.* I, p. 523 (1826). — 16 spec.

*adustus* (Mitt.) Par. — *Tortula*.

*amblyophyllus* Mont. — *Tortula*.

*arenaceus* Sull. *M. of U. St.* p. 28 (1856) et *lc. M.* p. 45, t. 29 (1864); *Lesq. et Jam. Man.* p. III (1884).

*Barbula arenacea* Jaeg. *Ad. II*, p. 671 (1877-78).

*B. obtusifolia* Schwaegr. (*fid. Hagen in Rev. bryol.* 1891, p. 15).

*D. ohioensis* W. P. Sch. *Syn. I ed.* p. 159 (1860).

*Didymodon arenaceus* Kindb. *Eur. a. N. Am. Bryin.* p. 277 (1897).

Ps. 2. *Rupic.* — *Eur.* : Norvegia. *Am. sept.* : Ohio, Kansas, reg. centr.

*atro-virens* Jur. — *Tortula*.

*bogosicus* C. M. *in Nuov. Giorn. bot. ital.* 1872, p. 12.

2. — *Afr.* : Abyssin.

*Brebissoni* Jur. — *Dialytrichia*.

*brevicaulis* Brid. — *D. latifolius* var. *brevicaulis*.

*bulbosus* de Not. — *Pottia latifolia*.

*caespitosus* de Not. — *Pottia*.

*camptothecius* Kindb. *in Mac. Cat.* p. 48 (1892).

l. Ad rup. marit. — *Am. merid.* : Gaspesia.

*canescens* Jur. — *Tortula*.

*cernuus* (Hüb.) *Bryol. eur. fasc.* 18-20 (Vol. II), *Mon.* p. 8, t. 134 (1843); *W. P. Sch. Syn. II ed.* p. 186 (1876).

*Cynodon latifolius* Brid. *Mant. M.* p. 99 (1819).

*Cynodontium latifolium* Schwaegr. *Suppl. I, P. I*, p. 110, t. 28 (1811).

*Dermatodon cernuus* Hüb. *Musc. germ.* p. 117 (1833).

*Desmatodon inclinatus* Sendtn. *in Regensb. Denkschr. III*, p. 143 (1841).

*Didymodon latifolius* Wahlenb. *Fl. lapp.* p. 313, t. 30 (1812).

*Tortula cernua* Lindb. *M. scand.* p. 20 (1879).

*Trichostomum cernuum* Lindb. in *Oefv. af K. Vet.-Akad. Förelh.* p. 225 (1864).

*Tr. inclinatum* C. M. *Syn. I.*, p. 593 (1849).

1. In fiss. rup. et ad mur. humid. — *Eur.* : Anglia, Scand., Sudet., Salisb., Stir., Carinth., Tirol., Alp. Bavar., Helvet. *As.* : Tibet. occid. *Am. sept.* : Groenland, Anticosti, New Brunswick, reg. sept. occid., M. Rupestr., Columb. brit., Nevada.

« *var. xanthopus* Kindb. in *Mac. Cat.* p. 48 (1892).

*Terr.* — *Am. sept.* : Manitoba.

*chloronotos* Mitt. — *Crossidium*.

*confertifolius* (Mitt.) Jaeg. — *Didymodon* ?

*corticola* W. P. Sch. in *W. Sch. M. Abyss.*

— *Afr.* : Abyssin.

*cuneifolius* Jur. — *Tortula*.

*curtus* Brid. — *Ditrichum tortile* *var. pusilla*.

*dichotomus* Brid. — *Dialytrichia Brebissoni*.

*eucalypttratus* Limpr. — *D. latifolius* *var. glacialis*.

*flavicans* Bryol. eur. — *Tortula obtusifolia*.

*var. obtusifolius* Bryol. eur. — *Tortula obtusifolia* *var. brevifolia*.

*flexifolius* Hpe. — *Leptodontium*.

*fusco-mucronatus* (C. M.) Par. — *Tortula*.

Garberi Lesq. et Jam. *Man.* p. 112 (1884).

• *An Tortula* ? (*fid. Broth.*).

*Ps.* 2.

— *Am. sept.* : Florid.

*Gasilieni* Vent. — *Tortula atrovirens* *var. Gasilieni*.

*glacialis* Funck. — *D. latifolius* *var. glacialis*.

*gracilis* Anzi. — *D. latifolius* *var. brevicaulis*.

*griseus* Jur. — *Crossidium*.

*Guepini* Bryol. eur. — *Tortula*.

*Güntheri* Sendtn. — *D. cernuus*.

*inermis* Mitt. } — *Tortula Brandisii* ? (*fid. C. M.*).

                  } — *Tortula* (*fid. Jur.*).

*inclinatus* Sendtn. — *D. cernuus*.

*involutus* Mitt. — *Hyophila*.

*javanicus* Mitt. — *Tortula*.

*lanceolatus* de Not. — *Pottia*

*laterum* W. P. Sch. — *Tortula plinthobia*.

*latifolius* (Hedw.) Brid. *Bryol. univ. I.*, p. 524 (1826); *Bryol. eur. fasc.* 18-20 (*Vol. II*), *Mon.* p. 5, t. 129 (1843); *W. P. Sch. op. cit.* p. 184.

*Barbula latifolia* Kindb. *Eur. and N. Am. Bryin.* p. 252 (1897).

*D. latifolius* γ *lutescens* Brid. *op. cit.* p. 525.

*Dicranum lanceolatum* Schleich. *Cent. IV pl. helvet. n.* 12.

*D. latifolium* Hedw. *Descr. I.*, p. 89, t. 33 (1787); *Drumm. M. bor. am. I ed. n.* 109 (*fid. Kindb.*).

*Didymodon apiculatus* *var. β* W. Arn. in *Mem. Soc. d'hist. nat. Paris, II.* p. 283 (1825).

*D. pilifer* Wahlenb. *Fl. suec. I ed. II.* p. 754 (1826).

*Mnium latifolium* Gmel. in *L. Syst. nat. XIII ed., II, P. II.* p. 1328 (1791).

*Swartzia pilifera* Brid. in *Schrad. Journ. III, P. II.* p. 289 (1801).

*Tortula latifolia* Lindb. *M. scand. p.* 20 (1879).

*Trichostomum Hoppeanum* Schultz in *Syll. II.* p. 40 (1828).



*Tr. latifolium* Lindb. in *Oefv. af K. Vet.-Akad. Foerh.* p. 224 (1864).  
*Tr. piliferum* Hüb. *Musc. germ.* p. 293 excl. syn. (1833).

1. Ad terr. humid. usq. ad niv. aetern. — *Eur.* : Spitzberg., reg. alp. et subalp.; desideratur in Britann. *As.* : Kamschatka, Talysch, Alatau, Kashmir, Tchitral, Panjab, Sikkim, Caucas. *Am. sept.* : Groenland., Labrador, reg. sept. occid., M. Rupestr., Columb. brit., Washington, Oregon, Nevada, Colorado, Calif.

« var.  $\beta$  *brevicaulis* W. P. Sch. *op. cit.* p. 185.

*Barbula brevicaulis* (ut subspec.) Kindb. *Eur. and N. Am. Bryin.* p. 252 (1897).

*D. brevicaulis* Brid. *Mant. M.* p. 87 excl. syn. (1819) et *Bryol. univ. I.* p. 525 pp. (1826).

*D. gracilis* Anzi *Enum.* p. 10 (1875) *fid. Limpr.*

*D. Güntheri* Sendtn. in *sched.*; *C. M. Syn. I.* p. 588 (1849).

*D. latifolius* var. *piliferus* Rab. *Deutsch. Kryptfl. II*, 3, p. 100 excl. syn. (1848).

*Dicranum latifolium*  $\beta$  *piliferum* H. et T. *Musc. brit.* p. 22 (1818).

Ut et cum typ.

« var. *calyptratus* Kaur. mss.; Bryhn in *D. K. N. Vid. Selsk. Skr.* 1899, n. 3, p. 25.

— *Eur.* : Norveg.

« var. *glacialis* (Funck) W. P. Sch. *I ed.* p. 157 et *II ed. l. c.*

*Barbula glacialis* (ut subsp.) Kindb. *op. et l. cc.*

*D. eucalypttratus* Limpr. *Laubm. Deutschl...* I, p. 650 (1888).

*D. glacialis* Funck mss. *Crypt. Gew. n.* 691; *Brid. Bryol. univ. I.* p. 822 p.p. (1826).

*D. latifolius*  $\beta$  *muticus* Brid. *Sp. M. I.* p. 207 (1806); *Bryol. eur. l. c.* t. 130.

*D. spathulatus* Hartm. in *Fr. Summ. Veg.* p. 94.

*Didymodon apiculatus*  $\alpha$  W. Arn. in *Mem. Soc. d'hist. nat. Paris, II*, p. 283 (1825).

*D. pilifer*  $\beta$  *spathulatus* Wahlenb. *Fl. suec. I ed. II*, p. 755 (1826).

*D. spathulatus* de Not. in *Mem. Acad. Torin. XXXVIII*, p. 233 (1836).

*D. Theobaldi* Pfeff. in *Jahresb. Naturf. Ges. Graubünd.* 1868, p. 83.

*Tortula eucalypttrata* Lindb. in *Meddel. 13 Hefst.* p. 252 (1886) et in *Not.* 1886, p. 100.

*T. latifolia*  $\beta$  *mutica* ej. *M. scand.* p. 20 (1879).

*Trichostomum latifolium*  $\beta$  *muticum* C. M. l. c.

Ut typ. — *Eur.* : Cum typ. *As.* : Kashmir, Tchitral, Caucas. *Am. sept.* : Groenland., M. Rupestr.

var. *lutescens* Brid. — *D. latifolius*.

var. *muticus* ej. — *D. latifolius* var. *glacialis*.

var. *piliferus* Rab. — *D. latifolius* var. *brevicaulis*.

Laureri (Schultz) *Bryol. eur. fasc.* 18-20 (*Vol. II*) *Mon.* p. 9, t. 135 (1843); W. P. Sch. *op. cit.* p. 187.

*Didymodon Blyttii* Hartm. *Skand. Fl. IV ed.* p. 381 (1843).

*D. nitens* Liebm. in *Fl. dan. Suppl. I*, p. 12, t. 50 (1853).

*Tortula bryoides* Hook. in *Drumm. M. bor. am. I ed. n.* 135 (1828).

*T. Laureri* Lindb. in *Oefv. af K. Vet.-Akad. Foerh.* p. 243 (1864).

*Trichostomum latifolium*  $\beta$  *Laureri* Hüb. *Musc. germ.* p. 292 (1833).

*Tr. Laureri* Schultz in *Fl.* 1827, *P. I.* p. 163.

1. Ad terr. humos. — *Eur.* : Rar. : Spitzberg., alp. Norveg., Salisb., Stir., Carinth., Bavar., Algov., Tirol., Helvet. *As.* : Tian Shan. *Am. sept.* : Groenland., M. Rupestr., ins. Vancouver, Colorado.

*Laureri* Wils. — *D. Thomsonii*.

*leucostomus* Berggr. — *D. suberectus*.

*limbatus* Mitt. — *Tortula sublimbata*.

*longirostris* (Griff.) Mitt. *M. Ind. or.* p. 37 (1859).

*Hymenostylium longirostre* Griff. *Not. p.* 395 *et Ic. pl. asiat. II, t.* 95. f. 3 (1849).

Lignic. et ad rup. humid. — *As.* : Khasia.

*marginatus* Mitt. — *Tortula*.

*micrangius* (C. M.) Par. — *Ptychomitrium sinense*.

*montenegrinus* (Br. et Ssycz.) Par. — *Tortula*.

*mucronifolius* Mitt. — *Tortula*.

*neo-mexicanus* Sull. et Lesq. — *Tortula*.

*nervosus* Bryol. eur. — *Tortula atro-virens*.

var. *Gasilieni* Husn. — *Tortula atro-virens* var. *Gasilieni*.

*nitens* Liebm. — *D. Laureri*.

*obliquus* Bryol. eur. — *D. suberectus*.

var. *apiculatus* { Lindb. — *D. suberectus*.

var. *muticus* {

*obtusifolius* W. P. Sch. — *Tortula*.

*ohioënsis* ej. — *D. arenaceus*.

*plinthobius* Sull. — *Tortula*.

*Porteri* Jam. et Aust. — *Tortula*.

*recurvatus* Mitt. — *Tortula*.

*recurvus* (Griff.) Mitt. — *Didymodon*.

*reflexidens* (Hpe.) Jaeg. *Ad. I, p.* 257 (1871-72).

*Trichostomum reflexidens* Hpe. in *Linn.* 1859-60, p. 625.

— *Pac.* : Austral. or.

*revolvens* Husn. — *Barbula*.

*rigidus* Mitt. — *Aloina*.

*ruber* Mitt. — *Tortula*.

*rupestris* Funck. — *Didymodon rigidulus* var. *densus*.

*Sartorii* (C. M.) Par. *Ind. p.* 319 (1895).

*Barbula Sartorii* Jaeg. *Ad. II, p.* 671 (1877-78).

*Trichostomum Sartorii* C. M. in *Linn.* 1874, p. 637.

— *Am. sept.* : Mexic.

*Schimperi* (Mont.) Jaeg. — *Didymodon*.

*spathulatus* { Hartm. — *D. latifolius* var. *glacialis*.

Mitt. — *Hyophila commutata*.

*Starkei* de Not. — *Pottia Starkeana*.

*suberectus* (Hook.) Limpr. *Laubm. Deutschl. .... I, p.* 651 (1888).

*Barbula leucostoma* R. Br. (fid. Berggr. in *K. Sv. Vet. Akad. Foerh. XIII, n.* 7, p. 46); *Par. Ind. p.* 81 (1894).

*B. suberecta* Kindb. *Eur. and N. Am. Bryin. p.* 253 (1897).

*D. leucostomus* Berggr. *M. Spetsb. n.* 34.

*D. obliquus* Bryol. eur. fasc. 18-20 (*Vol. II*), *Mon. p.* 10, t. 136 (1843)

W. P. Sch. *op. cit. p.* 187.

*D. obliquus* {  $\beta$  *apiculatus* { Lindb. in *Oefv. af K. Sv. Vet.-Akad. Foerh. XXIII, p.* 553.

$\gamma$  *muticus* {

*D. Tortula Br. in sched. ; C. M. Syn. I, p.* 594 (1849).

*Tortula suberecta* Hook. in *Drum. M. bor. am. I ed. n.* 145 (1828).

*Trichostomum obliquum* C. M. *op. cit. p.* 594 *et II, p.* 629 (1851).

*Zygotrichia leucostoma* Brid. *Bryol. univ. I, pp.* 221 *et* 821 (1826).

- l. Ad terr. humos. — *Eur.* : Rariss. : Alp. Norveg., Salisb., Stir., Carinth., Bavar. et Algov., Tirol., Helvet. *As.* : Penins. Tschutschica. *Am. sept.* : Ins. Sabine, fret. Scoresby, ins. Melville, M. Rupestr.

*sublimbatus* Mitt. — *Tortula*.

*subtorquescens* C. M. et Kindb. — *Tortula*.

*subulatus* Jur — *Tortula*.

var. *angustatus* ej. — *Tortula subulata* var. *angustata*.

*systylius* *Bryol. eur. fasc.* 31 (*Vol. II*), *Mon. Suppl. t.* 131 (1846); *W. P. Sch. op. cit. p.* 185.

*Anacalypta latifolia*  $\beta$  *pilifera* *Hook. in Drumm. M. bor. am. I ed. n.* 70 (1828).

*Tortula systylia* *Lindb. M. scand. p.* 20 (1879).

*Trichostomum systylium* *C. M. Syn. I, p.* 589 (1849).

- l. In terr. humos. — *Eur.* : Norveg., alp. Salisb., Stir., Carinth., Bavar. et Algov., Tirol., Helvet. *As.* : Alatau, Caucas. *Am. sept.* : Groenland., Manitoba, M. Rupestr., Colorado, Calif.

*systylioides* *Ren. et Card. in Bot. Gaz.* 1900, p. 16, t. II.

*An Pottia ? (fd. Broth.).*

— *Am. sept.* : Labrador.

*Thomsonii* (*C. M.*) *Jaeg. Ad. I, p.* 251 (1871-72).

*D. Laureri* *Wils. in hb. Ind. or. n.* 273; *Mitt. M. Ind. or. p.* 37 (1859).

*Trichostomum Thomsonii* *C. M. in Bot. Zeit.* 1854, p. 359.

- l. — *As.* : Himalaya, Tibet, occid.

*tonkinensis* *Besch. in Bull. Soc. bot. Fr.* 1893, p. 80.

*An Tortula ? (fd. Broth.).*

Ad rhizom. *Filicum* — *As.* : Tonkin.

*Tortula* *C. M.* — *D. suberectus*.

*Wallichii* *Mitt.* — *Didymodon*.

DESMOSTOMUM *Hsch. in Linn.* 1841, p. 116.

*Absque nom. spec.* — *Entosthodon Bergianus*.













